

BANG-YEN CHEN

Cohomology of CR-submanifolds

Annales de la faculté des sciences de Toulouse 5^e série, tome 3, n° 2 (1981), p. 167-172

http://www.numdam.org/item?id=AFST_1981_5_3_2_167_0

© Université Paul Sabatier, 1981, tous droits réservés.

L'accès aux archives de la revue « Annales de la faculté des sciences de Toulouse » (<http://picard.ups-tlse.fr/~annales/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

COHOMOLOGY OF CR-SUBMANIFOLDS

Bang-Yen Chen ⁽¹⁾

(1) *Department of Mathematics, Michigan State University, East Lansing, Michigan 48824 - USA.*

Résumé : Nous introduisons canoniquement une classe de cohomologie de Rham pour une CR-sous-variété compacte d'une variété kaehlérienne. Cette classe de cohomologie est utilisée pour montrer que si un certain groupe de cohomologie de dimension paire d'une CR-sous-variété, N est trivial, alors, soit la distribution holomorphe de N n'est pas intégrable, soit la distribution totalement réelle de N n'est pas minimale.

Summary : We introduce a canonical de Rham cohomology class for a closed CR-submanifold in a Kaehler manifold. This cohomology class is used to prove that if some even-dimensional cohomology group of a CR-submanifold N is trivial, then either the holomorphic distribution of N is not integrable or the totally real distribution of N is not minimal.

1. - INTRODUCTION

Let \tilde{M} be a Kaehler manifold with complex structure J and N a Riemannian manifold isometrically immersed in \tilde{M} . Let \mathcal{D}_x be the maximal holomorphic subspace of the tangent space $T_x N$, i.e., $\mathcal{D}_x = T_x N \cap J(T_x N)$. If the dimension of \mathcal{D}_x is constant along N , then \mathcal{D}_x defines a differentiable distribution \mathcal{D} , called the *holomorphic distribution* of N . A submanifold N in \tilde{M} is called a *CR-submanifold* [1,2] if there exists on N a holomorphic distribution \mathcal{D} such that its orthogonal complement \mathcal{D}^\perp is a distribution satisfying $J\mathcal{D}_x^\perp \subset T_x^\perp N$, $x \in N$. \mathcal{D}^\perp is called the *totally real distribution* of N .

Let \mathcal{H} be a differentiable distribution on a Riemannian manifold N with Levi-Civita connection ∇ . We put

$$(1.1) \quad \overset{\circ}{\sigma}(X, Y) = (\nabla_X Y)^\perp$$

for any vector fields X, Y in \mathcal{H} , where $(\nabla_X Y)^\perp$ denotes the component of $\nabla_X Y$ in the orthogonal complementary distribution \mathcal{H}^\perp in N . Let X_1, \dots, X_r be an orthonormal basis of \mathcal{H} , $r = \dim_{\mathbb{R}} \mathcal{H}$. If we put

$$(1.2) \quad \overset{\circ}{H} = \frac{1}{r} \sum_{i=1}^r \overset{\circ}{\sigma}(X_i, X_i).$$

Then $\overset{\circ}{H}$ is a well-defined \mathcal{H}^\perp -valued vector field on N (up to sign), called the *mean-curvature vector* of \mathcal{H} . A distribution \mathcal{H} on N is called *minimal* if the mean-curvature vector $\overset{\circ}{H}$ of \mathcal{H} vanishes identically.

The main purpose of this paper is to introduce a canonical cohomology class and use it to prove the following.

THEOREM 1. *Let N be a closed CR-submanifold of a Kaehler manifold \tilde{M} . If $H^{2k}(N; \mathbb{R}) = 0$ for some $k \leq \dim_{\mathbb{C}} \mathcal{D}$, then either \mathcal{D} is not integrable or \mathcal{D}^\perp is not minimal.*

2. - THE CANONICAL CLASS OF CR-SUBMANIFOLDS

Let M be a Kaehler manifold and N a CR-submanifold of \tilde{M} . We denote by $\langle \cdot, \cdot \rangle$ the metric tensor of \tilde{M} as well as that induced on N . Let ∇ and $\tilde{\nabla}$ be the covariant differentiations on N and \tilde{M} , respectively. The Gauss and Weingarten formulas are given respectively by

$$(2.1) \quad \tilde{\nabla}_X Y = \nabla_X Y + \sigma(X, Y),$$

$$(2.2) \quad \tilde{\nabla}_X \xi = -A_\xi X + D_X \xi$$

for any vector fields X, Y tangent to N and any vector field ξ normal to N . The second fundamental form σ and the second fundamental tensor A_ξ satisfy $\langle A_\xi X, Y \rangle = \langle \sigma(X, Y), \xi \rangle$. We recall the following.

PROPOSITION 2 [2]. *The totally real distribution \mathcal{D}^\perp of any CR-submanifold in any Kaehler manifold is integrable.*

For a CR-submanifold N of a Kaehler manifold M , we choose an orthogonal local

frame $e_1, \dots, e_h, Je_1, \dots, Je_h$ of \mathcal{D} . Let $\omega^1, \dots, \omega^h, \omega^{h+1}, \dots, \omega^{2h}$ be the $2h$ 1-forms on N satisfying

$$(2.3) \quad \omega^j(Z) = 0, \omega^i(e_j) = \delta_{ij}, i, j = 1, \dots, 2h$$

for any $Z \in \mathcal{D}^\perp$ where $e_{h+j} = Je_j$. Then

$$(2.4) \quad \omega = \omega^1 \wedge \dots \wedge \omega^{2h}$$

defines a $2h$ -form on N . This form is a well-defined global $2h$ -form on N because \mathcal{D} is orientable. We give the following.

THEOREM 3. For any closed CR-submanifold N of a Kaehler manifold \tilde{M} , the $2h$ -form ω is closed which defines a canonical deRham cohomology class given by

$$(2.5) \quad c(N) = [\omega] \in H^{2h}(N; \mathbb{R}), h = \dim_{\mathbb{C}} \mathcal{D} .$$

Moreover, this cohomology class is nontrivial if \mathcal{D} is integrable and \mathcal{D}^\perp is minimal.

Proof. First we give the following.

LEMMA 4. If N is a CR-submanifold of a Kaehler manifold \tilde{M} , then the holomorphic distribution \mathcal{D} is minimal.

Let X and Z be vector fields in \mathcal{D} and \mathcal{D}^\perp , respectively. Then we have

$$(2.6) \quad \langle Z, \nabla_X X \rangle = \langle JZ, \tilde{\nabla}_X JX \rangle = -\langle \tilde{\nabla}_X JZ, JX \rangle = \langle A_{JZ} X, JX \rangle .$$

Thus we find

$$(2.7) \quad \langle Z, \nabla_{JX} JX \rangle = -\langle A_{JZ} JX, X \rangle = -\langle A_{JZ} X, JX \rangle .$$

Combining (2.6) and (2.7) we get $\langle \nabla_X X + \nabla_{JX} JX, Z \rangle = 0$ from which we conclude that the holomorphic distribution \mathcal{D} is minimal. This proves the lemma.

From (2.4) we have

$$(2.8) \quad d\omega = \sum_{i=1}^{2h} (-1)^i \omega^1 \wedge \dots \wedge d\omega^i \wedge \dots \wedge \omega^{2h} .$$

It is clear from (2.3) and (2.8) that $d\omega = 0$ if and only if

$$(2.9) \quad d\omega(Z_1, Z_2, X_1, \dots, X_{2h-1}) = 0$$

$$(2.10) \quad d\omega(Z_1, X_1, \dots, X_{2h}) = 0$$

for any vectors $Z_1, Z_2 \in \mathcal{D}^\perp$ and $X_1, \dots, X_{2h-1} \in \mathcal{D}$. However, it follows from straight-forward computation that (2.9) holds when and only when \mathcal{D}^\perp is integrable and (2.10) holds when and only when \mathcal{D} is minimal. But for a CR-submanifold in a Kaehler manifold these two conditions hold automatically (Proposition 2 and Lemma 4). Therefore, the $2h$ -form ω is closed. Consequently, ω defines a deRham cohomology class $c(N)$ given by (2.5).

Let $e_{2h+1}, \dots, e_{2h+p}$ be an orthonormal local frame of \mathcal{D}^\perp and let $\omega^{2h+1}, \dots, \omega^{2h+p}$ be the p 1-forms on N satisfying $\omega^\alpha(X) = 0$ and $\omega^\alpha(e_\beta) = 0$ for any X in \mathcal{D} , where $\alpha, \beta = 2h+1, \dots, 2h+p$. Then by a similar argument for ω , we may conclude that if \mathcal{D} is integrable and \mathcal{D}^\perp is minimal, then the p -form $\omega^\perp = \omega^{2h+1} \wedge \dots \wedge \omega^{2h+p}$ is closed. Thus, the $2h$ -form ω is coclosed, i.e., $\delta\omega = 0$. Since N is a closed submanifold, ω is harmonic. Because ω is nontrivial, the cohomology class $[\omega]$ represented by ω is nontrivial in $H^{2h}(N; \mathbb{R})$. This proves the Theorem.

2. - PROOF OF THEOREM 1

Let N be a closed CR-submanifold of a complex m -dimensional Kaehler manifold M . Let $h = \dim_{\mathbb{C}} \mathcal{D}$ and $p = \dim_{\mathbb{R}} \mathcal{D}^\perp$. We choose an orthonormal local frame

$$e_1, \dots, e_h, e_{h+1}, \dots, e_{h+p}, e_{h+p+1}, \dots, e_m, J e_1, \dots, J e_m$$

in \tilde{M} in such a way that, restricted to N , $e_1, \dots, e_h, J e_1, \dots, J e_h$ are in \mathcal{D} and e_{h+1}, \dots, e_{h+p} are in \mathcal{D}^\perp . We denote by $\omega^1, \dots, \omega^m, \omega^{1*}, \dots, \omega^{m*}$, the dual frame of $e_1, \dots, e_m, J e_1, \dots, J e_m$. We put

$$\theta^A = \omega^A + \sqrt{-1} \omega^{A*}, \bar{\theta}^A = \omega^A - \sqrt{-1} \omega^{A*}, A = 1, \dots, m.$$

Then, restrict θ^A 's and $\bar{\theta}^A$'s to N , we have

$$(3.1) \quad \theta^\alpha = \bar{\theta}^\alpha = \omega^\alpha \quad \text{for } \alpha = h+1, \dots, h+p$$

$$\theta^r = \bar{\theta}^r = 0 \quad \text{for } r = h+p+1, \dots, m.$$

The Kaehler form $\tilde{\Omega}$ of \tilde{M} is a closed 2-form on \tilde{M} given by

$$(3.2) \quad \tilde{\Omega} = \frac{\sqrt{-1}}{2} \sum_A \theta^A \wedge \bar{\theta}^A.$$

Let $\Omega = i^*\tilde{\Omega}$ be the 2-form on N induced from $\tilde{\Omega}$ via the immersion $i : N \rightarrow \tilde{M}$. Then, (3.1) and (3.2) give

$$(3.3) \quad \Omega = \frac{\sqrt{-1}}{2} \sum_{i=1}^h \theta^i \wedge \bar{\theta}^i.$$

It is clear that Ω is a closed 2-form on N and it defines a cohomology class $[\Omega]$ in $H^2(N ; \mathbb{R})$. (2.4) and (3.3) imply that the canonical class $c(N)$ and the class $[\Omega]$ satisfy

$$(3.4) \quad [\Omega]^h = (-1)^h (h!) c(N).$$

If \mathcal{D} is integrable and \mathcal{D}^\perp is minimal, then Theorem 3 and (3.4) imply that $H^{2k}(N ; \mathbb{R}) \neq 0$ for $k = 1, 2, \dots, h$. (Q.E.D).

Because every hypersurface of a Kaehler manifold is a CR-hypersurface, Theorem 1 implies the following.

COROLLARY 5. *Let N be a $(2m-1)$ -dimensional closed manifold with $H^{2k}(N ; \mathbb{R}) = 0$ for some $k < m$. Then any immersion from N into a (complex) m -dimensional Kaehler manifold \tilde{M} is a CR-hypersurface such that either its holomorphic distribution is not integrable or its totally real distribution is not minimal.*

Remark. CR-products of a Kaehler manifold are examples of CR-submanifold whose holomorphic distributions are integrable and whose totally real distributions are minimal. Therefore, the assumption on cohomology groups are necessary for Theorem 1.

REFERENCES

- [1] A. BEJANCU. «*CR-submanifolds of Kaehler manifolds, I*». Proc. Amer. Math. Soc. 69 (1978), 134-142.
- [2] B.Y. CHEN. «*On CR-submanifolds of a Kaehler manifold, I*». J. Differential Geometry (to appear).

(Manuscrit reçu le 18 mai 1981)