

# ANNALES DE LA FACULTÉ DES SCIENCES DE TOULOUSE Mathématiques

LUCIEN GUILLOU

*On the structure of homeomorphisms of the open annulus*

Tome XX, n° 2 (2011), p. 367-378.

[http://afst.cedram.org/item?id=AFST\\_2011\\_6\\_20\\_2\\_367\\_0](http://afst.cedram.org/item?id=AFST_2011_6_20_2_367_0)

© Université Paul Sabatier, Toulouse, 2011, tous droits réservés.

L'accès aux articles de la revue « Annales de la faculté des sciences de Toulouse Mathématiques » (<http://afst.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://afst.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du  
Centre de diffusion des revues académiques de mathématiques  
<http://www.cedram.org/>

## On the structure of homeomorphisms of the open annulus

LUCIEN GUILLOU<sup>(1)</sup>

*Dedicated to Jose Maria Montesinos on the occasion  
of his 65th birthday*

---

**ABSTRACT.** — Let  $h$  be a without fixed point lift to the plane of a homeomorphism of the open annulus isotopic to the identity and without wandering point. We show that  $h$  admits a  $h$ -invariant dense open set  $O$  on which it is conjugate to a translation and we study the action of  $h$  on the compactly connected components of the closed and without interior set  $\mathbf{R}^2 \setminus O$ .

**RÉSUMÉ.** — Soit  $h$  un relevé au plan sans point fixe d'un homéomorphisme de l'anneau ouvert isotope à l'identité et sans point errant. Nous montrons que  $h$  admet un ouvert dense et  $h$ -invariant  $O$  sur lequel il est conjugué à une translation et nous étudions l'action de  $h$  sur les composantes compactement connexes du fermé d'intérieur vide  $\mathbf{R}^2 \setminus O$ .

---

### 1. Introduction

**1.1.** In the paper [BCL] the authors consider homeomorphisms  $H$  of the open annulus  $S^1 \times \mathbf{R}$  isotopic to the identity and preserving the Lebesgue measure. Given such a homeomorphism and a lift  $h : \mathbf{R}^2 \rightarrow \mathbf{R}^2$  to the universal cover they show (in their proposition 3.1) that if the closure of the rotation set of  $h$  is contained in  $]0, +\infty[$ , then  $h$  is conjugate to a translation. (Here the rotation set refers to a definition, adapted to this non compact

---

(\*) Reçu le 02/07/2010, accepté le 30/09/2010

<sup>(1)</sup> Université Grenoble 1, Institut Fourier B.P. 74, Saint-Martin-d'Hères 38402 France.  
Lucien.Guillou@ujf-grenoble.fr

situation, proposed by Le Calvez [LC] and using only recurrent points of  $H$  in its construction).

They remark that this statement is sharp, and give an example of a measure preserving homeomorphism  $H$  of  $S^1 \times \mathbf{R}$  isotopic to the identity, such that, for some lift  $h$  of  $H$ , the rotation set of  $h$  is included in  $]0, +\infty[$ , but  $h$  is not conjugate to a translation (see 1.2 below).

In the present note we wish to investigate the structure of such homeomorphisms. More generally, we will consider a homeomorphism  $H$  of the annulus  $S^1 \times \mathbf{R}$  isotopic to the identity, without wandering point which admits a lift  $h$  to  $\mathbf{R}^2$  without fixed point. We will show that some of the features of example 1.2 are indeed preserved in that general situation.

We will prove:

*A) There exists an  $h$ -invariant dense open set homeomorphic to  $\mathbf{R}^2$ ,  $O \subset \mathbf{R}^2$ , such that  $h$  restricted to  $O$  is conjugate to a translation. (See paragraph 2).*

*B) Let  $W = \mathbf{R}^2 \setminus O$  which is a closed subset with no interior in  $\mathbf{R}^2$ . We have:*

*- B1) No **closed** compactly connected component (cf. 3.1 below) of  $W$  is invariant under  $h$ . (Cf. Prop. 3.4).*

*- B2) We now assume that the compactly connected components of  $W$  are **closed**. Then for every such component  $C$ , if  $X = \liminf h^n(C)$ ,*

*either  $X$  is empty (that is  $h^n(C) \rightarrow \infty$ , meaning that, for every compact  $K \subset \mathbf{R}^2$  there exists an integer  $n(K)$  such that  $h^n(C) \cap K = \emptyset$  for  $n \geq n(K)$ )*

*or it is not empty and no point of  $X$  is accessible from  $\mathbf{R}^2 \setminus \overline{\bigcup_{n \in \mathbf{Z}} h^n(C)}$ . (Cf. Prop. 3.10).*

## 1.2. The Le Roux example [BCL, Appendix A]:

We will describe the lift  $h$  of this example to  $\mathbf{R}^2$ . Let  $I_k$  be the vertical segment  $\{(\frac{1}{2k}, y) | y \geq |k|\}$  and  $A$  be  $\bigcup_{k \in \mathbf{Z} \setminus \{0\}} I_k$  and let  $W = \bigcup_{n \in \mathbf{Z}} T^n(A)$  where  $T(x, y) = (x + 1, y)$ . Then  $\mathbf{R}^2 \setminus W$  is homeomorphic to  $\mathbf{R}^2$  and can be foliated by lines equivariantly with respect to  $T$ . The homeomorphism  $h$  is chosen to act equivariantly, without fixed point, preserving each line of the foliation and satisfying  $h(I_k) = I_{k-1}$  for  $k \neq 0, 1$ . On each leaf of

the foliation,  $h$  is equivariantly conjugate to a translation hence  $h$  preserves measures without atoms and charging the open sets. On  $S^1 \times \mathbf{R}$  seen as  $S^2$  minus the two poles,  $H$  preserves such a measure which is finite. That measure is nothing but the Lebesgue measure up to conjugation thanks to a classical result of Oxtoby and Ulam.

To see that  $h$  is not conjugate to a translation notice that the compact segment going from  $x_0 = (-\frac{1}{2}, 1)$  to its translate  $T(x_0) = (\frac{1}{2}, 1)$  has to meet all its images by all iterates of  $h$  since  $W$  is  $h$ -invariant.

We owe to P. Le Calvez the remark that this example can also be described without any reference to the Oxtoby-Ulam theorem. Consider the part of the phase space (which is homeomorphic to  $S^1 \times \mathbf{R}$ ) of the free undamped pendulum above the upper separatrix: it is homeomorphic to  $S^1 \times [0, +\infty[$ . We now focus on the time 1 of the corresponding autonomous hamiltonian and on an orbit of this diffeomorphism on the separatrix. Folding each complementary interval of this orbit on the separatrix and identifying all points of the orbit and the equilibrium point of the separatrix to a single point, we get an example conjugate to the preceding one after deleting that single point.

**Acknowledgement.** — Some arguments of this paper can be traced back to an old article of T. Homma and H. Kinoshita [HK], which makes for a hard reading but a rewarding one. Many thanks to Patrice Le Calvez and Alexis Marin for careful readings of a first version of this paper.

## 2. Brouwer homeomorphisms

Homeomorphisms of the plane preserving orientation and without fixed point are called **Brouwer homeomorphisms** (see [G1] for more on these). These homeomorphisms have only wandering points and more generally satisfy the following particular version of Franks' lemma (in a reformulation due to Le Roux [LR1, Lemma 7]). Recall first that a subset  $A$  of  $\mathbf{R}^2$  is **free** if  $h(A) \cap A = \emptyset$ .

**LEMMA 2.1.** — *Let  $U$  and  $V$  be two free connected open sets. Then the subset of integers such that  $h^n(U) \cap V \neq \emptyset$  is an interval of  $\mathbf{Z}$ .*

*Proof.* — The usual formulation of this lemma concerns the case where  $U$  and  $V$  are open discs. To prove the present lemma from this case, suppose there exists  $k < n < m$  such that  $h^k(U) \cap V \neq \emptyset$ ,  $h^n(U) \cap V = \emptyset$  and  $h^m(U) \cap V \neq \emptyset$ . Let  $u_1 \in U$  such that  $v_1 = h^k(u_1) \in V$  and  $u_2 \in U$  such that  $v_2 = h^m(u_2) \in V$  and let  $D$  and  $D'$  be discs in  $U$  and  $V$  respectively

such that  $u_1, u_2 \in D$  and  $v_1, v_2 \in D'$ . Then  $h^k(D) \cap D' \neq \emptyset, h^n(D) \cap D' = \emptyset$  and  $h^m(D) \cap D' \neq \emptyset$  in contradiction to Franks' lemma.  $\square$

A **Brouwer line** for a Brouwer homeomorphism  $h$  is a properly embedded free line  $l$  such that  $l$  separates  $h^{-1}(l)$  and  $h(l)$ . We will start with the following result from [G2].

**THEOREM 2.2.** — *Let  $H : S^1 \times \mathbf{R} \rightarrow S^1 \times \mathbf{R}$  be a homeomorphism isotopic to the identity such that :*

- $H$  admits a fixed point free lift  $h : \mathbf{R}^2 \rightarrow \mathbf{R}^2$ .
- $H$  does not have any wandering point.

*Then there exists a properly embedded line in  $S^1 \times \mathbf{R}$  joining one end of the annulus to the other which lifts in  $\mathbf{R}^2$  to a Brouwer line.*

Notice that such a Brouwer line projects properly and onto on  $\{0\} \times \mathbf{R}$  (and also, a properly embedded line in  $\mathbf{R}^2$  which projects properly and onto on  $\{0\} \times \mathbf{R}$  is a Brouwer line if it is free, that is, the requirement that  $l$  separates  $h^{-1}(l)$  and  $h(l)$  is automatically satisfied).

Given any Brouwer line  $l$ , if we let  $U$  be the open region between  $l$  and  $h(l)$ , then the set  $O = \bigcup_{n \in \mathbf{Z}} h^n(\text{Cl}U)$  is homeomorphic to  $\mathbf{R}^2$  and the restriction of  $h$  to  $O$  is conjugate to a translation. Therefore to prove statement A of the introduction, it is enough to prove that if the Brouwer homeomorphism  $h$  is a lift of a homeomorphism  $H$  of the open annulus without wandering point, then  $\mathbf{R}^2 \setminus O$  has no interior for a convenient choice of Brouwer line  $l$ . To this end, we choose a Brouwer line  $l$  as given by Theorem 2.2 that we orient so that  $l$  induces by projection the usual orientation on  $\{0\} \times \mathbf{R}$ . The following Lemma is then enough to conclude the proof of statement A (this lemma is an extension of the lemma in Winkelkemper [W]).

**LEMMA 2.3.** — *Let  $B_n$  (resp.  $B'_n$ ) be the component of  $\mathbf{R}^2 \setminus h^n(l)$  to the right (resp. to the left) of  $h^n(l)$ . Then the closed  $h$ -invariant set  $W = \bigcap_{n=-\infty}^{+\infty} B_n$  (resp.  $W' = \bigcap_{n=-\infty}^{+\infty} B'_n$ ) has no interior.*

*Proof.* — Exchanging  $h$  and  $h^{-1}$  if necessary, we can suppose  $h(l)$  on the right of  $l$ . Suppose  $U \subset W$  is an open subset which we can choose small enough to be free and projecting homeomorphically on  $S^1 \times \mathbf{R}$ ; since  $U \subset W$ ,  $h^{-n}(U)$  lies on the right of  $l$  for all  $n \geq 0$ . Given the properties of  $l$ , there is a  $m > 0$  such that  $U$  lies on the left of  $T^m(l)$ , then  $h^{-n}(U)$  lies on

the left of  $h^{-n}(T^m(l)) = T^m(h^{-n}(l))$  which is on the left of  $T^m(l)$  for  $n > 0$ . So that all  $h^{-n}(U)$ ,  $n \geq 0$ , lie on the left of  $T^m(l)$  and on the right of  $l$ . There are only a finite number of translates of  $U$  between  $l$  and  $T^m(l)$ , say  $U = U_1, U_2, \dots, U_k$  and each one is wandering. Since  $H$  has no wandering point on  $S^1 \times \mathbf{R}$ , there exists  $n_1 > 0$  such that  $h^{-n_1}(U_1)$  meets some  $U_i$ , say  $U_{j(1)}$ . Let  $V_1 = h^{-n_1}(U_1) \cap U_{j(1)}$ . There exists also  $n_2 > 0$  such that  $h^{-n_2}(V_1)$  meets one of its translates  $V_{j(2)} \subset U_{i_2}$ . Let  $V_2 = h^{-n_2}(V_1) \cap V_{j(2)}$ . Continuing in that way we find a sequence  $V_1, V_2, \dots$  of non empty sets each  $V_i$  being contained in some  $U_{j(i)}$ ,  $1 \leq j(i) \leq k$ . We must have  $j(i) = j(i')$  for some  $i$  and  $i'$ ,  $i < i'$ . Then, since  $V_{i'} \subset h^{-p}(V_i)$  for  $p = n_{i+1} + \dots + n_{i'}$ , we have  $U_{j(i')} \cap h^{-p}(U_{j(i)}) \neq \emptyset$  contradicting the freeness of  $U_{j(i)}$ .  $\square$

### 3. Compactly connected components

In this paragraph we consider any Brouwer homeomorphism  $h$  and an associated oriented Brouwer line  $l$  such that  $W = \bigcap_{n=-\infty}^{+\infty} B_n$  and  $W' = \bigcap_{n=-\infty}^{+\infty} B'_n$  have no interior (where as above  $B_n$  (resp.  $B'_n$ ) is the component of  $\mathbf{R}^2 \setminus h^n(l)$  to the right (resp. to the left) of  $h^n(l)$ ).

Notice that the sets  $W$  and  $W'$  are disjoint, that the invariant set  $O = \mathbf{R}^2 \setminus (W \cup W')$  is homeomorphic to  $\mathbf{R}^2$  and that on this set  $h$  is conjugate to a translation. Similar considerations can be applied to each one of  $W$  and  $W'$  and we will only describe those pertaining to  $W$ .

The set  $W$  is generally not connected. It is also non-compact (since it is invariant and points are wandering under  $h$ ) and we will have to consider its compactly connected components. Let us recall (see [Moore, page 76] and also [LR2, Définition 9.1])

DEFINITION 3.1. — A space  $Z$  is compactly connected if any two points in  $Z$  are contained in a subcontinuum of  $Z$ . Distinct maximal compactly connected subsets of a space  $X$  are disjoint and are called the compactly connected components of  $X$ ; these components fill in  $X$ . Notice that these compactly connected components can be non closed.


LEMMA 3.2. — *The compactly connected components of  $W$  are unbounded.*

*Proof.* — We work in the Alexandroff compactification of  $\mathbf{R}^2$ , that is  $\mathbf{R}^2 \cup \{\infty\} \cong S^2$ . First,  $W \cup \{\infty\}$  is compact and connected as the decreasing intersection of the compact connected  $B_n \cup \{\infty\}$ . Suppose now that  $W$  admits a compactly connected component  $C$  contained in some open ball

$B(0, R)$ . Let then  $\tilde{C}$  be the component of  $W \cap B(0, R)$  which contains  $C$ . If  $\text{Adh}\tilde{C} \subset B(0, R)$  then  $\text{Adh}\tilde{C} = \tilde{C}$  is connected and compact so it is a connected compact component of  $W$ . As such, it is the intersection of the open and closed subsets of  $W$  which contain  $\tilde{C}$  [B, II §4.4], and there exists an open and closed neighborhood of  $\tilde{C}$  inside  $W \cap B(0, R)$ . (Indeed, if all open and closed neighborhoods  $(U_i)_{i \in I}$  of  $\tilde{C}$  in  $W$  meet  $\text{Fr}B(0, R)$ , we get a family pf closed sets  $(U_i \cap \text{Fr}B(0, R))$  in  $\text{Fr}B(0, R)$  such that every finite sub-family has a non empty intersection and therefore, the intersection of all these sets, which is  $\tilde{C}$ , is non empty in contradiction to the local hypothesis.) But this contradicts the connectivity of  $W \cup \{\infty\}$ . Therefore  $\text{Adh}\tilde{C}$  meets  $\text{Fr}B(0, R)$ . But  $\text{Adh}\tilde{C}$  is compact and connected in  $W$  so  $\text{Adh}\tilde{C} \subset C$  and  $C$  meets  $\text{Fr}B(0, R)$ : a contradiction.

Let us call  $C$  a **closed** compactly connected component of  $W$  and  $p$  an accessible point of  $C$  from  $\mathbf{R}^2 \setminus C$  :  $p$  is the extremity of an arc  $\gamma$  such that  $\gamma \setminus \{p\} \subset \mathbf{R}^2 \setminus C$ . We can suppose that  $\gamma$  is a free simple arc. Each  $h^n(l)$  has to meet  $\gamma$  and  $h(\gamma)$  for  $n$  larger than some  $n_0$  which we can suppose to be  $-1$ , replacing  $l$  by  $h^{n_0+1}(l)$  if necessary. Let  $p_n$  denote the last point of  $h^n(l)$  on  $\gamma$  as we move towards  $p$ . Then the arc  $\gamma_n = p_n p$  on  $\gamma$  is disjoint from all  $h^i(l), i \leq n$  except for  $p_n \in h^n(l)$ .

Let  $q_0 = h(p_{-1})$  and  $\alpha_0$  be the subarc  $p_0 q_0$  of  $l$ . Since  $\mathbf{R}^2 \setminus (W \cup W')$  is simply connected (even homeomorphic to  $\mathbf{R}^2$ ), it is divided by the arc  $\gamma_0 \cup \alpha_0 \cup h(\gamma_{-1})$  into two domains and we call  $\Omega$  the one which does not contain  $h^{-1}(l)$ .



PROPOSITION 3.3. — *The domain  $\Omega$  is free.*

*Proof.* — Suppose there exist  $x \in \Omega \cap h(\Omega)$  and let  $\beta$  be an arc from  $a$  to  $h^{-1}(x)$  with  $a \in \text{int}p_0 p$  and  $\beta \setminus \{a\} \subset \Omega$ . Since  $h$  preserves orientation,  $h(y) \notin \Omega$  for  $y$  close to  $a$  on  $\beta$ . As  $h(\beta) \cap h(p_0 p) = h(a)$  and  $h(\beta) \cap \alpha_0 = \emptyset$  (since  $h(\beta)$  is on the right of  $h(l)$  and so, on the right of  $l$  which contains  $\alpha_0$ ),

there exist some  $b \in \beta$  such that the subarc  $h(ab)$  of  $h(\beta)$  joins  $h(p_0p)$  to  $p_0p$  inside  $\mathbf{R}^2 \setminus (W \cup W' \cup \Omega)$  and the Jordan curve  $\alpha_0 \cup q_0h(a) \cup h(ab) \cup h(b)p_0$  contains the whole Brouwer line  $l$  or  $C$  (according to  $p_0$  or  $p$  is contained inside that Jordan curve) which is absurd since these sets are unbounded.  $\square$

PROPOSITION 3.4. — *The closed compactly connected component  $C$  cannot be  $h$ -invariant.*

*Proof.* — Assume by contradiction that  $h(C) = C$  and let then  $\tilde{K} \subset C$  be a continuum containing  $p$  and  $h(p)$ . Then  $\Omega$  is bounded and being simply connected has a boundary  $\text{Fr}\Omega$  which is connected and separating the plane. We first show  $K = \overline{\Omega} \cap \tilde{K}$  is compact and connected. It is enough to show that  $\text{Fr}\Omega \cap C$  is connected for then, if  $K = (\text{Fr}\Omega \cap C) \cap \tilde{K}$  is not connected then  $(\text{Fr}\Omega \cap C) \cup \tilde{K} \subset C$  separates the plane which contradicts the fact that  $C$  has no interior and does not separate. Let us note  $\delta = \gamma_0 \cup \alpha_0 \cup h(\gamma_{-1})$  so that  $\text{Fr}\Omega \cap C = \text{Fr}\Omega \setminus (\delta \setminus \{p, h(p)\})$ . If this last set is not connected we have two possibilities, the first one is that it has three components or more, and then  $\text{Fr}\Omega$  is not connected, the second one is that it has two components, containing  $p$  and  $h(p)$  respectively, which do not disconnect the plane and then  $\text{Fr}\Omega$  does not disconnect.

Therefore  $\Sigma = \overline{\bigcup_{n \in \mathbf{Z}} h^n(K)} \subset W$  is a closed connected set which is invariant under  $h$  and therefore non compact. As  $W$  does not separate  $\mathbf{R}^2$  and has no interior, the same is true of  $\Sigma$  and  $\mathbf{R}^2 \setminus \Sigma$  is homeomorphic to  $\mathbf{R}^2$ . The proper line  $l$  separates  $\mathbf{R}^2 \setminus \Sigma$  into two regions homeomorphic to  $\mathbf{R}^2$  and we name  $R$  the one between  $l$  and  $\Sigma$ . The region  $R$  itself is cut by the subarc  $p_0p$  of  $\gamma$  into two regions  $A$  and  $B$  where we call  $A$  the one containing  $\Omega$  and  $B$  the one containing  $h^{-1}(\Omega) \cap R$ . By definition  $p_0p$  is on the frontier of  $A$  and  $B$ . Notice that  $A$  (and  $B$ ) are non compact since we can follow  $l$  to infinity in one direction or the other staying in  $A$  (or  $B$ ). Note that  $A$  contains  $h^k(\Omega)$ ,  $k \geq 0$  and  $B$  contains  $h^{-k}(\Omega) \cap R$ ,  $k \geq 1$ .

LEMMA 3.5. —  *$\text{Fr}A \cap \text{Fr}B \cap \Sigma$  is non compact.*

*Proof.* — Let  $\Sigma_A$  (resp.  $\Sigma_B$ ) be the set of points of  $\Sigma$  which admit a neighborhood contained in  $A \cup \Sigma$  (resp.  $B \cup \Sigma$ ). The sets  $A \cup \Sigma_A$  and  $B \cup \Sigma_B$  are disjoint and open, therefore their complement in  $R \cup \Sigma \cup (p_0p \setminus \{p\})$  (which complement is the set of points of  $\Sigma$  for which every neighborhood meets  $A$  and  $B$ , that is  $\text{Fr}A \cap \text{Fr}B \cap \Sigma$ ) separates  $R \cup \Sigma \setminus (p_0p \setminus \{p\})$  and  $R \cup \Sigma \setminus (p_0p \setminus \{p\})$  can be written as the disjoint union  $(A \cup \Sigma_A) \amalg (B \cup \Sigma_B) \amalg (\text{Fr}A \cap \text{Fr}B \cap \Sigma)$ .

On the other hand, if  $\text{Fr}A \cap \text{Fr}B \cap \Sigma$  was compact in  $\mathbf{R}^2$  or equivalently in  $R \cup \Sigma$  (which is homeomorphic to  $\mathbf{R}^2$ ), thinking of  $l$  as a straight line



and of  $p_0p$  as a segment orthogonal to  $l$  (as it is legitimate by Schoenflies theorem), one can find a large rectangle in  $R \cup \Sigma$  with a side parallel to  $l$ , containing  $\text{Fr}A \cap \text{Fr}B \cap \Sigma$  and whose boundary cuts  $p_0p$  transversally in a single point. The boundary of this rectangle joins a point of  $A$  near  $p_0p$  to a point of  $B$  near  $p_0p$  in contradiction to the above decomposition of  $R \cup \Sigma \setminus (p_0p \setminus \{p\})$ .

Given Lemma 3.5, let us choose some point  $x$  in  $\text{Fr}A \cap \text{Fr}B \cap \Sigma$  and outside  $K$ . Then  $x \notin \overline{\Omega}$  and we choose an open euclidean ball  $2U \subset R \cup \Sigma$  centered at  $x$  free and disjoint of  $\overline{\Omega}$ . ( $U$  will denote the ball of radius one half the one of  $2U$ ). As  $x$  belongs to  $\Sigma$ ,  $U$  meets some  $h^m(K)$  and so some  $h^m(\Omega)$  and (exchanging  $h$  and  $h^{-1}$  if necessary) we can suppose  $m > 0$  and therefore that  $h^m(\Omega) \subset A$ . Since  $U$  meets  $B$ , we want to show that  $2U$  meets some  $h^{-n}(\Omega)$ , for some  $n > 0$ , for then  $2U$  and  $\Omega$  will give a contradiction to Lemma 2.1.

To that end, let us choose on  $\text{Fr}U$  two arcs, one on  $\text{Fr}U \cap A$  and the other on  $\text{Fr}U \cap B$  (these exist since  $U$  meets  $A$  and  $B$  which are open connected non compact) and choose an arc  $\alpha_0$  inside  $R \setminus (\Sigma \cup U)$  joining these two arcs and meeting transversally  $p_0p$  into a single point. Complete  $\alpha_0$  by a sub-arc  $\alpha_1$  of  $\text{Fr}U$ . This gives a Jordan curve  $\alpha$  inside  $R \cup \Sigma$  which contains  $p$  in its interior. Since points are wandering there exists  $N > 0$  such that  $h^{-N}(p) \in \Sigma$  belongs to the exterior of  $\alpha$ .

Now, if  $U$  does not meet any  $h^{-k}(\Omega)$ ,  $k > 0$ , the connected set  $\hat{K} = \bigcup_{i=1}^N h^{-i}(K)$  either joins  $p$  which is inside  $\alpha$  to  $h^{-N}(p)$  which is outside  $\alpha$  without meeting  $\alpha$  (in contradiction to the Jordan curve theorem), or it meets  $\alpha_1$  ( $\hat{K}$ , contained in  $\Sigma$ , does not meet  $\alpha_0$ ) and then  $\overline{U}$  meets some  $h^{-i}(K) \subset \hat{K}$  and so  $2U$  meets some  $h^{-i}(\Omega)$  and we are done. This concludes the proof of Proposition 3.4.

COROLLARY 3.6. —  $h^n(C) \cap C = \emptyset$  for all  $n \in \mathbf{Z} \setminus \{0\}$ .

*Proof.* — If  $h^n(C) \cap C \neq \emptyset$  then  $h^n(C) = C$  in contradiction to 3.4 applied to  $h^n$  which has the same  $W$  as  $h$ .  $\square$

Recall that given a sequence  $\{X_n\}_{n \in \mathbf{N}}$  of subspaces of a topological space  $Z$ , a point  $x \in Z$  belongs to  $\liminf X_n$  if every neighborhood of  $x$  meets  $X_n$  for an infinite number of  $n$  and to  $\limsup X_n$  if every neighborhood of  $x$  meets  $X_n$  for all but a finite number of  $n$ .

We will now suppose that  $X = \liminf h^n(C)$  is not empty. It is then a closed and non compact subset of  $W$  (since it is  $h$ -invariant). We aim to Proposition 3.10 below. Our first step is :


PROPOSITION 3.7. — *The set  $X$  is also  $\limsup h^n(C)$ . That is, every open set  $U$  which meets an infinite number of  $h^n(C)$ , meets  $h^n(C)$  for all  $n$  greater than some  $n_0 = n_0(U)$ .*

*Remark.* — This Proposition answers a question of F. Le Roux [LR2, footnote 7]

*Proof.* — We will use repeatedly the following immediate consequence of a result of Le Roux [LR2, Lemme 9.3], we repeat the proof here for completeness.

PROPOSITION 3.8. —  $X \cap h^n(C) = \emptyset$  for all  $n \in \mathbf{Z}$ .

*Proof.* — Since  $X$  is  $h$ -invariant, it is enough to show that  $X \cap C = \emptyset$ . Let us suppose  $X \cap C \neq \emptyset$ , and let  $U$  be a free neighborhood of  $x \in X \cap C$  such that  $\overline{U} \cap h(C) = \emptyset$ . As  $x \in X$ , there exists  $n > 1$  so that  $U \cap h^n(C) \neq \emptyset$ . Let  $y \in C$  such that  $h^n(y) \in U$ . There exists a continuum  $K \subset C$  which contains  $x$  and  $y$ . Since  $h(C)$  (as  $C$ ) is free, we can find a free connected neighborhood  $V$  of  $h(K) \subset h(C)$  such that  $U \cap V = \emptyset$ . But  $x \in U \cap h^{-1}(V)$  and  $h^n(y) \in U \cap h^{n-1}(V)$  so that  $U$  and  $V$  contradict Lemma 2.1.



Let  $V$  be a free open disc and  $D$  a component of  $V \setminus \overline{\bigcup_{n \in \mathbf{Z}} h^n(C)}$ .

LEMMA 3.9. — *If  $FrD$  meets  $h^n(C)$  and  $h^m(C)$ , then  $|n - m| \leq 1$  and  $FrD$  cannot meet  $X$  if it meets some  $h^n(C)$ .*

*Proof.* — To prove the first assertion, note that since  $X \cap h^n(C) = \emptyset$  for all  $n$ , given  $x \in h^n(C)$  there exists a disc neighborhood  $U$  of  $x$  which does not meet any other  $h^p(C)$  and a ray from  $x$  to some point in  $D \cap U$  leads

to an accessible point of  $h^n(C)$  from  $D$ . So let us suppose  $|n - m| > 1$  and let  $\alpha$  be an arc from  $a \in h^n(C)$  to  $b \in h^m(C)$  such that  $\alpha \setminus \{a, b\} \subset D$  and let  $K$  be a continuum in  $h^n(C)$  containing  $a$  and  $h^{n-m}(b)$ . We assert that  $K \cup \alpha$  is free. Indeed,  $K$  is free as a subset of  $h^n(C)$ ,  $\alpha$  is free as  $V$  is free and  $h(K) \cap \alpha = \emptyset = h^{-1}(K) \cap \alpha$  since  $n \pm 1 \neq m$ . But  $b \in h^{m-n}(K \cup \alpha) \cap (K \cup \alpha)$ , and a small enough neighborhood of  $K \cup \alpha$  will contradict Lemma 2.1 if  $|n - m| > 1$ .

Let us suppose now that  $\text{Fr}D$  meets  $X$  and some  $h^n(C)$  and let again  $U$  be a disc neighborhood of some point  $x \in \text{Fr}D \cap X$  small enough so that  $U \cap h^k(C) = \emptyset$  if  $|k| \leq |n| + 1$ . A ray issued from  $x$  will either give an accessible point of some  $h^m(C)$ ,  $|m| > |n| + 1$  from  $D$ , but this is impossible according to the first part of the proof, or an accessible point of  $X$  from  $D$ . In that case, let  $\alpha$  be an arc from some point  $a \in h^n(C)$  to  $b \in X$  with  $\alpha \setminus \{a, b\} \subset D$  and let  $U'$  be a free neighborhood of  $b$  such that  $U' \cap h^k(C) = \emptyset$ , for  $|k| \leq |n| + 1$  and such that  $U' \cap h^{\pm 1}(\alpha) = \emptyset$ . The arc  $\alpha$  can be extended to an arc  $\tilde{\alpha} \subset \alpha \cup U'$  which joins  $a \in h^n(C)$  to some  $\tilde{b} \in h^m(C)$ ,  $|m| > |n| + 1$ . If  $K \subset h^n(C)$  is a continuum containing  $a$  and  $h^{n-m}(\tilde{b})$ , then  $K \cup \tilde{\alpha}$  is a free continuum such that  $\tilde{b} \in h^{m-n}(K \cup \tilde{\alpha}) \cap (K \cup \tilde{\alpha})$  and a free neighborhood of this continuum gives a contradiction to Lemma 2.1.  $\square$

We now return to the proof of Proposition 3.7. Let  $V$  be a free neighborhood of  $x \in \liminf h^n(C)$ . There exist  $m$  and  $n > m + 1$  such that  $V$  meets  $h^n(C)$  and  $h^m(C)$ . Let  $\alpha$  be an arc in  $V$  going from  $a_m \in h^m(C)$  to  $a_n \in h^n(C)$  disjoint from  $h^m(C)$  and  $h^n(C)$  except for its extremities. Let  $D$  be the component of  $V \setminus \overline{\bigcup_{n \in \mathbf{Z}} h^n(C)}$  which meets  $\alpha$  and has  $a_n$  on its frontier. By 3.9,  $\text{Fr}D$  meets  $h^{n+1}(C)$  or  $h^{n-1}(C)$ . In the first case, let  $a_{n+1}$  be the last point of  $h^{n+1}(C)$  seen on  $\alpha$  when going from  $a_n$  to  $a_m$ . If  $D'$  is the component of  $V \setminus \overline{\bigcup_{n \in \mathbf{Z}} h^n(C)}$  which meets the subarc  $a_m a_{n+1}$  of  $\alpha$  and has  $a_{n+1}$  on its frontier, then  $\text{Fr}D'$  does not meet  $h^n(C)$  by construction of  $\alpha$  and therefore, according to 3.9, meets  $h^{n+2}(C)$ . Iterating this process we see that  $\alpha$  meets all  $h^k(C)$ ,  $k \geq n$ . In the other case, the same reasoning shows that  $\alpha$  meets all the  $h^k(C)$  for  $m \leq k \leq n$ . As  $V$  meets an infinite number of  $h^k(C)$  we conclude in either case that  $V$  meets all  $h^k(C)$  for  $k$  large enough and therefore  $x \in \limsup h^n(C)$ .

**Assumption:** We assume for the rest of this paper that the compactly connected components of  $W$  (in fact, we will only consider those of  $X$ ) are closed.

PROPOSITION 3.10. — *No point of  $X$  is accessible from  $\mathbf{R}^2 \setminus \overline{\bigcup_{n \in \mathbf{Z}} h^n(C)}$ .*

*Proof.* — We begin with a lemma :

**LEMMA 3.11.** — *There is no free arc  $\alpha$  joining  $C$  to  $X$  contained in  $\mathbf{R}^2 \setminus \overline{\bigcup_{n \in \mathbf{Z}} h^n(C)}$  except for its extremities.*

*Proof of 3.11.* — Let a free arc  $\alpha$  join  $p \in C$  to  $q \in X$  and consider a neighborhood  $D$  of  $q$  such that  $\alpha \cup D$  is still free and  $D \cap h(C) = \emptyset = D \cap h^{-1}(C)$  (recall that  $X$  is disjoint from  $h(C)$  and  $h^{-1}(C)$  by proposition 3.8). Then  $\alpha \cup D$  contains a point  $h^n(p')$  for some  $n > 1$  and some  $p' \in C$ . Let  $K \subset C$  be a continuum containing  $p$  and  $p'$  and consider the continuum  $L = K \cup \alpha \cup D$ . It is free but  $h^n(p') \in h^n(L) \cap L$  and a small enough neighborhood of  $L$  gives a contradiction to Lemma 2.1.  $\square$

At this point we will finish the proof of 3.10 following the lines of the proof of a similar result (with  $C$  replaced by a disc) in [LR2, Proposition 5.5].

Let us suppose there exist a point  $q$  of  $X$  accessible from  $\mathbf{R}^2 \setminus \overline{\bigcup_{n \in \mathbf{Z}} h^n(C)}$  by some arc  $\alpha$  and let  $Z$  be the connected component of  $X$  which contains  $q$ . A point  $x$  of  $\mathbf{R}^2 \setminus \overline{\bigcup_{n \in \mathbf{Z}} h^n(C)}$  will be called a *neighborhood point* of  $Z$  if there exists a free closed euclidean disc  $D$  with center  $x$  such that  $\text{int}D \cap Z \neq \emptyset$ . The set of all such points is an open set  $V$  and  $V \cup Z$  is an open neighborhood of  $Z$ .

A point of  $x \in V$  will be said of *type  $C$*  if there is some euclidean disc  $D$  of center  $x$  as in the previous definition and an arc in  $D$  from  $x$  to  $Z$  which meets some  $h^n(C)$  and of *type  $Z$*  if there exists such a disc  $D$  and an arc in  $D$  from  $x$  to  $Z$  which does not meet any  $h^n(C)$ . It follows from Lemma 3.11 that this type is well defined.

We show that all points of  $V$  are of type  $C$ . Indeed, it is easily verified that the type is locally constant on  $V$  and so is constant on every connected component of  $V$ . But  $V \cup Z$  and  $\mathbf{R}^2 \setminus Z$  are connected and therefore their intersection  $V$  also as follows from the Mayer-Vietoris sequence of the pair  $(\mathbf{R}^2 \setminus Z, V \cup Z)$ . Furthermore, since  $Z \subset X$ , certainly  $V$  meets some  $h^n(C)$  and all points of  $V$  are of type  $C$ .

Now, if the point  $x$  on the arc  $\alpha$  is close enough to  $q$ , the subarc  $xq$  of  $\alpha$  is contained in a free euclidean disc which meets  $Z$ , and,  $x$  being of type  $C$ , meets some  $h^n(C)$ . Contradiction.

## Bibliography

- [Br] BROUWER (L.E.J.). — Beweis des ebenen Translationssatzes, *Math. Ann.*, 72, p. 37-54 (1912).
- [BCL] BÉGUIN (F.), CROVISIER (S.) and LEROUX (F.). — Pseudo-rotations of the open annulus, *Bull. Braz. Math. Soc.* 37, p. 275-306 (2006).
- [B] BOURBAKI (N.). — *Topologie générale*, Hermann, Paris (1971).
- [G1] GUILLOU (L.). — Théorème de translation plane de Brouwer et généralisations du théorème de Poincaré-Birkhoff, *Topology*, 33, p. 331-351 (1994).
- [G2] GUILLOU (L.). — Free lines for homeomorphisms of the open annulus, *Trans. AMS*, 360, p. 2191-2204 (2008).
- [HK] HOMMA (T.), TERASAKA (H.). — On the structure of the plane translation of Brouwer, *Osaka Math. J.* 5, p. 233-266 (1953).
- [LC] LE CALVEZ (P.). — Rotation numbers in the infinite annulus, *Proc. Amer. Math. Soc.* 129, p. 3221-3230 (2001).
- [LR1] LEROUX (F.). — Bounded recurrent sets for planar homeomorphisms, *Ergodic theory Dynam. Systems* 19, p. 1085-1091 (1999).
- [LR2] LEROUX (F.). — Structure des homéomorphismes de Brouwer, *Geometry and Topology* 9, p. 1689-1774, (2005).
- [M] MOORE (R.L.). — *Foundations of Point Set Theory*, AMS Colloquium Publications volume XIII (1962).
- [W] WINKELNKEMPER (H.). — A generalisation of the Poincaré-Birkhoff theorem, *Proc. AMS*, 102, p. 1028-1030 (1988).