



Annales de la Faculté des Sciences de Toulouse

MATHÉMATIQUES

ALEXEI BORODIN AND SERGEI KOROTKIKH
Inhomogeneous spin q -Whittaker polynomials

Tome XXXIII, n° 1 (2024), p. 1–68.

<https://doi.org/10.5802/afst.1761>

© les auteurs, 2024.

Les articles des *Annales de la Faculté des Sciences de Toulouse* sont mis à disposition sous la licence Creative Commons Attribution (CC-BY) 4.0
<http://creativecommons.org/licenses/by/4.0/>



Publication membre du centre
Mersenne pour l'édition scientifique ouverte
<http://www.centre-mersenne.org/>
e-ISSN : 2258-7519

Inhomogeneous spin q -Whittaker polynomials ^(*)ALEXEI BORODIN ⁽¹⁾ AND SERGEI KOROTKIKH ⁽²⁾

ABSTRACT. — We introduce and study an inhomogeneous generalization of the spin q -Whittaker polynomials from [15]. These are symmetric polynomials, and we prove a branching rule, skew dual and non-dual Cauchy identities, and an integral representation for them. Our main tool is a novel family of deformed Yang–Baxter equations.

RÉSUMÉ. — Nous introduisons et étudions une généralisation inhomogène des polynômes spin de q -Whittaker de [15]. Ce sont des polynômes symétriques, et nous prouvons une règle de branchement, des identités de Cauchy asymétriques duales et non-duales, et une représentation intégrale pour ces polynômes. Nous prouvons une règle de branchement, des identités de Cauchy asymétriques, duales et non-duelles, et une représentation intégrale pour ces polynômes. Notre outil principal est une nouvelle famille d'équations de Yang–Baxter déformées.

1. Introduction**Background**

Connections between integrable lattice models and the theory of symmetric functions have been long known to be important to both domains; one might, for example, consult the following works (the list is certainly incomplete): Fomin–Kirillov [22], Tsilevich [35], Brubaker–Bump–Friedberg [17], Zinn–Justin [38], Korff [25], Wheeler–Zinn–Justin [36, 37], Motegi–Sakai [31], Motegi [30], Cantini–de Gier–Wheeler [21], Garbali–de Gier–Wheeler [23],

(*) Reçu le 23 avril 2021, accepté le 11 février 2022.

(1) Department of Mathematics, Massachusetts Institute of Technology, Cambridge MA 02139, USA — borodin@math.mit.edu

(2) Department of Mathematics, Massachusetts Institute of Technology, Cambridge MA 02139, USA — korotkih@mit.edu

A. B. was partially supported by the NSF grant DMS-1853981 and the Simons Investigator program. A. B. and S. K. were also partially supported by the NSF FRG grant DMS-1664619.

Article proposé par Laurent Miclo.

Borodin [4, 5], Borodin–Petrov [9, 10], Borodin–Wheeler [13, 15], Garbali–Wheeler [24], Buciumas–Scrimshaw [18], Mucciconi–Petrov [32], Petrov [33].

In particular, in [4, 9, 10, 15] families of *spin* Hall–Littlewood and q -Whittaker symmetric functions were introduced. They generalized the classical families of the Hall–Littlewood and q -Whittaker symmetric polynomials (both of which are special cases of the Macdonald symmetric functions, see Macdonald [28]), and the additional *spin parameter* was related to the spin in the underlying integrable vertex models. It was shown, in particular, that symmetric functions from those families satisfy explicit branching rules as well as (skew) Cauchy and dual Cauchy identities. A different version of the spin q -Whittaker polynomials was also recently suggested in [32]; it appears to enjoy a few valuable properties that the original one lacked.

The introduction of the spin Hall–Littlewood functions was prompted by their usefulness in asymptotic analysis of a variety of probabilistic systems that include certain integrable models of directed polymers in random media and exclusions processes in (1+1)-dimensions, see Borodin–Corwin–Petrov–Sasamoto [6, 7], where they arose as an orthogonal basis of eigenfunctions for Hamiltonians of certain associated (“dual”) quantum integrable systems. Probabilistic applications of the spin q -Whittaker polynomials were demonstrated by Bufetov–Mucciconi–Petrov [20] and Mucciconi–Petrov [32].

In [10] an inhomogeneous version of the spin Hall–Littlewood functions was introduced. The inhomogeneities appear naturally from the point of view of the vertex models involved, and they are also crucial for some of the probabilistic applications, such as the analysis of the Asymmetric Simple Exclusion Process (ASEP) and the stochastic six vertex model with varying initial conditions, see Aggarwal–Borodin [2] and Aggarwal [1], and large time asymptotics of the exponential jump model of Borodin–Petrov [11]. Curiously, while the inhomogeneities significantly change the symmetric functions, they do not affect the product factors of the (skew) Cauchy identities.

The main goal of the present work is to define and study inhomogeneous spin q -Whittaker polynomials. So far it has been unclear how to do that. Even though the approach of [15], that allowed to obtain the homogeneous spin q -Whittaker polynomials from the Hall–Littlewood ones via the procedure of *fusion*, is also applicable in the inhomogeneous case, the resulting functions do not appear to satisfy the dual Cauchy identity involving both q -Whittaker and Hall–Littlewood functions, which is the most natural requirement. Indeed, we show that the “correct” inhomogeneous spin q -Whittaker polynomials (those that do satisfy the dual Cauchy identity with the inhomogeneous spin Hall–Littlewood functions) are different from the functions naively obtained via fusion. These two families (the “naive” and the “correct” one) can be united in a (non-dual) Cauchy identity, however.


With equal inhomogeneities our inhomogeneous polynomials degenerate to those of [15]. It remains to be seen if the modified definitions of [32] can be applied in the inhomogeneous setting, and if this would again lead to more desirable properties.

In the next subsection we list the properties of these new inhomogeneous symmetric polynomials that we were able to prove. The new key ingredient of our approach is a *deformed Yang–Baxter equation* for the vertex weights of the underlying integrable vertex model; it is also discussed below.

Main results


For a pair of sequences of complex parameters $\mathbf{S} = (s_0, s_1, s_2, \dots)$ and $\Xi = (\xi_0, \xi_1, \xi_2, \dots)$ we define functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$, which depend on n variables $\kappa_1, \dots, \kappa_n$ and are labeled by pairs of partitions λ, μ . Here are the main facts we prove about them.

- The functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ are symmetric polynomials in $\kappa_1, \dots, \kappa_n$.
- For a single variable, $\mathbb{F}_{\lambda/\mu}(\kappa \mid \Xi, \mathbf{S})$ has a fully factorized expression. To write it down, we first define an auxiliary function $ZW_{\lambda/\mu}(\kappa \mid \Xi, \mathbf{S})$ as a partition function of a one-row vertex model of the form



This partition function is the product of weights of all the vertices in this row, where each vertex has four nonnegative integral labels on the edges incident to it. The labels on the vertical edges are the multiplicities in λ' and μ' : for $j \geq 1$, $m_j(\mu') = \mu_j - \mu_{j+1}$, $m_j(\lambda') = \lambda_j - \lambda_{j+1}$. For any $k \geq 1$ we defined $s_k^{(1)} := \sqrt{s_k s_{k+1} \xi_{k+1} / \xi_k}$. The weights of the vertices are given by (note that all the square roots

will disappear upon substitution into these weights)



$$l = \mathbb{1}_{i+j=k+l} \mathbb{1}_{i \geq l} s^{2l} t^{-2l} \frac{(s^2/t^2; q)_{i-l} (t^2; q)_l}{(s^2; q)_i} \frac{(q; q)_i}{(q; q)_{i-l} (q; q)_l}.$$

Even though the row is semi-infinite, the number of vertices of weight not equal to 1 is readily seen to be finite.

Then the polynomial $\mathbb{F}_{\lambda/\mu}(\kappa \mid \Xi, \mathbf{S})$ is defined by the following expression:

$$\begin{aligned} \mathbb{F}_{\lambda/\mu}(\kappa \mid \Xi, \mathbf{S}) &:= (-1)^{|\mu| - |\lambda|} \frac{(\tau_{\Xi} \mathbf{S})^\mu}{\mathbf{S}^\lambda} \frac{c_{\tau_{\Xi} \mathbf{S}}(\mu)}{c_{\mathbf{S}}(\lambda)} (\kappa s_0 / \xi_0)^{\lambda_1 - \mu_1} \\ &\quad \times \frac{(\kappa^{-1} s_1 \xi_1; q)_{\lambda_1 - \mu_1}}{(q; q)_{\lambda_1 - \mu_1}} ZW_{\lambda/\mu}(\kappa \mid \Xi, \mathbf{S}), \end{aligned}$$

where $\tau_{\Xi} \mathbf{S} = (s_1^{(1)}, s_2^{(1)}, \dots)$, and

$$c_{\mathbf{S}}(\lambda) := \prod_{i \geq 1} \frac{(s_i^2; q)_{\lambda_i - \lambda_{i+1}}}{(q; q)_{\lambda_i - \lambda_{i+1}}}, \quad \mathbf{S}^\lambda := \prod_{i \geq 1} s_i^{\lambda_i}.$$

- The following branching rule holds:

$$\begin{aligned} \mathbb{F}_{\lambda/\nu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) \\ = \sum_{\mu} \mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_m \mid \Xi, \mathbf{S}) \mathbb{F}_{\mu/\nu}(\kappa_{m+1}, \dots, \kappa_n \mid \tau_{\mathbf{S}}^m \Xi, \tau_{\Xi}^m \mathbf{S}), \end{aligned}$$

where $\tau_{\mathbf{S}}^k \Xi = (\xi_0^{(k)}, \xi_1^{(k)}, \xi_2^{(k)}, \dots)$, $\tau_{\Xi}^k \mathbf{S} = (s_0^{(k)}, s_1^{(k)}, s_2^{(k)}, \dots)$ with

$$\xi_i^{(k)} := \sqrt{\xi_{i+k} s_{i+k} \xi_i / s_i}, \quad s_i^{(k)} := \sqrt{s_{i+k} \xi_{i+k} s_i / \xi_i},$$

Together with the partition function expression for the one-variable polynomials, this allows to write the general $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ in terms of the partition function for the vertex model in Figure 1.1, see Section 5.1.

- The functions $\mathbb{F}_{\lambda}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) := \mathbb{F}_{\lambda/\emptyset}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ satisfy the following stability relation:

$$\mathbb{F}_{\lambda}(\kappa_1, \dots, \kappa_{n-1}, s_n \xi_n \mid \Xi, \mathbf{S}) = \mathbb{F}_{\lambda}(\kappa_1, \dots, \kappa_{n-1} \mid \Xi, \mathbf{S}).$$

- We have the following dual Cauchy identity:

$$\sum_{\lambda} \tilde{\mathbb{F}}_{\lambda}^*(u_1, \dots, u_m \mid \Xi, \mathbf{S}) \mathbb{F}_{\lambda}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) = \prod_{i=1}^n \prod_{j=1}^m \frac{1 - u_j \kappa_i}{1 - u_j \xi_i s_i},$$

where $\tilde{\mathbb{F}}_{\lambda'}^*$ denotes a certain stable version of the inhomogeneous spin Hall–Littlewood functions described in Section 5.2. This identity is a particular case of a more general dual skew Cauchy identity, which is proved in Theorem 5.10. In these dual Cauchy identities the number of nonzero terms is always finite, thus no convergence conditions are needed.

- Let $\bar{\mathbf{S}} = (s_0^{-1}, s_1^{-1}, s_2^{-1}, \dots)$, then the following Cauchy-type summation identity holds:

$$\sum_{\lambda} \mathbb{F}_{\lambda}^*(\chi_1, \dots, \chi_m \mid \bar{\mathbf{S}}, \mathbf{S}) \mathbb{F}_{\lambda}(\kappa_1, \dots, \kappa_n \mid \mathbf{S}, \mathbf{S}) = \prod_{i=1}^n \prod_{j=1}^m \frac{(\kappa_i; q)_{\infty} (s_i^2 \chi_j; q)_{\infty}}{(s_i^2; q)_{\infty} (\kappa_i \chi_j; q)_{\infty}},$$

where \mathbb{F}_{λ}^* denotes a certain rescaling of the functions \mathbb{F}_{λ} . This identity could be either viewed as holding in suitable rings of power series or numerically under certain restrictions on the parameters, see Theorem 6.1. There is also a more general skew version of this identity, provided in Theorem 6.1 as well.

- The polynomials \mathbb{F}_{μ} admit an integral representation of the form

$$\begin{aligned} \mathbb{F}_{\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) &= \oint_{\mathcal{C}} \frac{dz_1}{2\pi i z_1} \cdots \oint_{\mathcal{C}} \frac{dz_k}{2\pi i z_k} \prod_{\alpha < \beta} \frac{z_{\alpha} - z_{\beta}}{z_{\alpha} - q z_{\beta}} \\ &\quad \times \prod_{\alpha=1}^k \left(\frac{\xi_0^{-1}}{z_{\alpha} - \xi_{\mu'_{\alpha}}^{-1} s_{\mu'_{\alpha}}} \prod_{j=1}^{\mu'_{\alpha}-1} \frac{1 - s_j \xi_j z_{\alpha}}{z_{\alpha} \xi_j - s_j} \prod_{i=1}^n \frac{1 - z_{\alpha} \kappa_i}{1 - z_{\alpha} \xi_i s_i} \right), \end{aligned}$$

where $k = \mu_1$, and the integration contour \mathcal{C} is depicted in Figure 5.3 and described immediately before Theorem 5.13.


Comparing with [15], for $s_0 = s_1 = \dots = s$ and $\xi_0 = \xi_1 = \dots = \xi$ our functions $\tilde{\mathbb{F}}_{\lambda/\mu}$ and $\mathbb{F}_{\lambda/\mu}$ reduce to the spin q -Whittaker and stable spin Hall–Littlewood functions from [15]; in particular, our functions degenerate to the usual Hall–Littlewood and q -Whittaker polynomials when $s = 0$.

Deformed Yang–Baxter equations

The key tool that allowed us to prove most of the results above is a family of identities that we call *deformed Yang–Baxter equations*. Their detailed exposition can be found in Section 3. While the conventional Yang–Baxter equations for vertex models have the general form

$$\mathcal{R}_{\mathbf{x}}^{(12)} \mathcal{R}_{\mathbf{y}}^{(13)} \mathcal{R}_{\mathbf{z}}^{(23)} = \mathcal{R}_{\mathbf{z}}^{(23)} \mathcal{R}_{\mathbf{y}}^{(13)} \mathcal{R}_{\mathbf{x}}^{(12)},$$

where $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are parameters subject to certain relations (in our setting the parameters $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are pairs of complex numbers), and $\mathcal{R}_{\mathbf{x}}^{(12)}, \mathcal{R}_{\mathbf{y}}^{(13)}, \mathcal{R}_{\mathbf{z}}^{(23)}$


 Figure 1.1. The model used to compute the function $\mathbb{F}_{\lambda/\mu}$.

are operators acting in two of the three tensor factors in a tensor product $V_1 \otimes V_2 \otimes V_3$ with suitable linear spaces V_i . It turns out that in certain situations one can add an additional degree of freedom resulting in equations of the form

$$\mathcal{R}_{\mathbf{x}(\tau)}^{(12)} \mathcal{R}_{\mathbf{y}(\tau)}^{(13)} \mathcal{R}_{\mathbf{z}(\tau)}^{(23)} = \mathcal{R}_{\mathbf{z}'(\tau)}^{(23)} \mathcal{R}_{\mathbf{y}'(\tau)}^{(13)} \mathcal{R}_{\mathbf{x}'(\tau)}^{(12)},$$

where the subscripts $\mathbf{x}(\tau), \mathbf{y}(\tau), \mathbf{z}(\tau)$ and $\mathbf{x}'(\tau), \mathbf{y}'(\tau), \mathbf{z}'(\tau)$ now depend on a new parameter τ . Note that the deformed equation has different operators in the left-hand and right-hand sides.

In this work we provide two examples of the deformed Yang–Baxter equation; one of them implies that the functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ are symmetric, while the other one is crucial for proving the Cauchy identities. Additionally, we describe a method we use to deform the previously known Yang–Baxter equations. Such deformations might be useful in other contexts. For instance, in the end of Section 3 we briefly mention a different application: One can use a deformation of a higher rank Yang–Baxter equation to extend the results of [14] on integral representations for certain observables of a colored stochastic higher spin six-vertex model. Unfortunately, so far our understanding of the deformed Yang–Baxter equation is based solely on computational tricks we use to derive them; we are not aware of either a representation theoretic meaning of such equations or a complete list of situations where one might expect to see them.

Applications to probability

The inhomogeneous spin q -Whittaker functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ can be defined in terms of a new stochastic integrable vertex model with parameters attached to diagonals, see Section 5. Earlier results [9, 10, 14] suggest that this model could be studied via Cauchy-type identities, which would lead to explicit expressions for averages of certain natural observables. Indeed, such an approach is also applicable to our new model. However, very recently another method of analyzing stochastic vertex models was introduced by Bufetov–Korotkikh [19]. It is less constructive but yields more general results; its key feature is a certain local relation for q -moments of the height functions for the model. The adaptation of this method to the setting of the present paper is done in [26], and it leads to novel probabilistic results that include, in particular, large scale asymptotics for partition functions of a new integrable extension of the Beta polymer model of Barraquand–Corwin [3].

Layout of the paper

In Section 2, we recall the definition and properties of the (stochastic) higher spin six vertex model and its extensions obtainable by fusion. In particular, we list several instances of the Yang–Baxter equation used throughout the work. In Section 3, we introduce the deformed Yang–Baxter equations; our construction there is based on certain rationality properties of the vertex weights. Section 4 is devoted to *row operators*. On one hand, these operators encode partition functions of single rows of a vertex model subject to certain boundary conditions, while on the other hand, the row operators capture the branching structure of our symmetric functions. In the same section we also prove exchange relations between the row operators using the (deformed) Yang–Baxter equations and a “zipper” argument. Having developed the framework of row operators, in Section 5 we define the inhomogeneous spin q -Whittaker functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ and provide proofs of the above-listed properties. Finally, in Section 6 we state and prove the (skew) Cauchy identity between the functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \mathbf{S}, \mathbf{S})$ and $\mathbb{F}_{\lambda/\mu}(\chi_1, \dots, \chi_m \mid \bar{\mathbf{S}}, \mathbf{S})$.

Additionally, in the end of the paper we include an appendix with a condensed list of the numerous expressions used in the work, to help the potential reader to follow the text.

Notation

In this work we follow the standard notations regarding partitions: A *partition* λ is a monotone infinite sequence $(\lambda_1, \lambda_2, \lambda_3, \dots)$ of nonnegative integers satisfying

$$\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \dots \geq 0, \quad |\lambda| := \lambda_1 + \lambda_2 + \lambda_3 + \dots < \infty.$$

The coordinates λ_i are called parts of the partition. For a partition λ , its *length* $l(\lambda)$ is equal to the number of nonzero parts λ_i , while $m_k(\lambda)$ is used to denote the number of parts λ_i equal to k . The *conjugate* λ' of a partition λ is defined by $\lambda'_i = \#\{j \geq 1 : \lambda_j \geq i\}$.

For a pair of partitions λ, μ , we write $\mu \subset \lambda$ iff for any $i \geq 1$ we have $\mu_i \leq \lambda_i$. Further, we say the partition λ *interlaces* the partition μ , and write $\lambda \succ \mu$, if $\lambda_i \geq \mu_i \geq \lambda_{i+1}$ for all $i \geq 1$. Note that in this case we also have $0 \leq l(\lambda) - l(\mu) \leq 1$.

Throughout the text we use the following notation for q -Pochhammer symbol:

$$(x; q)_n := \prod_{i=1}^n (1 - xq^{i-1}),$$

with the convention $(x; q)_0 := 1$. We also often assume that $|q| < 1$, so the notation $(x; q)_\infty$ makes sense both numerically and formally in the space of power series in q .

2. Preliminaries on vertex models and the Yang–Baxter equation

Vertex models considered in this work consist of collections of vertices with oriented edges between them. A configuration of a model is an assignment of integer labels to all edges in a way such that the (*arrow*) *conservation law* holds: for each vertex the sum of labels of the incoming edges equals the sum of labels of the outgoing ones. To each configuration we assign a weight, which is equal to the product of local weights corresponding to each vertex and depending on the labels around the vertex. In this section we describe various local weights, as well as recall several known properties of them.

2.1. Higher spin vertex weights

We start with the (stochastic) higher spin six vertex model. Our exposition in this section follows [10] and [15], up to a change of notation discussed in Remark 2.1 below.

Here \mathcal{R}_z denotes the \mathcal{R} -matrix defined by

$$\begin{aligned}\mathcal{R}_z(0, 0; 0, 0) &= \mathcal{R}_z(1, 1; 1, 1) = 1 - qz, \\ \mathcal{R}_z(0, 1; 0, 1) &= 1 - z, \quad \mathcal{R}_z(0, 1; 1, 0) = z(1 - q), \\ \mathcal{R}_z(1, 0; 0, 1) &= 1 - q, \quad \mathcal{R}_z(1, 0; 1, 0) = q(1 - z),\end{aligned}$$

and for all values not listed above $\mathcal{R}_z(i, j; k, l) = 0$.

Throughout this work, instead of writing equations like (2.3), we extensively use graphical notation for partition functions of vertex models. In general, we consider a collection of vertices and oriented edges (possibly of different kinds) connected to them, with some edges connecting a pair of vertices, and some edges being connected to only one vertex with the other end being free. The edges of the former type are called *internal edges*, while the edges of the latter type are referred to as *boundary edges*. Additionally, to each vertex we assign a family of vertex weights, with each weight corresponding to a particular configuration of labels of the edges connected with the vertex, for example (2.1), and to each boundary edge we assign a label which is called the *boundary condition* corresponding to the edge. Given such data, the corresponding *partition function* is defined as the sum of the products of the weights of all vertices, taken over all possible assignments of labels to the internal edges satisfying the conservation law.

For example, the Yang–Baxter equation (2.3) for the weights $w_{u;s}^s$ can be graphically interpreted as follows:

$$\begin{array}{c} b_1 \\ \uparrow \\ a_3 \text{ --- } w_{y;s}^s \\ \text{---} \\ a_2 \text{ --- } w_{x;s}^s \\ \downarrow \\ a_1 \end{array} \begin{array}{c} b_2 \\ \nearrow \\ \mathcal{R}_{x/y} \\ \searrow \\ b_3 \end{array} = \begin{array}{c} b_1 \\ \uparrow \\ a_3 \text{ --- } w_{x/y}^s \\ \text{---} \\ a_2 \text{ --- } w_{y;s}^s \\ \downarrow \\ a_1 \end{array} \begin{array}{c} b_2 \\ \nearrow \\ \mathcal{R}_{x/y} \\ \searrow \\ b_3 \end{array} \quad (2.4)$$

The vertices with horizontal and vertical edges have the weights $w_{u;s}^s$ given by (2.1), while the vertices with diagonal edges, which we call *tilted* vertices, correspond to the \mathcal{R} -matrix:

$$\begin{array}{c} j \\ \nearrow \\ \mathcal{R}_z \\ \searrow \\ i \end{array} \begin{array}{c} k \\ \nearrow \\ \\ \searrow \\ l \end{array} = \mathcal{R}_z(i, j; k, l).$$

Note that the diagrams we use have two kinds of edges, which visually differ by their thickness. Throughout this work we follow the convention that thin edges carry labels from $\{0, 1\}$, while thick edges are labeled by any non-negative integer.

Remark 2.1. — The letter s in the superscript of the notation $w_{u;s}^s$ stands for “stochastic”, and should not be confused with the spin parameter s of a vertex. We use such a notation to distinguish our weights from the weights $w_{u;s}$ often used in this context, cf. [4, 10, 15]. Our weights $w_{u;s}^s$ coincide with the weights $L_{u,s}$ used in [10], and they are related to the weights $w_{u;s}$ by [10, (2.2) and (2.3)]:

$$w_{u;s}^s(i, j; k, l) = (-s)^l \frac{(s^2; q)_k}{(q; q)_k} \frac{(q; q)_i}{(s^2; q)_i} w_{u;s}(i, j; k, l).$$

The word “stochastic” refers to the fact that the sum of these weights over different labels of the outgoing edges (and fixed values of the incoming ones) is identically equal to 1, cf. Remark 2.2 below.

2.2. Fusion

The weights $w_{u;s}^s$ can be used to reach more general vertex models by a procedure called *fusion* that we review below. See [15, Section 4] and [10, Section 5] for a more detailed exposition of the fusion procedure, as well as further references.

Fix $J > 0$ and consider a column consisting of J vertices having weights $w_{u;s}^s, w_{qu;s}^s, \dots, w_{q^{J-1}u;s}^s$ starting from the bottom. This configuration is called a J -column, and using the Yang–Baxter equation one can show that, under specific boundary left and right conditions, this column of vertices behaves like a single vertex of a new type, with left and right edges having labels from $\{0, \dots, J\}$.

More precisely, for $j \geq 0$ set

$$Z_j(J) := \sum_{a_1 + \dots + a_J = j} q^{\sum_{r=1}^J (r-1)a_r} = \frac{q^{\frac{j(j-1)}{2}} (q; q)_J}{(q; q)_j (q; q)_{J-j}},$$

where the sum is taken over all sequences $(a_1, \dots, a_J) \in \{0, 1\}^J$ satisfying $\sum_{r=1}^J a_r = j$, and the equality is a particular case of the q -binomial identity. For $i, j, k, l \in \mathbb{Z}_{\geq 0}$ such that $i + j = k + l$ and $j, l \leq J$ define vertex weights

$\mathcal{W}_{u;s}^{(J)}$ by

$$\mathcal{W}_{u;s}^{(J)}(i, j; k, l) := \frac{Z_l(J)}{Z_j(J)} q^{-\sum_{r=1}^J (r-1)b_r} \times \sum_{a_1 + \dots + a_J = j} q^{\sum_{r=1}^J (r-1)a_r} \left(\begin{array}{c} k \\ \uparrow \\ a_J \text{ --- } w_{q^{J-1}u;s}^s \text{ --- } b_J \\ \vdots \\ a_2 \text{ --- } w_{qu;s}^s \text{ --- } b_2 \\ a_1 \text{ --- } w_{u;s}^s \text{ --- } b_1 \\ \vdots \\ i \end{array} \right). \quad (2.5)$$

Here $(b_1, \dots, b_J) \in \{0, 1\}^J$ is any vector satisfying $\sum_{r=1}^J b_r = l$. It turns out that the right-hand side of (2.5) does not depend on the particular choice of (b_1, \dots, b_J) , since a direct computation for $J = 2$ shows that the right-hand side is not changed when b_i and b_{i+1} are swapped, see [10, Proposition 5.4]. This independence property is called q -exchangeability and it allows to stack (2.5) horizontally, identifying a row of J -columns with a single row of “fused” vertices: for $i_1, \dots, i_L, j, k_1, \dots, k_L, l \in \mathbb{Z}_{\geq 0}$ such that $\sum_{m=1}^L i_m + j = \sum_{m=1}^L k_m + l$ we have

$$\mathcal{W}_{u_1;s_1}^{(J)}(i_1, j_0; k_1, j_1) \mathcal{W}_{u_2;s_2}^{(J)}(i_2, j_1; k_2, j_2) \dots \mathcal{W}_{u_L;s_L}^{(J)}(i_L, j_{L-1}; k_L, j_L) = \frac{Z_l(J)}{Z_j(J)} q^{-\sum_{r=1}^J (r-1)b_r} \times \sum_{a_1 + \dots + a_r = j} q^{\sum_{r=1}^J (r-1)a_r} \left(\begin{array}{c} k_1 \quad k_L \\ \uparrow \quad \uparrow \\ a_J \text{ --- } w_{q^{J-1}u_1;s_1}^s \dots w_{q^{J-1}u_L;s_L}^s \text{ --- } b_J \\ \vdots \\ a_2 \text{ --- } w_{qu_1;s_1}^s \dots w_{qu_L;s_L}^s \text{ --- } b_2 \\ a_1 \text{ --- } w_{u_1;s_1}^s \dots w_{u_L;s_L}^s \text{ --- } b_1 \\ \vdots \\ i_1 \quad i_L \end{array} \right), \quad (2.6)$$

where we set $j_r = j + \sum_{m=1}^r i_m - \sum_{m=1}^r k_m$. Note that by the q -exchangeability we can also rewrite (2.5) as

$$\mathcal{W}_{u;s}^{(J)}(i, j; k, l) := \sum_{b_1 + \dots + b_L = l} \sum_{a_1 + \dots + a_J = j} \frac{q^{\sum_{r=1}^J (r-1)a_r}}{Z_j(J)} \left(\begin{array}{c} \begin{array}{c} \uparrow k \\ a_J \xrightarrow{w_{q^{J-1}u;s}^s} b_J \\ \vdots \\ a_2 \xrightarrow{w_{qu;s}^s} b_2 \\ a_1 \xrightarrow{w_{u;s}^s} b_1 \\ \downarrow i \end{array} \end{array} \right). \quad (2.7)$$

The resulting weights $\mathcal{W}_{u;s}^{(J)}$ can be given by an explicit expression obtained in [29], here we use a slightly different form of this expression given in [12, Theorem C.1.1] ⁽¹⁾ and originated from [16]:

$$\begin{aligned} \mathcal{W}_{u;s}^{(J)}(i, j; k, l) &= \mathbb{1}_{i+j=k+l} u^{l-j} q^{iJ} s^{j+l} \\ &\times \sum_{p=0}^{\min(j,k)} \phi(k-p, k+l-p; suq^J, su) \phi(p, j; sq^{-J}/u, q^{-J}), \end{aligned} \quad (2.8)$$

where

$$\phi(a, b; x, y) := (y/x)^a \frac{(x; q)_a (y/x; q)_{b-a}}{(y; q)_b} \frac{(q; q)_b}{(q; q)_a (q; q)_{b-a}}.$$


Actually, one can construct the weights $\mathcal{W}_{u;s}^{(J)}$ by fusing vertices with the weights \mathcal{R}_z , expressing $\mathcal{W}_{u;s}^{(J)}$ via a $J \times M$ rectangular grid of the vertices with weights $\mathcal{R}_{z_{i,j}}$ for a certain choice of parameters $z_{i,j}$ and specific boundary conditions. We are not using this construction for the current work; more details on this approach can be found in [8, Appendix].

2.3. Simplification of \mathcal{W} -weights

The weights $\mathcal{W}_{u;s}^{(J)}$ are rather complicated, but in this work we only use a particular “ q -Hahn” specialization of them: When $u = s$ the weights $\mathcal{W}_{u;s}^{(J)}$

⁽¹⁾ Our weights $\mathcal{W}_{u;s}^{(J)}(i, j; k, l)$ coincide with the extension $W_{J,M}(u/s; q; i\mathbf{e}_1, j\mathbf{e}_1, k\mathbf{e}_1, l\mathbf{e}_1)|_{q^M = s^{-2}}$ of the weights $W_{L,M}(x/y; q; \mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$ from [12, App. C].

simplify, with all but the $p = j$ term vanishing in (2.8), providing the following expression



$$l = W_{t,s}^s(i, j; k, l) := \mathcal{W}_{s;s}^{(J)}(i, j; k, l) \Big|_{q^J = t^{-2}}$$

$$= \mathbb{1}_{i+j=k+l} \mathbb{1}_{i \geq l} s^{2l} t^{-2l} \frac{(s^2/t^2; q)_{i-l} (t^2; q)_l}{(s^2; q)_i} \frac{(q; q)_i}{(q; q)_{i-l} (q; q)_l}. \quad (2.9)$$


Here we have used the fact that for fixed i, j, k, l the weight $\mathcal{W}_{u;s}^{(J)}(i, j; k, l)$ depends rationally on q^J for sufficiently large J , so we can introduce a new parameter t and replace q^J by t^{-2} .

Similarly to Remark 2.1, the weights $W_{t,s}^s$ defined above differ from the weights W_x from [15, (35)], and these two families of weights are related by

$$W_{s,t}^s(i, j; k, l) = (-s)^l \frac{(s^2; q)_k}{(q; q)_k} \frac{(q; q)_i}{(s^2; q)_i} W_{-s/t^2}(i, j; k, l), \quad (2.10)$$

where the global spin parameter from [15] is assumed to be equal to s .

The vertices with the weights $W_{t,s}^s$ satisfy a version of the Yang–Baxter equation: for every $a_1, a_2, a_3, b_1, b_2, b_3 \in \mathbb{Z}_{\geq 0}$ we have



$$(2.11)$$

where the tilted vertex is just a rotation by 45° of the vertex with the weights W_{t_1,t_2}^s . This Yang–Baxter equation is given in [15, Corollary 4.3]: it can be proved either by a direct computation, or using the Yang–Baxter equation (2.4) together with fusion.

Remark 2.2. — Apart from the Yang–Baxter equation, the weights $w_{u;s}^s$, $W_{t,s}^s$ and $\mathcal{W}_{u;s}^{(J)}$ satisfy another property, namely, they are *stochastic*: we have

$$\sum_{k,l} w_{u;s}^s(i, j; k, l) = 1, \quad \sum_{k,l} W_{t,s}^s(i, j; k, l) = 1, \quad \sum_{k,l} \mathcal{W}_{u;s}^{(J)}(i, j; k, l) = 1,$$

with fixed i, j in the appropriate label sets. The stochasticity of $w_{u;s}^s$ is proved by a direct computation, while for $W_{t,s}^s$ and $\mathcal{W}_{u;s}^{(J)}$ it follows from (2.7) and the stochasticity of $w_{u;s}^s$:

$$\begin{aligned} & \sum_{k,l} \mathcal{W}_{u;s}^{(J)}(i, j; k, l) \\ & := \sum_k \sum_{b_1, \dots, b_J \in \{0,1\}} \sum_{a_1 + \dots + a_J = j} \frac{q^{\sum_{r=1}^J (r-1)a_r}}{Z_j(J)} \left(\begin{array}{c} k \\ \uparrow \\ a_J \text{ --- } w_{q^{J-1}u;s}^s \text{ --- } b_J \\ \vdots \\ a_2 \text{ --- } w_{qu;s}^s \text{ --- } b_2 \\ a_1 \text{ --- } w_{u;s}^s \text{ --- } b_1 \\ \downarrow \\ i \end{array} \right) \\ & = \sum_{a_1 + \dots + a_J = j} \frac{q^{\sum_{r=1}^J (r-1)a_r}}{Z_j(J)} = 1, \end{aligned}$$

see [10, Section 5] for a more detailed explanation.

We use the weights $w_{u;s}^s$ and $W_{t,s}^s$ instead of $w_{u;s}$ and W_x from [4, 10, 11] because the deformations of the Yang–Baxter equation in Section 3 below look more natural for the stochastic weights.

2.4. Dual weights

Apart from vertices with edges oriented along the up-right direction we also use vertices with edges oriented along the up-left direction. We call them *dual vertices*, and their weights are closely related to the weights of the regular vertices. Such vertices play an important role for symmetric functions arising from vertex models, especially for the proofs of Cauchy type summation identities for such functions. Dual vertex weights were used in [15], and they are equivalent to the conjugated weights from [10]. Since our weights w^s and W^s differ from the weights used in those works, our

dual vertex weights also differ, though we do not state the exact matching between these weights; it can be readily recovered from the definitions below, Remark 2.1 and (2.10).

Define dual vertex weights $w_{u;s}^{s*}$ by the following relation:

$$\begin{aligned}
 \begin{array}{c} k \\ \uparrow \\ w_{u;s}^{s*} \\ \downarrow \\ i \\ \leftarrow j \quad \rightarrow l \end{array} &= w_{u;s}^{s*}(i, l; k, j) := s^{-2l} \frac{(q; q)_i (s^2; q)_k}{(s^2; q)_i (q; q)_k} w_{u;s}^s(k, j; i, l) \\
 &= s^{-2l} \frac{(q; q)_i (s^2; q)_k}{(s^2; q)_i (q; q)_k} j \begin{array}{c} i \\ \uparrow \\ w_{u;s}^s \\ \downarrow \\ k \\ \leftarrow \quad \rightarrow l \end{array}. \quad (2.12)
 \end{aligned}$$

The explicit values of weights $w_{u;s}^{s*}$ can be computed from (2.2) and they are summarized below:

$$\begin{array}{cccc}
 \begin{array}{c} g \\ \uparrow \\ w_{u;s}^{s*} \\ \downarrow \\ g \\ \leftarrow 0 \quad \rightarrow 0 \end{array} & \begin{array}{c} g+1 \\ \uparrow \\ w_{u;s}^{s*} \\ \downarrow \\ g \\ \leftarrow 0 \quad \rightarrow 1 \end{array} & \begin{array}{c} g-1 \\ \uparrow \\ w_{u;s}^{s*} \\ \downarrow \\ g \\ \leftarrow 1 \quad \rightarrow 0 \end{array} & \begin{array}{c} g \\ \uparrow \\ w_{u;s}^{s*} \\ \downarrow \\ g \\ \leftarrow 1 \quad \rightarrow 1 \end{array} \\
 \frac{1 - suq^g}{1 - su} & \frac{-s^{-1}u(1 - s^2q^g)}{1 - su} & \frac{1 - q^g}{1 - su} & \frac{q^g - s^{-1}u}{1 - su} \\
 & & & (2.13)
 \end{array}$$

As before, all unlisted configurations have weight 0, with the conservation law taking the form

$$w_{u;s}^{s*}(i, l; k, j) = 0, \quad \text{unless } i + l = k + j.$$

Looking at the explicit values, one can establish another relation between the weights $w_{u,s}^s$ and $w_{u,s}^{s*}$:

$$\begin{array}{c} k \\ \uparrow \\ \text{---} w_{u,s}^{s*} \text{---} \\ \downarrow \\ i \end{array} \begin{array}{c} l \\ \leftarrow \\ \text{---} \\ \rightarrow \\ (1-j) \end{array} = \frac{s-u}{s(1-us)} \begin{array}{c} k \\ \uparrow \\ \text{---} w_{u^{-1};s}^s \text{---} \\ \downarrow \\ i \end{array} \begin{array}{c} (1-l) \\ \leftarrow \\ \text{---} \\ \rightarrow \\ (1-l) \end{array} . \quad (2.14)$$

This is essentially the relation between the initial and conjugated weights in [10] that was used to prove the Cauchy identity there.

Using the definition (2.12), the dual weights $w_{u,s}^{s*}$ can be interpreted as rescaled $w_{u,s}^s$ weights reflected across the horizontal axis and with the direction of all edges reversed. Thus, one can rewrite the Yang–Baxter equation (2.4) in terms of the dual weights by reversing the orientation of all the edges, reflecting the whole vertex diagrams on both sides of (2.4) across the horizontal axis and then renormalizing the weights. It turns out that the factors arising from this rescaling actually cancel out⁽²⁾, and we arrive at the following Yang–Baxter equation:

$$\begin{array}{c} b_1 \\ \uparrow \\ \text{---} w_{y;s}^{s*} \text{---} \\ \downarrow \\ \text{---} w_{x;s}^{s*} \text{---} \\ a_1 \end{array} \begin{array}{c} b_2 \\ \leftarrow \\ \text{---} \\ \rightarrow \\ b_3 \end{array} \mathcal{R}_{x/y}^* = \begin{array}{c} b_1 \\ \uparrow \\ \text{---} w_{x;s}^{s*} \text{---} \\ \downarrow \\ \text{---} w_{y;s}^{s*} \text{---} \\ a_1 \end{array} \begin{array}{c} b_2 \\ \leftarrow \\ \text{---} \\ \rightarrow \\ b_3 \end{array} \quad (2.15)$$

where


$$\begin{array}{c} i \\ \swarrow \\ \mathcal{R}_z^* \\ \searrow \\ j \end{array} \begin{array}{c} l \\ \swarrow \\ \text{---} \\ \searrow \\ k \end{array} = \mathcal{R}_{z^{-1}}(i, j; k, l).$$

Alternatively, one can recover the same Yang–Baxter equation using (2.14).

There is also another Yang–Baxter equation that involves both $W_{t,s}^s$ and $w_{u,s}^{s*}$.


⁽²⁾ Cancellation of the terms $(-s)^{-2l}$ uses the conservation law of the tilted vertex.

PROPOSITION 2.3. — Assume that x, y, s, t are variables satisfying $xs = yt$. Then for any $a_1, b_1 \in \{0, 1\}$ and $a_2, b_2, a_3, b_3 \in \mathbb{Z}_{\geq 0}$ we have



where the tilted vertices are rotated by 45° vertices with the weights $w_{y;t}^{s*}$.

Proof. — There are three ways to prove Proposition 2.3. First, it can be verified by a direct computation, which reduces to four equalities between partition functions corresponding to the values of a_1, b_1 , with each partition function consisting of at most two summands. Alternatively, using (2.14), we can reduce the Yang–Baxter equation (2.16) to the equation




which follows from (2.4) by fusion. Finally, one can see (2.16) or, more precisely, (2.17) as a particular case of the Yang–Baxter equation for the weights $\mathcal{W}_{u;s}^{(J)}$:

$$\begin{aligned} & \sum_{l_1, l_2, l_3} \mathcal{W}_{y;s}^{(J)}(a_3, a_2; l_3, l_2) \mathcal{W}_{x;s}^{(I)}(l_3, a_1; b_3, l_1) \mathcal{W}_{xy^{-1}q^{-J/2}, q^{-J/2}}^{(I)}(l_2, l_1; b_2, b_1) \\ &= \sum_{l_1, l_2, l_3} \mathcal{W}_{xy^{-1}q^{-J/2}, q^{-J/2}}^{(I)}(a_2, a_1; l_2, l_1) \mathcal{W}_{x;s}^{(I)}(a_3, l_1; l_3, b_1) \mathcal{W}_{y;s}^{(J)}(l_3, l_2; b_3, b_2), \end{aligned}$$

which in turn can be seen as the $n = 1$ case of the master Yang–Baxter equation from [12, C.1.2]. \square

In this work we also use a dual analogue of the weights $W_{t,s}^s$. Define



$$\begin{aligned}
 & := s^{-2l} \frac{(q; q)_i (s^2; q)_k}{(s^2; q)_i (q; q)_k} \cdot t^{2k} \frac{(t^2; q)_j (q; q)_l}{(q; q)_j (t^2; q)_l} W_{t,s}^s(k, j; i, l) \\
 & = \mathbb{1}_{i+l=j+k} \mathbb{1}_{i \geq j} t^{2i-2j} \frac{(s^2/t^2; q)_{i-j} (t^2; q)_j}{(s^2; q)_i (q; q)_{i-j} (q; q)_j},
 \end{aligned} \tag{2.18}$$

where for the last computation we used (2.9) and the relation $i - j = k - l$ implied by $\mathbb{1}_{i+l=j+k}$.

3. Deformations of Yang–Baxter equations

The Yang–Baxter equations listed in the previous section have the same structure: they change positions of vertices while leaving the vertices themselves and their weights unchanged. In this section we introduce deformed Yang–Baxter equations, which change the weights of vertices in a nontrivial way.

We start with a straightforward deformation of (2.11).

PROPOSITION 3.1. — *We have the following equality of rational functions in t_1, t_2, t_3, η :*

$$\begin{aligned}
 & \sum_{l_1, l_2, l_3} W_{\eta t_2, \eta t_3}^s(a_3, a_2; l_3, l_2) W_{t_1, t_3}^s(l_3, a_1; b_3, l_1) W_{\eta t_1, \eta t_2}^s(l_2, l_1; b_2, b_1) \\
 & = \sum_{l_1, l_2, l_3} W_{t_1, t_2}^s(a_2, a_1; l_2, l_1) W_{\eta t_1, \eta t_3}^s(a_3, l_1; l_3, b_1) W_{t_2, t_3}^s(l_3, l_2; b_3, b_2),
 \end{aligned}$$

or, equivalently,

$$(3.1)$$

where the tilted vertex is a vertex with weights W^s , rotated by 45° .

Proof. — Note that

$$= \frac{(\eta^2 t^2; q)_l}{(t^2; q)_l} \frac{(s^2; q)_i}{(\eta^2 s^2; q)_i} j \text{ --- } l \cdot$$

Applying this equality to (3.1) and cancelling common factors corresponding to internal edges, we reduce (3.1) to (2.11) with both sides multiplied by

$$\frac{(\eta^2 t_1^2; q)_{b_1}}{(t_1^2; q)_{b_1}} \frac{(t_3^2; q)_{a_3}}{(\eta^2 t_3^2; q)_{a_3}}. \quad \square$$

There is also another possible deformation, which is crucial for the main results of this work. This time we deform the Yang–Baxter equation (2.16):

PROPOSITION 3.2. — *Let η, x, y, s, t be parameters satisfying the relation $xs = yt$. The following equality of rational functions holds:*

$$\begin{aligned} & \sum_{l_1, l_2, l_3} W_{t,s}^s(a_3, a_2; l_3, l_2) w_{\eta x; \eta s}^{s*}(l_3, l_1; b_3, a_1) w_{y;t}^{s*}(l_2, b_1; b_2, l_1) \\ &= \sum_{l_1, l_2, l_3} w_{\eta y; \eta t}^{s*}(a_2, l_1; l_2, a_1) w_{x;s}^{s*}(a_3, b_1; l_3, l_1) W_{t,s}^s(l_3, l_2; b_3, b_2), \end{aligned}$$

or, equivalently,

where the tilted vertex is a vertex with weights w^{s^*} , rotated by 45° .

The proof is based on the following observation: note that the weights $w_{u;s}^s(g, j; g+j-l, l)$ and $w_{u;s}^{s^*}(g, l; g+l-j, j)$ depend on g as rational functions in $\alpha = q^g$. This suggests that we can set q to be a generic non-zero complex number, extend the weights to get a model where certain edges have complex labels rather than integer ones, and then specialize these complex labels in a specific way to get a deformation of the initial model.

In more detail, assume that $q \in \mathbb{C} \setminus \{0\}$ and consider the vertex weights $w_{u;s}^{s^*}$. Using the explicit values (2.13) of these weights, we can define their extension denoted by

with $\alpha, \beta \in \mathbb{C} \setminus \{0\}$ being complex labels assigned to the vertical edges. The values of $\widehat{w}_{u;s}^{s^*}$ are given by

$$\begin{array}{cccc}
 \begin{array}{c} \alpha \\ \uparrow \\ \widehat{w}_{u;s}^{s^*} \\ \leftarrow 0 \quad 0 \\ \downarrow \\ \alpha \end{array} &
 \begin{array}{c} q\alpha \\ \uparrow \\ \widehat{w}_{u;s}^{s^*} \\ \leftarrow 0 \quad 1 \\ \downarrow \\ \alpha \end{array} &
 \begin{array}{c} q^{-1}\alpha \\ \uparrow \\ \widehat{w}_{u;s}^{s^*} \\ \leftarrow 1 \quad 0 \\ \downarrow \\ \alpha \end{array} &
 \begin{array}{c} \alpha \\ \uparrow \\ \widehat{w}_{u;s}^{s^*} \\ \leftarrow 1 \quad 1 \\ \downarrow \\ \alpha \end{array} \\
 \frac{1 - su\alpha}{1 - su} &
 \frac{-s^{-1}u(1 - s^2\alpha)}{1 - su} &
 \frac{1 - \alpha}{1 - su} &
 \frac{\alpha - s^{-1}u}{1 - su}
 \end{array} \tag{3.3}$$

and all other values are 0. One can immediately observe that for any $i, k \in \mathbb{Z}_{\geq 0}$ the weight $\widehat{w}_{u;s}^{s*}(q^i, l; q^k, j)$ is equal to $w_{u;s}^{s*}(i, l; k, j)$. Moreover, comparing (3.3) with (2.13) we can find a more general specialization, producing w^{s*} -weights with deformed parameters: For any complex η we have

$$\begin{array}{c} \eta^{2(1-j)} q^k \\ \uparrow \\ j \leftarrow \text{---} l \xrightarrow{\widehat{w}_{u;s}^{s*}} \\ \downarrow \\ \eta^{2(1-j)} q^i \end{array} = \frac{1 - \eta^2 s u}{1 - s u} \begin{array}{c} k \\ \uparrow \\ j \leftarrow \text{---} l \xrightarrow{w_{\eta u; \eta s}^{s*}} \\ \downarrow \\ i \end{array} . \quad (3.4)$$

Similarly, we can perform such an extension for the weights $W_{t,s}^s$. Here we have four edge labels which can be substituted by complex numbers, for the application below we modify the horizontal incoming and the vertical outgoing edges, while leaving the other two untouched. For $i, l \in \mathbb{Z}_{\geq 0}$ and $\alpha, \beta \in \mathbb{C} \setminus \{0\}$ define weights $\widehat{W}_{t,s}^s(i, \alpha; \beta, l)$ by setting

$$\begin{array}{c} q^{i-l} \alpha \\ \uparrow \\ \alpha \text{---} l \xrightarrow{\widehat{W}_{t,s}^s} \\ \downarrow \\ i \end{array} = \widehat{W}_{t,s}^s(i, \alpha; q^{i-l} \alpha, l) \\
 := \mathbb{1}_{i-l \geq 0} s^{2l} t^{-2l} \frac{(s^2/t^2; q)_{i-l} (t^2; q)_l}{(s^2; q)_i} \frac{(q; q)_i}{(q; q)_{i-l} (q; q)_l},$$

while $\widehat{W}_{t,s}^s(i, \alpha; \beta, l) = 0$ for all other choices of β . Note that these weight do not actually depend on α , so we can change α without affecting the weight.

Now we can apply the idea explained above to prove Proposition 3.2.

Proof of Proposition 3.2. — Since both sides are rational in η and q , it is enough to prove the equality when η, q are generic complex numbers. Fix $a_1, a_3, b_1, b_2 \in \mathbb{Z}_{\geq 0}$ and let $\Delta \in \mathbb{Z}$. Rewrite the Yang–Baxter equation (2.16)

as

$$(3.5)$$

where $\alpha = q^N$ for sufficiently large $N \in \mathbb{Z}_{\geq 0}$, and on both sides we are summing over all configurations of internal edges such that the vertex weights are nonzero.⁽³⁾ Note that both sides of (3.5) are rational functions in α , equal at infinitely many points, hence (3.5) holds for any α .

The proof is finished by (3.4) – multiplying both sides of (3.5) by $\frac{1-xs}{1-\eta^2 xs} = \frac{1-yt}{1-\eta^2 yt}$ and substituting $\alpha = \eta^{2(1-a_1)} q^{a_2}$ readily implies the claim. \square







Remark 3.3. — Proposition 3.1 can also be obtained in a manner similar to Proposition 3.2. In order to do that one needs to extend the weights $W_{t,s}^s$ with respect to another pair of labels

$$\alpha := \mathbb{1}_{k-j \geq 0} \alpha^{2 \log_q(s/t)}$$

$$\times \frac{(q\alpha; q)_{k-j} (s^2/t^2; q)_{k-j}}{(q; q)_{k-j}} \frac{(t^2; q)_{\infty}}{(\alpha t^2; q)_{\infty}} \frac{(q^{k-j} \alpha s^2; q)_{\infty}}{(s^2; q)_{\infty}},$$

⁽³⁾ There is still a finite number of such configurations, even with some edges now having complex labels.

table:

 $\frac{1 - sxq\sum_{r=1}^n I_r}{1 - sx}$	 $\frac{(s^2q^{I_i} - sx)q\sum_{r=i+1}^n I_r}{1 - sx}$	 $\frac{sx(q^{I_i} - 1)q\sum_{r=i+1}^n I_r}{1 - sx}$	(3.6)
 $\frac{1 - s^2q\sum_{r=1}^n I_r}{1 - sx}$	 $\frac{sx(q^{I_j} - 1)q\sum_{r=j+1}^n I_r}{1 - sx}$	 $\frac{s^2(q^{I_i} - 1)q\sum_{r=i+1}^n I_r}{1 - sx}$	

Here we assume that $i < j$ are integers between 1 and n , $\mathbf{I} = (I_1, \dots, I_n)$ denotes an n -tuple of nonnegative integers, \mathbf{e}_i denotes an n -tuple with 1 at the i th position and 0 elsewhere, and $\mathbf{0} = \mathbf{e}_0 := (0, \dots, 0)$. Note that these weights depend rationally on q^{I_c} for each $c \in \{1, \dots, n\}$, hence we can again substitute the labels of the thick edges by complex parameters. The resulting weights satisfy the following property:

$$\begin{array}{c}
 (T_{\eta^2, j} \mathbf{B}) \Big|_{\beta_c = q^{K_c}} \\
 \uparrow \widehat{w}_{u;s}^{\text{col}} \\
 \mathbf{e}_j \xrightarrow{\quad} \mathbf{e}_l = \mathbf{e}_j \xrightarrow{\quad} \mathbf{e}_l, \\
 \downarrow \\
 (T_{\eta^2, j} \mathbf{A}) \Big|_{\alpha_c = q^{I_c}}
 \end{array}
 \quad = \quad
 \begin{array}{c}
 \mathbf{K} \\
 \uparrow \\
 \mathbf{e}_j \xrightarrow{\quad} \mathbf{e}_l, \\
 \downarrow \\
 \mathbf{I}
 \end{array}
 ,$$

where $\mathbf{A} = (\alpha_1, \dots, \alpha_n)$ and $\mathbf{B} = (\beta_1, \dots, \beta_n)$ are tuples of non-zero complex numbers, and $T_{\eta^2, j}$ is the operator that multiplies the j th coordinate by η^2 and leaves the other coordinates unchanged. Using (2.14) one can readily see that for $n = 1$ the equality above is equivalent to (3.4).

Following the same notation, the colored analogue of the q -Hahn weights $W_{t,s}^s$ is given by, cf. [8, (6.17)],

$$\begin{array}{c}
 \begin{array}{c}
 \uparrow K \\
 | \\
 \text{---} W_{t,s}^{\text{col}} \text{---} \\
 | \\
 \downarrow I
 \end{array}
 \begin{array}{c}
 \leftarrow J \\
 \text{---} \\
 \rightarrow L
 \end{array}
 \end{array}
 = \mathbb{1}_{\mathbf{I}+\mathbf{J}=\mathbf{K}+\mathbf{L}} (s^2/t^2)^{|\mathbf{L}|} \frac{(s^2/t^2; q)_{|\mathbf{I}|-|\mathbf{L}|} (t^2; q)_{|\mathbf{L}|}}{(s^2; q)_{|\mathbf{I}|}}$$

$$\times \prod_{c=1}^n \mathbb{1}_{I_c \geq L_c} \frac{q^{L_{[1,c-1]}(I_c - L_c)}(q; q)_{I_c}}{(q; q)_{L_c} (q; q)_{I_c - L_c}},$$

where $|\mathbf{I}| := I_1 + \dots + I_n$ and $|\mathbf{L}| := L_1 + \dots + L_n$. Similarly to the $n = 1$ case, the colored versions of vertex weights above satisfy various forms of the Yang–Baxter equation. It turns out that we can deform some of these equations for the colored model using the same trick of replacing integer labels by complex ones. For instance, the colored analogue of (2.17) holds, and repeating the same argument as in Proposition 3.2 one can get the following deformed Yang–Baxter equation for $x/s = y/t$:

For $n = 1$ this deformed Yang–Baxter equation is equivalent to Proposition 3.2.

The deformed equation above can be used in the following way: Rotating the pictures on both sides of the equality by 90° , rescaling and considering

the limiting regime $s, x \rightarrow 0, x/s = y/t = \text{const}$, one gets

$$(3.7)$$

where the weights $W_{y;t}^{\mathcal{O}}$ are given by

$$= \mathbb{1}_{I+J=K+L} \frac{(t^2; q)_{|L|}}{(yt)_{|L|}} \times \prod_{c=1}^n \frac{q^{L_c(I_{[c+1,n]} - L_{[c+1,n]})} (q^{I_c - L_c + 1}; q)_{L_c}}{(q; q)_{L_c}},$$

while the weights $w_z^{\mathcal{O}}$ are certain degenerations of the weights $w_{u;s}^{\text{col}}$.

Now we can use a standard strategy: given a Yang–Baxter equation like (3.7), there is a way to produce a Cauchy-type summation identity by repeatedly applying the Yang–Baxter equation to vertices in row partition functions. Various examples of this procedure are described in [4, 10, 12, 15, 36], and in the case of varying parameters of the tilted crosses this is performed in Propositions 4.6 and 5.10 below. It turns out that this procedure applied to (3.7) results in a Cauchy-type summation identity of the form

$$\mathbb{E}_{\mathbf{u}|\Xi, \mathbf{S}}(\mathcal{O}) = f(\mathbf{u} \mid \Xi, \mathbf{S}),$$

where the expectation is taken with respect to a certain probability measure depending on parameters $\mathbf{u}, \Xi, \mathbf{S}$ and described in terms of the stochastic (inhomogeneous) colored higher spin six-vertex model, the observable \mathcal{O} is closely related to q -moments of the colored height function of the colored vertex model, and the functions f are partition functions of the colored vertex model with specific boundary conditions. This identity, in less generality, was first obtained in [14] in the case of the *homogeneous* colored vertex model, where the same strategy was used to produce a Cauchy-like summation identity starting from a non-deformed Yang–Baxter equation,

cf. [14, Theorem 4.7]. Our deformation allows to extend that result to the inhomogeneous case, answering [14, Remark 6.5].

However, there now exists a completely different approach used in [19] to achieve more general results about q -moments of the height functions; thus, we are not pursuing this topic here any further.

4. Row operators

In this section we introduce *row operators*, which form an algebraic framework for our results. The operators depend on one variable, κ or u depending on the operator, as well as two infinite sequences of nonzero complex inhomogeneity parameters $\Xi = (\xi_0, \xi_1, \xi_2, \dots)$ and $\mathbf{S} = (s_0, s_1, s_2, \dots)$, which are fixed throughout the section and are assumed to satisfy $s_{i+k}\xi_{i+k}s_i/\xi_i \neq 1$ for any $i, j \in \mathbb{Z}_{\geq 0}$.

In what follows, all non-tilted vertices in our diagrams will have one of the four types of weights: $w_{u;s}^{\mathbf{S}}, W_{t,s}^{\mathbf{S}}, w_{u;s}^{\mathbf{S}*}, W_{t,s}^{\mathbf{S}*}$; the pictorial notation for these vertices will always coincide with the pictures from (2.1), (2.9), (2.12) and (2.18) respectively. To simplify the figures below, we will interchangeably use the following equivalent notations

$$\begin{array}{c} k \\ \uparrow \\ j \text{---} \xrightarrow{w_{u;s}^{\mathbf{S}}} \text{---} l \\ \downarrow \\ i \end{array} = w^{\mathbf{S}} \left(\begin{array}{c} k \\ \uparrow \\ j \text{---} \xrightarrow{(u;s)} \text{---} l \\ \downarrow \\ i \end{array} \right) = \begin{array}{c} k \\ \uparrow \\ j \text{---} \xrightarrow{(u;s)} \text{---} l \\ \downarrow \\ i \end{array}$$

for the vertices with weights $w_{u;s}^{\mathbf{S}}$, and similarly for the other three types of weights. As the thickness and the direction of edges uniquely determine the weight family, this should not cause confusion.

4.1. Mixed shift

Before discussing the row operators, we would like to define a certain transformation, which is regularly applied to the sequences Ξ and \mathbf{S} throughout the further text. For an integer $k \in \mathbb{Z}_{\geq 0}$ and a pair of sequences $\mathcal{A} = (a_0, a_1, \dots)$, $\mathcal{B} = (b_0, b_1, \dots)$ such that $b_i \neq 0$ for any $i \in \mathbb{Z}_{\geq 0}$ introduce a *mixed shift* operator $\tau_{\mathcal{B}}$ by

$$\tau_{\mathcal{B}}^k \mathcal{A} = (a_0^{(k)}, a_1^{(k)}, \dots), \quad a_i^{(k)} = \sqrt{a_{i+k} b_{i+k} a_i / b_i}. \quad (4.1)$$

(t, s) is extensively used throughout the text, for convenience we rewrite (2.9) as

$$W^s \left(\begin{array}{c} \begin{array}{ccc} & k & \\ & \uparrow & \\ j & \text{---} & l \\ & \downarrow & \\ & i & \end{array} & \left(\sqrt{\frac{s\xi}{\kappa}}, s \right) \end{array} \right) = \mathbb{1}_{i+j=k+l} \mathbb{1}_{i \geq l} (\kappa s / \xi)^l \\
 \times \frac{(\kappa s / \xi; q)_{i-l} (\kappa^{-1} s \xi; q)_l}{(s^2; q)_i} \frac{(q; q)_i}{(q; q)_{i-l} (q; q)_l}. \quad (4.3)$$

We immediately note that the expression above depends polynomially on κ .

We also need to clarify what we mean by a partition function of the semi-infinite row. Let λ, μ be a pair of partitions with $l(\lambda), l(\mu) < L$ for some $L \in \mathbb{Z}_{\geq 0}$, and consider the partition function of the first L vertices of the row above, with the right outgoing edge having label 0. Then the L th vertex of this finite row has labels 0 around it and the weight W^s of this vertex is equal to 1, hence we can reduce L by 1 without changing the partition function. So, the partition function of the finite row consisting of the first L vertices does not depend on L as long as $l(\lambda), l(\mu) \leq L$, so we can drop L from the notation and work with the semi-infinite row, with all but finitely many vertices having weight 1. Note that the label of the horizontal edge between the $(L-1)$ st and the L th columns is also uniquely determined by the conservation law in columns $1, 2, \dots, L-1$, and the fact that it equals 0 is equivalent to the fact that $\lambda_1 = \mu_1 + a$.

The discussion above implies that all coefficients in (4.2) are finite products of weights W^s . By (4.3), all these weights are polynomials in κ , thus $\langle \lambda | \mathbb{T}_a(\kappa | \Xi, \mathbb{S}) | \mu \rangle$ can be viewed as a polynomial in κ . Moreover, note that by the conservation law in columns $1, 2, \dots, r-1$, the label of the horizontal edge immediately to the left of the r th column is uniquely determined and is equal to $\lambda_r - \mu_r$ (recall that $\lambda_1 - \mu_1 = a$). Hence, using vanishing of the weights $W_{t,s}^s(i, j; k, l)$ for $l > i$, we have

$$0 \leq \lambda_1 - \mu_1, \quad 0 \leq \lambda_r - \mu_r \leq \mu_{r-1} - \mu_r \quad \text{for } r \geq 2.$$

The condition above is equivalent to the *interlacing* $\lambda \succ \mu$ of partitions, that is, $\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots$.

So, we can sum up the discussion above as follows: $\langle \lambda | \mathbb{T}_a(\kappa | \Xi, \mathbb{S}) | \mu \rangle$, that is, the partition function from (4.2), depends polynomially on κ and satisfies

$$\langle \lambda | \mathbb{T}_a(\kappa | \Xi, \mathbb{S}) | \mu \rangle = 0 \quad \text{unless } \mu \prec \lambda \text{ and } \lambda_1 = \mu_1 + a. \quad (4.4)$$

$\tilde{B}^*(u \mid \Xi, \mathbf{S})$ ⁽⁴⁾ and set:

$$\mathbb{B}(\kappa \mid \Xi, \mathbf{S}) := \sum_{a \geq 0} (\kappa s_0 / \xi_0)^a \frac{(\kappa^{-1} s_0 \xi_0; q)_a}{(q; q)_a} \mathbb{T}_a(\kappa \mid \Xi, \mathbf{S}), \quad (4.7)$$

$$\mathbb{B}^*(\kappa \mid \Xi, \mathbf{S}) = \sum_{a \geq 0} \mathbb{T}_a^*(\kappa \mid \Xi, \mathbf{S}), \quad (4.8)$$

$$\tilde{B}^*(u \mid \Xi, \mathbf{S}) = \sum_{a=0}^1 (-u \xi_0 / s_0)^a T_a^*(u \mid \Xi, \mathbf{S}). \quad (4.9)$$

One can readily see that both combinations $\mathbb{B}^*(\kappa \mid \Xi, \mathbf{S})$ and $\tilde{B}^*(u \mid \Xi, \mathbf{S})$ are well-defined linear operators in $\text{End}(\mathbb{V})$, though the linear combination $\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ is not. In particular, compositions like $\tilde{B}^*(u \mid \Xi, \mathbf{S})\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ are not *a priori* well-defined. To remedy this issue, we treat $\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ as an infinite matrix in \mathbb{V} with respect to the basis $\{|\lambda\rangle\}_\lambda$ ⁽⁵⁾: in view of (4.4) we have

$$\langle \lambda | \mathbb{B}(\kappa \mid \Xi, \mathbf{S}) | \mu \rangle = (\kappa s_0 / \xi_0)^{\lambda_1 - \mu_1} \frac{(\kappa^{-1} s_0 \xi_0; q)_{\lambda_1 - \mu_1}}{(q; q)_{\lambda_1 - \mu_1}} \langle \lambda | \mathbb{T}_{\lambda_1 - \mu_1}(\kappa \mid \Xi, \mathbf{S}) | \mu \rangle. \quad (4.10)$$

Moreover, $\langle \lambda | \mathbb{B}(\kappa \mid \Xi, \mathbf{S}) | \mu \rangle = 0$ unless $\mu \prec \lambda$.

To justify compositions of such operators, consider a space $\widetilde{\text{End}}(\mathbb{V})$ consisting of infinite matrices M with respect to the basis $\{|\lambda\rangle\}_\lambda$, satisfying

$$\langle \lambda | M | \mu \rangle = 0 \quad \text{unless } l(\lambda) \leq l(\mu) + A \text{ and } \lambda_1 \geq \mu_1 - B,$$

for certain integers $A, B \in \mathbb{Z}_{\geq 0}$ depending on M . Then the usual matrix multiplication is well defined on the space $\widetilde{\text{End}}(\mathbb{V})$ ⁽⁶⁾, and $\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ can be considered as an element of $\widetilde{\text{End}}(\mathbb{V})$ with $(A, B) = (1, 0)$. Moreover, using the conservation law in a way similar to the derivation of (4.4), we have

$$\langle \lambda | \tilde{B}^*(u \mid \Xi, \mathbf{S}) | \mu \rangle = 0 \quad \text{unless } \lambda \prec \mu \text{ and } \mu_1 - \lambda_1 \in \{0, 1\}. \quad (4.11)$$

Thus, $\tilde{B}^*(u \mid \Xi, \mathbf{S})$ is also an element of $\widetilde{\text{End}}(\mathbb{V})$, with $(A, B) = (0, 1)$, so arbitrary compositions of operators $\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ and $\tilde{B}^*(u \mid \Xi, \mathbf{S})$ are well defined as elements of $\widetilde{\text{End}}(\mathbb{V})$.

The space $\widetilde{\text{End}}(\mathbb{V})$ provides only a partial resolution to the problem of interpreting $\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ as an operator, since in general the operator

⁽⁴⁾ Here we mostly follow the notation used in [15], but our notation $\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ corresponds to $\mathbb{C}(x)$ from [15].

⁽⁵⁾ so we will often call $\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ an operator

⁽⁶⁾ Any matrix element of the product can be computed as a sum over partitions whose Young diagrams fit inside a finite rectangle.

$\mathbb{B}^*(\kappa \mid \Xi, \mathbf{S})$ is not an element of $\widetilde{\text{End}}(\mathbb{V})$. Moreover, it turns out that the computation of the product $\mathbb{B}^*(\chi \mid \Xi, \mathbf{S})\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ in general requires infinite summations, cf. Remark 6.6 below. Since we are not using the operators $\mathbb{B}^*(\kappa \mid \Xi, \mathbf{S})$ for a large part of the text, we postpone this issue until Theorem 6.1, where we will treat it using convergence of formal power sums.

The linear combinations $\mathbb{B}(\kappa \mid \Xi, \mathbf{S})$ and $\mathbb{B}^*(\kappa \mid \Xi, \mathbf{S})$ are closely related to each other:

PROPOSITION 4.1. — *For any pair of partitions λ, μ , we have*

$$\langle \mu \mid \mathbb{B}^*(\kappa \mid \Xi, \mathbf{S}) \mid \lambda \rangle = \frac{(\tau_{\Xi} \mathbf{S})^{2\mu}}{\mathbf{S}^{2\lambda}} \frac{c_{\tau_{\Xi} \mathbf{S}}(\mu)}{c_{\mathbf{S}}(\lambda)} \langle \lambda \mid \mathbb{B}(\kappa \mid \tau_{\Xi} \Xi, \tau_{\Xi} \mathbf{S}) \mid \mu \rangle, \quad (4.12)$$

where $\tau_{\mathbf{S}}$ and τ_{Ξ} are mixed shift operators, and

$$c_{\mathbf{S}}(\lambda) := \prod_{i \geq 1} \frac{(s_i^2; q)_{\lambda_i - \lambda_{i+1}}}{(q; q)_{\lambda_i - \lambda_{i+1}}}, \quad \mathbf{S}^{2\lambda} := \prod_{i \geq 1} (s_{i-1})^{2\lambda_i}.$$

Proof. — For the duration of the proof, set

$$\tau_{\Xi} \Xi = (\xi'_0, \xi'_1, \xi'_2, \dots), \quad \tau_{\Xi} \mathbf{S} = (s'_0, s'_1, s'_2, \dots).$$

In particular, we have $\xi'_i s'_i = \xi_{i+1} s_{i+1}$ and $\xi'_i / s'_i = \xi_i / s_i$.

Recall that applying the conservation law to the row partition function $\langle \lambda \mid \mathbb{T}_a(\kappa \mid \Xi, \mathbf{S}) \mid \mu \rangle$ we can uniquely determine the labels of horizontal edges: the horizontal edge between columns $c - 1$ and c has label $\lambda_c - \mu_c$. Thus, using (4.10), we can write an explicit expression for $\mathbb{B}(\kappa \mid \tau_{\Xi} \Xi, \tau_{\Xi} \mathbf{S})$ in terms of the vertex weights W^s :

$$\begin{aligned} \langle \lambda \mid \mathbb{B}(\kappa \mid \tau_{\Xi} \Xi, \tau_{\Xi} \mathbf{S}) \mid \mu \rangle &= (\kappa s_0 / \xi_0)^{\lambda_1 - \mu_1} \frac{(\kappa^{-1} s'_0 \xi'_0; q)_{\lambda_1 - \mu_1}}{(q; q)_{\lambda_1 - \mu_1}} \\ &\times \prod_{c \geq 1} W^s_{\sqrt{s'_c \xi'_c / \kappa}, s'_c} (m_c(\mu'), \lambda_c - \mu_c; m_c(\lambda'), \lambda_{c+1} - \mu_{c+1}). \end{aligned} \quad (4.13)$$

Similarly, applying the same argument for the dual weights, we get

$$\begin{aligned} \langle \mu \mid \mathbb{B}^*(\kappa \mid \Xi, \mathbf{S}) \mid \lambda \rangle &= \left(\frac{\kappa}{s_0 \xi_0} \right)^{\lambda_1} \\ &\times \prod_{c \geq 1} W^{s*}_{\sqrt{s_c \xi_c / \kappa}, s_c} (m_c(\lambda'), \lambda_{c+1} - \mu_{c+1}; m_c(\mu'), \lambda_c - \mu_c). \end{aligned} \quad (4.14)$$

Now a direct computation using the explicit expressions from (2.9) and (2.18) shows that for any parameters t_1, t_2 and η we have

$$W_{t_1, t_2}^{s*}(i, l; k, j) = t_2^{-2l} \frac{(q; q)_i}{(t_2^2; q)_i} \frac{(\eta^2 t_2^2; q)_k}{(q; q)_k} \cdot t_1^{2k} \frac{(t_1^2; q)_j}{(q; q)_j} \frac{(q; q)_l}{(\eta^2 t_1^2; q)_l} W_{\eta t_1, \eta t_2}^s(k, j; i, l).$$

Substituting $t_1 = \sqrt{s_c \xi_c / \kappa}$, $t_2 = s_c$ and $\eta = \sqrt{\frac{s_{c+1} \xi_{c+1}}{s_c \xi_c}}$, we arrive at

$$\begin{aligned} W_{\sqrt{s_c \xi_c / \kappa}, s_c}^{s*}(i, l; k, j) &= s_c^{-2l} (\kappa^{-1} s_c \xi_c)^k \frac{(q; q)_i (s_c'^2; q)_k (\kappa^{-1} s_c \xi_c; q)_j}{(s_c^2; q)_i (q; q)_k (q; q)_j} \\ &\quad \times \frac{(q; q)_l}{(\kappa^{-1} s_c' \xi_c'; q)_l} W_{\sqrt{s_c' \xi_c' / \kappa}, s_c'}^s(k, j; i, l). \end{aligned}$$

Applying the equation above to (4.13), we see that all the Pochhammer symbols depending on j, l cancel out due to the relation $s_c' \xi_c' = s_{c+1} \xi_{c+1}$, so we get

$$\begin{aligned} \langle \lambda | \mathbb{B}(\kappa | \tau_S \Xi, \tau_\Xi S) | \mu \rangle &= \left(\frac{\kappa}{s_0 \xi_0} \right)^{\lambda_1} \frac{S^{2\lambda}}{(\tau_\Xi S)^{2\mu}} \frac{c_S(\lambda)}{c_{\tau_\Xi S}(\mu)} \\ &\quad \times \prod_{c \geq 1} W_{\sqrt{s_c \xi_c / \kappa}, s_c}^{s*}(m_c(\lambda'), \lambda_{c+1} - \mu_{c+1}; m_c(\mu'), \lambda_c - \mu_c), \end{aligned}$$

where we have used the relation

$$\frac{S^{2\lambda}}{(\tau_\Xi S)^{2\mu}} = (s_0 \xi_0)^{\mu_1} s_0^{2\lambda_1 - 2\mu_1} \prod_{c=1}^{\infty} s_c^{2(\lambda_{c+1} - \mu_{c+1})} \prod_{c=1}^{\infty} (s_c \xi_c)^{\mu_{c+1} - \mu_c}.$$

The proof is finished by comparing with (4.14). \square

Remark 4.2. — Up to a multiplication by a factor consisting of $(S)^{2\lambda}$ and $c_S(\lambda)$, our operators $\mathbb{B}(\kappa | \Xi, S)$, $\mathbb{B}^*(\kappa | \Xi, S)$ and $\tilde{B}(u | \Xi, S)$ degenerate to the operators $\mathbb{C}, \mathbb{B}^*, \tilde{B}$ from [15] when $S = (s, s, s, \dots)$ and $\Xi = (1, 1, 1, \dots)$. We also want to clarify that the usage of the letter T for row operators is not universally standard: for example, [10] and [15] use letters A, B, C, D for analogues of such operators. Partially to avoid confusion with this alternative notation we are using the tilde in the notation $\tilde{B}^*(u | \Xi, S)$. Another reason is the connection with stable spin Hall–Littlewood functions described in Section 5.2 below.

4.4. Fusion of the operators \tilde{B}^*

Here we prove that in some cases the operator $\mathbb{B}^*(\kappa | \Xi, S)$ can be obtained using operators $\tilde{B}^*(u | \Xi, S)$. This will only be used later as a step in the proof of Theorem 6.1.

PROPOSITION 4.3. — *For any $J \in \mathbb{Z}_{\geq 1}$ we have*

$$\tilde{B}^*(1 | S, S) \tilde{B}^*(q | S, S) \dots \tilde{B}^*(q^{J-1} | S, S) = \mathbb{B}^*(q^J | \bar{S}, S), \quad (4.15)$$

where

$$\bar{S} := (s_0^{-1}, s_1^{-1}, \dots).$$

where $b_1, \dots, b_J \in \{0, 1\}$ satisfy $\sum_{r=1}^J b_r = l$, and we have used the relation

$$Z_j(J) = (-1)^j q^{Jj} \frac{(q^{-J}; q)_j}{(q; q)_j}.$$

Let λ, μ be partitions and take $L > l(\lambda), l(\mu)$. Similarly to (2.6) we can fuse L dual columns by stacking (4.17) horizontally L times with the parameters s being equal to s_1, s_2, \dots, s_L . Setting the bottom (resp. top) boundary condition to a configuration corresponding to $|\lambda\rangle$ (resp. $\langle\mu|$), and assuming that all boundary edges on the right have labels equal to 0, we see that the parameter L becomes irrelevant and we can take $L \rightarrow \infty$ to get, using notations (4.5) and (4.6),

$$\begin{aligned} (-1)^j \langle\mu|\mathbb{T}_j^*(q^J | \bar{\mathbf{S}}, \mathbf{S})|\lambda\rangle &= \sum_{a_1 + \dots + a_J = j} q^{\sum_{r=1}^J (r-1)a_r} \\ &\times \langle\mu|T_{a_1}^*(1 | \mathbf{S}, \mathbf{S})T_{a_2}^*(q | \mathbf{S}, \mathbf{S}) \dots T_{a_J}^*(q^{J-1} | \mathbf{S}, \mathbf{S})|\lambda\rangle. \end{aligned} \quad (4.18)$$

Recall that

$$\begin{aligned} \mathbb{B}^*(q^J | \bar{\mathbf{S}}, \mathbf{S}) &= \sum_{j \geq 0} \mathbb{T}_j^*(q^J | \bar{\mathbf{S}}, \mathbf{S}), \\ \tilde{B}^*(q^{r-1} | \mathbf{S}, \mathbf{S}) &= \sum_{a_r \in \{0, 1\}} (-1)^{a_r} q^{(r-1)a_r} T_{a_r}^*(q^{r-1} | \mathbf{S}, \mathbf{S}), \quad r = 1, \dots, J, \end{aligned}$$

so the claim follows after multiplying (4.18) by $(-1)^j$ and summing over j . \square

4.5. Infinite 0th column

There is a useful interpretation of the linear combinations $\mathbb{B}(\kappa | \Xi, \mathbf{S})$ and $\tilde{B}^*(u | \Xi, \mathbf{S})$ in terms of the operators \mathbb{T}_a and T_a^* with an additional 0th column with ‘‘infinite’’ vertical labels. For a partition λ and an integer $N \geq \lambda_1$, let $\widehat{\lambda}^{(N)}$ denote the partition $(N, \lambda_1, \lambda_2, \dots)$, and set $\widehat{\Xi} = (1, \xi_0, \xi_1, \dots)$, $\widehat{\mathbf{S}} = (1, s_0, s_1, \dots)$.

PROPOSITION 4.4. — *For any partitions λ, μ and any $a \in \mathbb{Z}_{\geq 0}$, we have*

$$\langle\lambda|\mathbb{B}(\kappa | \Xi, \mathbf{S})|\mu\rangle = \frac{(s_0^2; q)_\infty}{(\kappa s_0 \xi_0^{-1}; q)_\infty} \lim_{N \rightarrow \infty} \langle\widehat{\lambda}^{(N+a)}|\mathbb{T}_a(\kappa | \widehat{\Xi}, \widehat{\mathbf{S}})|\widehat{\mu}^{(N)}\rangle,$$

where the convergence holds in the space of formal power series in q . Similarly, for any $i \in \{0, 1\}$ we have

$$\langle\mu|\tilde{B}^*(u | \Xi, \mathbf{S})|\lambda\rangle = (1 - u s_0 \xi_0) \lim_{N \rightarrow \infty} \langle\widehat{\mu}^{(N-i)}|T_i^*(u | \widehat{\Xi}, \widehat{\mathbf{S}})|\widehat{\lambda}^{(N)}\rangle,$$

where the convergence holds in the space of formal power series in q . In both cases the convergence also holds numerically, when $q \in \mathbb{C}$ satisfies $|q| < 1$ and $s_{i+k}\xi_{i+k}s_i/\xi_i \neq q^{-m}$ for any $i, k, m \in \mathbb{Z}_{\geq 0}$.

Proof. — We start with the first equality. Recall that from (4.10) we have

$$\langle \lambda | \mathbb{B}(\kappa | \Xi, \mathbf{S}) | \mu \rangle = (\kappa s_0 / \xi_0)^\Delta \frac{(\kappa^{-1} s_0 \xi_0; q)_\Delta}{(q; q)_\Delta} \langle \lambda | \mathbb{T}_\Delta(\kappa | \Xi, \mathbf{S}) | \mu \rangle,$$

where $\Delta = \lambda_1 - \mu_1$. On the other hand, separating the first vertex of the row partition function used to define the operator \mathbb{T}_a , we have

$$\langle \widehat{\lambda}^{(N+a)} | \mathbb{T}_a(\kappa | \widehat{\Xi}, \widehat{\mathbf{S}}) | \widehat{\mu}^{(N)} \rangle = W^s \left(\begin{array}{c} N+a-\lambda_1 \\ \uparrow \\ a \text{ --- } \text{---} \Delta \\ \downarrow \\ N-\mu_1 \end{array} \begin{array}{c} (\sqrt{\frac{\xi_0 s_0}{\kappa}}, s_0) \end{array} \right) \langle \lambda | \mathbb{T}_\Delta(\kappa | \Xi, \mathbf{S}) | \mu \rangle.$$

Hence, the claim follows from the limiting relation (cf. (4.3))

$$\begin{aligned} \lim_{N \rightarrow \infty} W^s \sqrt{\xi_0 s_0 / \kappa, s_0} (N - \mu_1, i; N - \lambda_1 + i, \Delta) \\ = (\kappa s_0 / \xi_0)^\Delta \frac{(\kappa s_0 / \xi_0; q)_\infty}{(s_0^2; q)_\infty} \frac{(\kappa^{-1} s_0 \xi_0; q)_\Delta}{(q; q)_\Delta}. \end{aligned}$$

The other equality follows similarly, using the limiting relation (cf. (2.13))

$$\lim_{N \rightarrow \infty} w_{u\xi_0; s_0}^{s*} (N - \lambda_1, i; N - \mu_1 - i, \Delta) = \frac{1}{1 - u\xi_0/s_0} (-u\xi_0/s_0)^\Delta$$

for $i, \Delta \in \{0, 1\}$. □

4.6. Exchange relations

The vertex weights we are using are distinguished by the Yang–Baxter equation, which makes the models with these weights integrable. In terms of row operators, the Yang–Baxter equation turns into so-called *exchange relations*. In this section we cover the exchange relations between operators $\mathbb{B}(\kappa | \Xi, \mathbf{S})$ and $\widetilde{B}^*(u | \Xi, \mathbf{S})$, which will result in various algebraic properties of the symmetric functions constructed in the next section. We start with relations resembling commutativity of operators.

PROPOSITION 4.5. — *The following relations hold:*

$$\begin{aligned} \mathbb{B}(\kappa_1 | \Xi, \mathbf{S}) \mathbb{B}(\kappa_2 | \tau_5 \Xi, \tau_\Xi \mathbf{S}) &= \mathbb{B}(\kappa_2 | \Xi, \mathbf{S}) \mathbb{B}(\kappa_1 | \tau_5 \Xi, \tau_\Xi \mathbf{S}), \\ \widetilde{B}^*(u_1 | \Xi, \mathbf{S}) \widetilde{B}^*(u_2 | \Xi, \mathbf{S}) &= \widetilde{B}^*(u_2 | \Xi, \mathbf{S}) \widetilde{B}^*(u_1 | \Xi, \mathbf{S}). \end{aligned} \tag{4.19}$$

Proof. — We start with the second relation. It suffices to prove that for any partitions μ, λ the following relation holds:

$$\langle \mu | \tilde{B}^*(u_1 | \Xi, S) \tilde{B}^*(u_2 | \Xi, S) | \lambda \rangle = \langle \mu | \tilde{B}^*(u_2 | \Xi, S) \tilde{B}^*(u_1 | \Xi, S) | \lambda \rangle.$$

Note that the tilted cross weights \mathcal{R}^* in the Yang–Baxter equation (2.15) depend only on the ratio x/y . Hence, after setting $s = s_r$, $x = u_1 \xi_r$, $y = u_2 \xi_r$, the Yang–Baxter equation (2.15) gives

where all non-tilted vertices have w^{s^*} -weights. Stacking L columns horizontally and iterating the Yang–Baxter equation above for $r = 1, \dots, L$, we can move the tilted cross $\mathcal{R}^*_{u_1/u_2}$ through L columns to get

Note that in the identity above we have also specified the left and right boundary conditions to be 0. In this case the conservation law forces the tilted crosses on both sides to be trivial of the form $\begin{smallmatrix} 0 & & 0 \\ & \times & \\ 0 & & 0 \end{smallmatrix}$ and have weight 1, hence for any partitions λ, μ such that $l(\lambda), l(\mu) < L$ we get

$$\begin{aligned} & \sum_{\nu} \langle \mu | T_0^*(u_2 | \Xi, S) | \nu \rangle \langle \nu | T_0^*(u_1 | \Xi, S) | \lambda \rangle \\ &= \sum_{\nu} \langle \mu | T_0^*(u_1 | \Xi, S) | \nu \rangle \langle \nu | T_0^*(u_2 | \Xi, S) | \lambda \rangle, \end{aligned}$$

with sums on both sides being finite. Since L was arbitrary, we can remove the restrictions on $l(\lambda)$ and $l(\mu)$. One finishes the proof by replacing $\Xi, S, \lambda, \mu, \nu$ by $\widehat{\Xi}, \widehat{S}, \widehat{\lambda}^{(N)}, \widehat{\mu}^{(N)}, \widehat{\nu}^{(N)}$ and applying Proposition 4.4.

To prove the first equation we will use a similar argument, but this time the tilted cross will change while we move it through the columns. Again, it suffices to prove

$$\langle \lambda | \mathbb{B}(\kappa_1 | \Xi, \mathbf{S}) \mathbb{B}(\kappa_2 | \tau_{\mathbf{S}} \Xi, \tau_{\Xi} \mathbf{S}) | \mu \rangle = \langle \lambda | \mathbb{B}(\kappa_2 | \Xi, \mathbf{S}) \mathbb{B}(\kappa_1 | \tau_{\mathbf{S}} \Xi, \tau_{\Xi} \mathbf{S}) | \mu \rangle.$$

This time we use the deformed Yang–Baxter equation from Proposition 3.1. Setting $t_1 = \sqrt{\frac{s_r \xi_r}{\kappa_1}}$, $t_2 = \sqrt{\frac{s_r \xi_r}{\kappa_2}}$, $t_3 = s_r$, $\eta = \sqrt{s_{r+1} \xi_{r+1}} / \sqrt{s_r \xi_r}$ in (3.1), we obtain


$$=$$

where all the vertices use W^s -weights with indicated parameters, and we set

$$\tau_{\mathbf{S}} \Xi = (\xi'_0, \xi'_1, \dots), \quad \tau_{\Xi} \mathbf{S} = (s'_0, s'_1, \dots).$$

Note that due to the equality $\xi'_r s'_r = \xi_{r+1} s_{r+1}$ the weights of the tilted cross on the left-hand side for $r = i$ coincide with the weights of the tilted cross on the right-hand side for $r = i + 1$. This suggests that we can stack columns with $r = 1, \dots, L$ horizontally, place a tilted cross with parameters $(\sqrt{\frac{s_{L+1} \xi_{L+1}}{\kappa_1}}, \sqrt{\frac{s_{L+1} \xi_{L+1}}{\kappa_2}})$ on the right, and iterate the equation above, moving the tilted cross and changing the parameters of it to $(\sqrt{\frac{s_1 \xi_1}{\kappa_1}}, \sqrt{\frac{s_1 \xi_1}{\kappa_2}})$. The resulting equation with 0 boundary conditions on the left and on the

right is given below:



Note that on both sides of the equation above the tilted cross must again be trivial with weight 1, hence for any partitions λ, μ such that $l(\lambda), l(\mu) < L$ we obtain

$$\langle \mu | \mathbb{T}_0(\kappa_1 | \Xi, \mathbb{S}) \mathbb{T}_0(\kappa_2 | \tau_{\mathbb{S}} \Xi, \tau_{\Xi} \mathbb{S}) | \lambda \rangle = \langle \mu | \mathbb{T}_0(\kappa_2 | \Xi, \mathbb{S}) \mathbb{T}_0(\kappa_1 | \tau_{\mathbb{S}} \Xi, \tau_{\Xi} \mathbb{S}) | \lambda \rangle.$$

Since L was arbitrary, we can remove the restrictions on $l(\lambda), l(\mu)$. The proof is then finished by replacing $\Xi, \mathbb{S}, \lambda, \mu$ with $\widehat{\Xi}, \widehat{\mathbb{S}}, \widehat{\lambda}^{(N)}, \widehat{\mu}^{(N)}$, multiplying both sides by

$$\frac{(s_0^2; q)_{\infty} (s_0 s_1 \xi_1 / \xi_0; q)_{\infty}}{(\kappa_1 s_0 / \xi_0; q)_{\infty} (\kappa_2 s_0 / \xi_0; q)_{\infty}},$$

and applying Proposition 4.4. \square

The next exchange relation involves operators $\mathbb{B}(\kappa | \Xi, \mathbb{S})$ and $\widetilde{B}^*(u | \Xi, \mathbb{S})$.

PROPOSITION 4.6. — *The following exchange relation holds:*

$$\widetilde{B}^*(u | \Xi, \mathbb{S}) \mathbb{B}(\kappa | \tau_{\mathbb{S}} \Xi, \tau_{\Xi} \mathbb{S}) = \frac{1 - u\kappa}{1 - u\xi_1 s_1} \mathbb{B}(\kappa | \tau_{\mathbb{S}} \Xi, \tau_{\Xi} \mathbb{S}) \widetilde{B}^*(u | \tau_{\mathbb{S}} \Xi, \tau_{\Xi} \mathbb{S}). \quad (4.20)$$

Proof. — The proof is similar to the proof of Proposition 4.5: we will take a suitable Yang–Baxter equation, iterate it for several columns, choose certain specific boundary conditions, and use Proposition 4.4. For the duration of the proof we again set


$$\tau_{\mathbb{S}} \Xi = (\xi'_0, \xi'_1, \dots), \quad \tau_{\Xi} \mathbb{S} = (s'_0, s'_1, \dots).$$

We start with the deformed Yang–Baxter equation from Proposition 3.2, where we set

$$s = s'_r = \sqrt{s_r s_{r+1} \xi_{r+1} / \xi_r}, \quad t = \sqrt{\frac{s_{r+1} \xi_{r+1}}{\kappa}},$$

$$x = u \xi'_r = u \sqrt{\xi_r \xi_{r+1} s_{r+1} / s_r}, \quad y = u \sqrt{\kappa \xi_{r+1} s_{r+1}}, \quad \eta = \sqrt{\frac{\xi_r s_r}{\xi_{r+1} s_{r+1}}}$$

to get



where \nearrow vertices have W^s -weights, and the remaining vertices have w^{s^*} -weights.

Note that the tilted cross in the left-hand side for $r = i$ coincides with the tilted cross in the right-hand side for $r = i + 1$, so we can stack several

We complete the proof by the following application of Proposition 4.4:

$$\begin{aligned}
 & \langle \lambda | \widetilde{B}^*(u | \Xi, \mathbf{S}) \mathbb{B}(\kappa | \tau_{\mathbf{S}} \Xi, \tau_{\Xi} \mathbf{S}) | \mu \rangle \\
 &= (1 - u \xi_0 s_0) \frac{(s'_0{}^2; q)_{\infty}}{(\kappa s'_0 / \xi'_0; q)_{\infty}} \lim_{N \rightarrow \infty} \langle \widehat{\lambda}^{(N-1)} | T_1^*(u | \widehat{\Xi}, \widehat{\mathbf{S}}) \mathbb{T}_0(\kappa | \tau_{\widehat{\mathbf{S}}} \widehat{\Xi}, \tau_{\widehat{\Xi}} \widehat{\mathbf{S}}) | \widehat{\mu}^{(N)} \rangle \\
 &= (1 - u \xi_0 s_0) \frac{1 - u \kappa}{1 - u \xi_0 s_0} \frac{(s'_0{}^2; q)_{\infty}}{(\kappa s'_0 / \xi'_0; q)_{\infty}} \\
 & \quad \times \lim_{N \rightarrow \infty} \langle \widehat{\lambda}^{(N-1)} | \mathbb{T}_0(\kappa | \tau_{\widehat{\mathbf{S}}} \widehat{\Xi}, \tau_{\widehat{\Xi}} \widehat{\mathbf{S}}) T_1^*(u | \tau_{\widehat{\mathbf{S}}} \widehat{\Xi}, \tau_{\widehat{\Xi}} \widehat{\mathbf{S}}) | \widehat{\mu}^{(N)} \rangle \\
 &= \frac{1 - u \kappa}{1 - u \xi_1 s_1} \langle \lambda | \mathbb{B}(\kappa | \tau_{\mathbf{S}} \Xi, \tau_{\Xi} \mathbf{S}) \widetilde{B}^*(u | \tau_{\mathbf{S}} \Xi, \tau_{\Xi} \mathbf{S}) | \mu \rangle,
 \end{aligned}$$

where we have used the relation $\xi'_0 s'_0 = \xi_1 s_1$. \square

5. Inhomogeneous spin q -Whittaker functions

In this section for any pair of partitions λ, μ we define a function $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n | \Xi, \mathbf{S})$ in variables $\kappa_1, \dots, \kappa_n$ as a partition function of a certain vertex model with vertices having W -weights. These functions are symmetric polynomials in $\kappa_1, \dots, \kappa_n$, and they depend on inhomogeneity parameters $\Xi = (\xi_0, \xi_1, \xi_2, \dots)$ and $\mathbf{S} = (s_0, s_1, s_2, \dots)$. We also prove several algebraic properties of these functions using the exchange relations from the previous section.

5.1. Vertex model construction

For a pair of partitions λ, μ with $l(\lambda), l(\mu) < L$ consider a grid of vertices consisting of n rows and L columns. For convenience, we number the rows from top to bottom, while the columns are numbered from the left to the right, starting from 1 in both cases. To a vertex at the intersection of the i th row and the j th column we assign the weight $W_{t,s}^{\xi}$ with parameters $t = \sqrt{\frac{\xi_{i+j} s_{i+j}}{\kappa_i}}$ and $s = \sqrt{s_{i+j} \xi_{i+j} s_j / \xi_j}$. Define boundary conditions as follows:

- the labels of all right edges are equal to 0;
- the left incoming edge in the i th row has label a_i ;
- the bottom edge in the j th column has label $m_j(\mu') = \mu_j - \mu_{j+1}$;
- the top edge in the j th column has label $m_j(\lambda') = \lambda_j - \lambda_{j+1}$.

The resulting partition function $ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n | \Xi, \mathbf{S})$ is depicted in Figure 5.1, where we use the notation $s_j^{(i)} := \sqrt{s_{i+j} \xi_{i+j} s_j / \xi_j}$.

Similarly to the row partition functions from the previous section, the conservation law forces all labels to the right of the $l(\lambda)$ th column in the partition function $ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ to be 0, with all vertices to the right of the $l(\lambda)$ th column having weight 1. Hence, the partition function does not actually depend on L and, taking $L \rightarrow \infty$, we can instead consider a semi-infinite grid, with n rows and an infinite number of columns.

Using the partition function $ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ we set

$$\mathbb{F}_{\lambda/\mu}^{\mathbf{s}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) := \sum_{a_1, \dots, a_n} \left(\prod_{i=1}^n (\kappa_i s_0 / \xi_0)^{a_i} \frac{(\kappa_i^{-1} s_i \xi_i; q)_{a_i}}{(q; q)_{a_i}} \right) \times ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}), \quad (5.1)$$

where the sum is over $(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n$ such that $\sum_{r=1}^n a_r = \lambda_1 - \mu_1$. Renormalizing, we define *inhomogeneous spin q -Whittaker* functions by

$$\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) := (-1)^{|\mu| - |\lambda|} \frac{(\tau_{\Xi}^n \mathbf{S})^\mu}{\mathbf{S}^\lambda} \frac{c_{\tau_{\Xi}^n \mathbf{S}}(\mu)}{c_{\mathbf{S}}(\lambda)} \mathbb{F}_{\lambda/\mu}^{\mathbf{s}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}), \quad (5.2)$$

where

$$c_{\mathbf{S}}(\lambda) := \prod_{i \geq 1} \frac{(s_i^2; q)_{\lambda_i - \lambda_{i+1}}}{(q; q)_{\lambda_i - \lambda_{i+1}}}, \quad \mathbf{S}^\lambda := \prod_{i \geq 1} s_i^{\lambda_i}.$$

In the particular case $\mu = \emptyset$ we also use the following notation⁽⁷⁾

$$\mathbb{F}_{\lambda}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) := \mathbb{F}_{\lambda/\emptyset}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}).$$

Remark 5.1. — When $\Xi = \xi^\infty := (\xi, \xi, \dots)$ and $\mathbf{S} = s^\infty := (s, s, \dots)$, the functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ turn into the spin q -Whittaker functions introduced in [15]. More precisely, if we let $\mathbb{F}_{\lambda/\mu}^{BW}(x_1, \dots, x_n)$ denote the functions from [15], then

$$\mathbb{F}_{\lambda/\mu}^{BW}(x_1, \dots, x_n) = \mathbb{F}_{\lambda/\mu}(-\xi x_1, \dots, -\xi x_n \mid \xi^\infty, s^\infty).$$

Remark 5.2. — The letter \mathbf{s} in the notation $\mathbb{F}_{\lambda/\mu}^{\mathbf{s}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ stands for “stochastic”; it is explained by the following identity:

$$\prod_{i=1}^n \frac{(\kappa_i s_0 \xi_0^{-1}; q)_\infty}{(s_i \xi_i s_0 / \xi_0; q)_\infty} \sum_{\lambda} \mathbb{F}_{\lambda}^{\mathbf{s}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) = 1.$$

This easily follows from Remark 2.2 in the following way. First we consider the partition function from Figure 5.1 with $\mu = \emptyset$. Since $W_{i,s}^{\mathbf{s}}(i, j, k, l) = 0$ unless $i \geq l$, any configuration with non-zero weight must have edge labels

⁽⁷⁾ Throughout the text we freely use λ instead of λ/\emptyset in similar contexts; this should not cause any confusion.




Figure 5.1. Partition function $ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$.

equal to 0 for all the edges to the right of the diagonal $i + j = n + 1$. Hence we can consider only the first n columns of the partition function, and instead of 0 boundary condition on the right we can take a sum over all possible labels of the outgoing edges on the right. Taking additionally a sum over possible boundary conditions on the top we end up with a partition function having a fixed incoming configuration and a “free” boundary condition for the outgoing edges. Applying the stochasticity of the weights $W_{t,s}^s$ consequently at each vertex starting from the top-right corner and going down and left, we get

$$\sum_{\lambda} ZW_{\lambda/\emptyset}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) = 1.$$

Hence,

$$\sum_{\lambda} \mathbb{F}_{\lambda}^{\mathbf{s}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) = \sum_{a_1, \dots, a_n \geq 0} \prod_{i=1}^n (\kappa_i s_0 / \xi_0)^{a_i} \frac{(\kappa_i^{-1} s_i \xi_i; q)_{a_i}}{(q; q)_{a_i}}$$

and the identity follows from the q -binomial theorem

$$\sum_{a \geq 0} x^a \frac{(y; q)_a}{(q; q)_a} = \frac{(xy; q)_{\infty}}{(x; q)_{\infty}}.$$




Figure 5.2. Partition function $Zw_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathcal{S})$.

5.2. Inhomogeneous stable spin Hall–Littlewood functions

It is also useful to consider another family of functions that straightforwardly generalizes both inhomogeneous spin Hall–Littlewood functions from [10] and stable Hall–Littlewood functions from [23] and [15].

Let u_1, \dots, u_n be a collection of variables. Similarly to the construction of the functions $\mathbb{F}_{\lambda/\mu}$ above, consider a grid of vertices consisting of n rows and L columns, with the vertex at the intersection of the i th row and the j th column having the w^s -weight with parameters $(u_i \xi_j; s_j)$. For a pair of partitions λ, μ with $l(\lambda), l(\mu) < L$ define boundary conditions in the same way as before:

- the labels of all right edges are equal to 0;
- the left incoming edge in the i th row has label a_i for some $a_i \in \{0, 1\}$;
- the bottom edge in the j th column has label $m_j(\mu') = \mu_j - \mu_{j+1}$;
- the top edge in the j th column has label $m_j(\lambda') = \lambda_j - \lambda_{j+1}$.

The resulting partition function is denoted by $Zw_{\lambda/\mu}^{(a_1, \dots, a_n)}(u_1, \dots, u_n \mid \Xi, \mathcal{S})$, and its depiction is given in Figure 5.2.

The *inhomogeneous stable spin Hall–Littlewood functions* $\widetilde{F}_{\lambda/\mu}(u_1, \dots, u_n \mid \Xi, \mathbf{S})$ ⁽⁸⁾ are defined by

$$\begin{aligned} & \widetilde{F}_{\lambda'/\mu'}^s(u_1, \dots, u_n \mid \Xi, \mathbf{S}) \\ & := \sum_{a_1, \dots, a_n \in \{0,1\}} \prod_{i=1}^n (-u \xi_0 s_0)^{a_i} Z w_{\lambda'/\mu'}^{(a_1, \dots, a_n)}(u_1, \dots, u_n \mid \Xi, \mathbf{S}), \\ & \widetilde{F}_{\lambda'/\mu'}(u_1, \dots, u_n \mid \Xi, \mathbf{S}) \\ & := (-1)^{|\mu| - |\lambda|} \frac{S^\mu c_{\mathbf{S}}(\mu)}{S^\lambda c_{\mathbf{S}}(\lambda)} \widetilde{F}_{\lambda'/\mu'}^s(u_1, \dots, u_n \mid \Xi, \mathbf{S}). \end{aligned} \tag{5.3}$$

Note that we use conjugated partitions λ' and μ' : this is needed to obtain usual Hall–Littlewood polynomials as a degeneration of $\widetilde{F}_{\lambda/\mu}$, see [15]. Also note that the parameter s_0 is actually artificial: both $\widetilde{F}_{\lambda'/\mu'}^s(u_1, \dots, u_n \mid \Xi, \mathbf{S})$ and S^λ/S^μ have factor $s_0^{\lambda_1 - \mu_1}$ with no other dependence on s_0 , thus $\widetilde{F}_{\lambda'/\mu'}(u_1, \dots, u_n \mid \Xi, \mathbf{S})$ does not depend on s_0 at all.

For $\Xi = 1^\infty$ and $\mathbf{S} = s^\infty$ the functions $\widetilde{F}_{\lambda/\mu}(u_1, \dots, u_n \mid \Xi, \mathbf{S})$ degenerate to the stable spin Hall–Littlewood functions used in [15] and originating from [23]. Actually, the stability property holds for the inhomogeneous analogue as well: we have

$$\widetilde{F}_{\lambda/\mu}(u_1, \dots, u_{n-1}, 0 \mid \Xi, \mathbf{S}) = \widetilde{F}_{\lambda/\mu}(u_1, \dots, u_{n-1} \mid \Xi, \mathbf{S}),$$

which can be readily verified using the expression for the partition function $Z w_{\lambda'/\mu'}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ and the explicit values of weights $w_{u,s}^s$ for $u = 0$.

At the same time, the functions $\widetilde{F}_{\lambda/\mu}(u_1, \dots, u_n \mid \Xi, \mathbf{S})$ are closely related to the inhomogeneous spin Hall–Littlewood functions F defined in [10, Definition 4.4]. Actually, we already have those functions in our setting: one has

$$\begin{aligned} & (-1)^{|\mu| - |\lambda|} \frac{S^\mu c_{\widehat{\mathbf{S}}}(\widehat{\mu}^{(N)})}{S^\lambda c_{\widehat{\mathbf{S}}}(\widehat{\lambda}^{(N+n)})} Z w_{\widehat{\lambda}^{(N+n)}/\widehat{\mu}^{(N)}}^{(1,1,\dots,1)}(u_1, \dots, u_n \mid \widehat{\Xi}, \widehat{\mathbf{S}}) \\ & = F_{\lambda'/\mu'}(u_1, \dots, u_n \mid \Xi, \mathbf{S}), \end{aligned}$$

since both sides are defined using the same partition function, up to a renormalization of the weights mentioned in Remark 2.1 and resulting in the prefactor on the left-hand side. Here $N \in \mathbb{Z}_{\geq 1}$ is a sufficiently large integer, which implicitly enters the right-hand side via the index λ'/μ' of the function F : In [10] partitions are replaced by signatures, that is, nonincreasing

⁽⁸⁾ We follow the notation $\widetilde{F}_{\lambda/\mu}$ to be consistent with [15] and to differentiate from the non-stable version of these functions, which is commonly denoted by $F_{\lambda/\mu}$.

integer sequences, and in the right-hand side we treat λ' and μ' as signatures of lengths $N + n$ and N , respectively, by adding an appropriate number of 0's at the end. We also use the notation from Proposition 4.4 to shift the sequences of parameters Ξ and \mathbf{S} to the right, so that the partition function from Figure 5.2 starts with the column having parameters $(u_i \xi_0, s_0)$.

Moreover, it turns out that one can interpret the functions $\tilde{F}_{\lambda/\mu}(u_1, \dots, u_n \mid \Xi, \mathbf{S})$ as a special case of the inhomogeneous spin Hall–Littlewood functions $F_{\lambda/\mu}$. To see it, note that

$$\left(\begin{array}{c} \begin{array}{ccc} & k & \\ & \uparrow & \\ (s_0)^{-l} & 1 & \xrightarrow{(u\xi_0, s_0)} l \\ & \downarrow & \\ & i & \end{array} \\ \end{array} \right) \Big|_{s_0 \rightarrow 0} = (-u\xi_0 s_0)^l \Big|_{s_0 \rightarrow 1}.$$

Applying this to the 0th column of the partition function $Zw_{\hat{\lambda}^{(N+n)}/\hat{\mu}^{(N)}}^{(1,1,\dots,1)}(u_1, \dots, u_n \mid \hat{\Xi}, \hat{\mathbf{S}})$, one can rewrite (5.3) as

$$\begin{aligned} \left(s_0^{\mu_1 - \lambda_1} Zw_{\hat{\lambda}^{(N+n)}/\hat{\mu}^{(N)}}^{(1,1,\dots,1)}(u_1, \dots, u_n \mid \hat{\Xi}, \hat{\mathbf{S}}) \right) \Big|_{s_0 \rightarrow 0} \\ = \tilde{F}_{\lambda'/\mu'}^{\mathbf{S}}(u_1, \dots, u_n \mid \Xi, \mathbf{S}) \Big|_{s_0 \rightarrow 1} \end{aligned}$$

or, using the relation

$$\frac{(q; q)_{N - \mu_1}}{(q; q)_{N + n - \lambda_1}} \left(s_0^{\lambda_1 - \mu_1} \frac{S^\mu}{S^\lambda} \frac{c_{\hat{\mathbf{S}}}(\hat{\mu}^{(N)})}}{c_{\hat{\mathbf{S}}}(\hat{\lambda}^{(N+n)})} \right) \Big|_{s_0 \rightarrow 0} = \left(\frac{S^\mu}{S^\lambda} \frac{c_{\mathbf{S}}(\mu)}{c_{\mathbf{S}}(\lambda)} \right) \Big|_{s_0 \rightarrow 1},$$

one can renormalize everything to get

$$\frac{(q; q)_{N - \mu_1}}{(q; q)_{N + n - \lambda_1}} F_{\lambda'/\mu'}(u_1, \dots, u_n \mid \Xi, \mathbf{S}) \Big|_{s_0 = 0} = \tilde{F}_{\lambda'/\mu'}(u_1, \dots, u_n \mid \Xi, \mathbf{S}),$$

where in the left-hand side we again implicitly treat λ' and μ' as signatures of lengths $N + n$ and N respectively, while the right-hand side does not depend on s_0 .

The observation above allows us to immediately translate certain results from [10] to our setting. For example, we immediately see that the functions $\tilde{F}_\lambda(u_1, \dots, u_n \mid \Xi, \mathbf{S})$ are symmetric in u_1, \dots, u_n since the same result holds for the functions $F_{\lambda/\mu}(u_1, \dots, u_n \mid \Xi, \mathbf{S})$, see [10, Proposition 4.5]. We also have an explicit formula in the case $\mu = \emptyset$, which is similar to [10,

Theorem 4.14] and [23, Remark 2]:

$$\begin{aligned} & \widetilde{F}_\lambda(u_1, \dots, u_n \mid \Xi, \mathbb{S}) \\ &= \frac{1}{(q; q)_{n-l(\lambda)}} \sum_{\sigma \in \mathbb{S}_n} \sigma \left(\prod_{1 \leq \alpha < \beta \leq n} \frac{u_\alpha - qu_\beta}{u_\alpha - u_\beta} \prod_{i=1}^n \widetilde{\varphi}_{\lambda_i}(u_i \mid \Xi, \mathbb{S}) \right), \end{aligned}$$

where the sum is over permutations of n elements that act on functions in u_1, \dots, u_n by permuting variables, and for $k \geq 1$ we set

$$\widetilde{\varphi}_0(u \mid \Xi, \mathbb{S}) = 1 - q, \quad \widetilde{\varphi}_k(u \mid \Xi, \mathbb{S}) = \frac{\xi_0 u(1 - q)}{1 - s_k \xi_k u} \prod_{j=1}^{k-1} \frac{\xi_j u - s_j}{1 - s_j \xi_j u}.$$

5.3. Row operator representations

To prove algebraic properties of the functions $\mathbb{F}_{\lambda/\mu}$, instead of using the whole defining partition functions, it is more convenient to construct them row by row using the row operators introduced earlier.

PROPOSITION 5.3. — *For any partitions λ, μ we have*

$$\begin{aligned} & \mathbb{F}_{\lambda/\mu}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \\ &= \langle \lambda \mid \mathbb{B}(\kappa_1 \mid \tau_{\Xi} \Xi, \tau_{\Xi} \mathbb{S}) \mathbb{B}(\kappa_2 \mid \tau_{\Xi}^2 \Xi, \tau_{\Xi}^2 \mathbb{S}) \dots \mathbb{B}(\kappa_n \mid \tau_{\Xi}^n \Xi, \tau_{\Xi}^n \mathbb{S}) \mid \mu \rangle \\ &= \frac{\mathbb{S}^{2\lambda}}{(\tau_{\Xi} \mathbb{S})^{2\mu}} \frac{c_{\mathbb{S}}(\lambda)}{c_{\tau_{\Xi} \mathbb{S}}(\mu)} \\ & \quad \times \langle \mu \mid \mathbb{B}^*(\kappa_n \mid \tau_{\Xi}^{n-1} \Xi, \tau_{\Xi}^{n-1} \mathbb{S}) \dots \mathbb{B}^*(\kappa_2 \mid \tau_{\Xi} \Xi, \tau_{\Xi} \mathbb{S}) \mathbb{B}^*(\kappa_1 \mid \Xi, \mathbb{S}) \mid \lambda \rangle, \end{aligned} \quad (5.4)$$

where τ_{Ξ} and $\tau_{\mathbb{S}}$ are the mixed shift operators defined by (4.1).

Proof. — The second equality follows at once from Proposition 4.1, so we focus on the first equality.

Recall that

$$\tau_{\Xi}^i \Xi = (\xi_0^{(i)}, \xi_1^{(i)}, \dots), \quad \tau_{\Xi}^i \mathbb{S} = (s_0^{(i)}, s_1^{(i)}, \dots),$$

where $s_j^{(i)} = \sqrt{s_{i+j} \xi_{i+j} s_j / \xi_j}$, and $\xi_j^{(i)}$ satisfy $\xi_j^{(i)} s_j^{(i)} = \xi_{i+j} s_{i+j}$. Thus, the parameters of the (i, j) th vertex in the partition function $ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S})$ coincide with the parameters of the j th vertex in the row of vertices used to define $\mathbb{T}_{a_i}(\kappa_i \mid \tau_{\Xi}^i \Xi, \tau_{\Xi}^i \mathbb{S})$, or, equivalently, we have

$$\begin{aligned} & ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \\ &= \langle \lambda \mid \mathbb{T}_{a_1}(\kappa_1 \mid \tau_{\Xi} \Xi, \tau_{\Xi} \mathbb{S}) \dots \mathbb{T}_{a_n}(\kappa_n \mid \tau_{\Xi}^n \Xi, \tau_{\Xi}^n \mathbb{S}) \mid \mu \rangle. \end{aligned}$$

On the other hand, using relations $s_0^{(i)} \xi_0^{(i)} = s_i \xi_i$ and $s_0^{(i)} / \xi_0^{(i)} = s_0 / \xi_0$, we see that the coefficients in (5.1) coincide with the coefficients in (4.7), concluding the proof. \square

As a corollary we observe that the functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S})$ are polynomials in $\kappa_1, \dots, \kappa_n$. Indeed, matrix coefficients of the operator $\mathbb{C}(\kappa \mid \Xi, \mathbb{S})$ are polynomials in κ , so Proposition 5.3 implies polynomiality.

PROPOSITION 5.4. — *For any partitions λ, μ we have*

$$\begin{aligned} \widetilde{\mathbb{F}}_{\lambda'/\mu'}^{\mathbb{S}}(u_1, \dots, u_n \mid \Xi, \mathbb{S}) \\ = \frac{\mathbb{S}^{2\lambda} \mathbf{c}_{\mathbb{S}}(\lambda)}{\mathbb{S}^{2\mu} \mathbf{c}_{\mathbb{S}}(\mu)} \langle \mu \mid \widetilde{B}^*(u_n \mid \Xi, \mathbb{S}) \dots \widetilde{B}^*(u_2 \mid \Xi, \mathbb{S}) \widetilde{B}^*(u_1 \mid \Xi, \mathbb{S}) \mid \lambda \rangle. \end{aligned} \quad (5.5)$$

Proof. — Similarly to the proof of (5.3), we consider the partition function $Zw_{\lambda/\mu}^{(a_1, \dots, a_n)}(u_1, \dots, u_n \mid \Xi, \mathbb{S})$ row by row, interpreting it as a matrix coefficient of a composition of row operators. In addition, here we will change the weights $w^{\mathbb{S}}$ to the weights $w^{\mathbb{S}^*}$ using (2.12). We have already used this change of weights for a single column during the discussion of fusion of the row operators $\widetilde{B}^*(u \mid \Xi, \mathbb{S})$, see (4.16).

Noting that for the partition function $Zw_{\lambda/\mu}^{(a_1, \dots, a_n)}(u_1, \dots, u_n \mid \Xi, \mathbb{S})$ the sum of all labels of edges between columns $L-1$ and L is equal to $\lambda_L - \mu_L$, we can use (4.16) for each column in $Zw_{\lambda/\mu}^{(a_1, \dots, a_n)}(u_1, \dots, u_n \mid \Xi, \mathbb{S})$ to get

$$\begin{aligned} Zw_{\lambda/\mu}^{(a_1, \dots, a_n)}(u_1, \dots, u_n \mid \Xi, \mathbb{S}) \\ = \frac{s_0^{-\lambda_1} \mathbb{S}^{2\lambda} \mathbf{c}_{\mathbb{S}}(\lambda)}{s_0^{-\mu_1} \mathbb{S}^{2\mu} \mathbf{c}_{\mathbb{S}}(\mu)} \langle \mu \mid T_{a_n}^*(u_n \mid \Xi, \mathbb{S}) \dots T_{a_2}^*(u_2 \mid \Xi, \mathbb{S}) T_{a_1}^*(u_1 \mid \Xi, \mathbb{S}) \mid \lambda \rangle. \end{aligned}$$

The claim follows by comparing coefficients in (4.9) and (5.3). \square

5.4. Properties of the inhomogeneous spin q -Whittaker functions

So far we have been establishing an algebraic framework to analyse the functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S})$. Here we finally use it to prove our main results about these functions.

PROPOSITION 5.5 (Branching rule). — *For any partitions λ, μ and $1 \leq m < n$ we have*

$$\begin{aligned} \mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \\ = \sum_{\nu} \mathbb{F}_{\lambda/\nu}(\kappa_1, \dots, \kappa_m \mid \Xi, \mathbb{S}) \mathbb{F}_{\nu/\mu}(\kappa_{m+1}, \dots, \kappa_n \mid \tau_{\mathbb{S}}^m \Xi, \tau_{\Xi}^m \mathbb{S}), \end{aligned} \quad (5.6)$$

where the sum on the right-hand side is finite.

Proof. — A similar claim for the functions $\mathbb{F}_{\lambda/\mu}^s$ follows at once from Proposition 5.3 by an insertion of the summation $\sum_{\nu} |\nu\rangle\langle\nu| = \text{id}$:

$$\begin{aligned} \mathbb{F}_{\lambda/\mu}^s(\kappa_1, \dots, \kappa_n \mid \Xi, S) &= \langle \lambda \mid \mathbb{B}(\kappa_1 \mid \tau_S \Xi \tau_{\Xi} S) \dots \mathbb{B}(\kappa_n \mid \tau_S^n \Xi, \tau_{\Xi}^n S) \mid \mu \rangle \\ &= \sum_{\nu} \langle \lambda \mid \mathbb{B}(\kappa_1 \mid \tau_S \Xi, \tau_{\Xi} S) \dots \mathbb{B}(\kappa_m \mid \tau_S^m \Xi, \tau_{\Xi}^m S) \mid \nu \rangle \\ &\quad \times \langle \nu \mid \mathbb{B}(\kappa_{m+1} \mid \tau_S^{m+1} \Xi, \tau_{\Xi}^{m+1} S) \dots \mathbb{B}(\kappa_n \mid \tau_S^n \Xi, \tau_{\Xi}^n S) \mid \mu \rangle \\ &= \sum_{\nu} \mathbb{F}_{\lambda/\nu}^s(\kappa_1, \dots, \kappa_m \mid \Xi, S) \mathbb{F}_{\nu/\mu}^s(\kappa_{m+1}, \dots, \kappa_n \mid \tau_S^m \Xi, \tau_{\Xi}^m S). \end{aligned}$$

To reach (5.6), one multiplies the above equality by

$$(-1)^{|\mu| - |\lambda|} \frac{(\tau_{\Xi}^n S)^{\mu} c_{\tau_{\Xi}^n S}(\mu)}{S^{\lambda} c_S(\lambda)} = (-1)^{|\mu| - |\lambda|} \frac{(\tau_{\Xi}^n S)^{\mu} c_{\tau_{\Xi}^n S}(\mu)}{(\tau_{\Xi}^m S)^{\nu} c_{\tau_{\Xi}^m S}(\nu)} \frac{(\tau_{\Xi}^m S)^{\nu} c_{\tau_{\Xi}^m S}(\nu)}{S^{\lambda} c_S(\lambda)}. \quad \square$$

PROPOSITION 5.6. — *We have*

$$\mathbb{F}_{\lambda/\mu}(\kappa \mid \Xi, S) = \begin{cases} (-\kappa)^{|\lambda| - |\mu|} \prod_{i \geq 1} \xi_i^{\mu_i - \lambda_i} \left(\frac{s_i \xi_i}{s_{i-1} \xi_{i-1}} \right)^{\mu_i / 2} \\ \quad \times \frac{(\kappa^{-1} s_i \xi_i; q)_{\lambda_i - \mu_i} (\kappa s_i / \xi_i; q)_{\mu_i - \lambda_{i+1}} (q; q)_{\lambda_i - \lambda_{i+1}}}{(q; q)_{\lambda_i - \mu_i} (q; q)_{\mu_i - \lambda_{i+1}} (s_i^2; q)_{\lambda_i - \lambda_{i+1}}} & \text{if } \mu \prec \lambda, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. — This is a result of an explicit computation consisting of the following steps: we consider an expression of $\langle \lambda \mid \mathbb{C}(\kappa \mid \tau_S \Xi, \tau_{\Xi} S) \mid \mu \rangle$ as a product of the vertex weights W^s , which is done in (4.13) by using the conservation law, then plug in the explicit values of the weights W^s from (4.3), and finally renormalize the resulting functions $\mathbb{F}_{\lambda/\mu}^s(\kappa \mid \Xi, S)$ to obtain the functions $\mathbb{F}_{\lambda/\mu}(\kappa \mid \Xi, S)$ using (5.2). \square

Note that the square roots $\left(\frac{s_i \xi_i}{s_{i-1} \xi_{i-1}} \right)^{\mu_i / 2}$ in the expression from Proposition 5.6 are due to the renormalization (5.2), which involves square roots.

COROLLARY 5.7. — *For a pair of partitions λ, μ the function $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, S)$ vanishes unless for any $r \geq 1$ we have*

$$\mu'_r \leq \lambda'_r \leq \mu'_r + n.$$

In particular,

$$\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, S) = 0 \quad \text{unless } \mu \subset \lambda \text{ and } l(\lambda) \leq l(\mu) + n.$$

Proof. — We use the following elementary observation: if partitions μ and λ interlace, that is,

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \lambda_3 \geq \dots,$$

then for any $r \geq 1$ we have

$$\lambda'_r = \#\{i : \lambda_i \geq r\} \leq \#\{i : \mu_i \geq r\} + 1 = \mu'_r + 1.$$

Now the claim follows from Propositions 5.5 and 5.6: the function $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ vanishes unless there exists a sequence of partitions

$$\mu = \nu^{(0)} \prec \nu^{(1)} \prec \dots \prec \nu^{(n)} = \lambda. \quad \square$$

COROLLARY 5.8. — *We have the following stability property*

$$\mathbb{F}_{\lambda}(\kappa_1, \dots, \kappa_{n-1}, s_n \xi_n \mid \Xi, \mathbf{S}) = \mathbb{F}_{\lambda}(\kappa_1, \dots, \kappa_{n-1} \mid \Xi, \mathbf{S}).$$

Proof. — From Proposition 5.6 we have

$$\mathbb{F}_{\lambda}(\kappa \mid \Xi, \mathbf{S}) = \begin{cases} (-\kappa)^{|\lambda|} \xi_0^{-\lambda_1} \frac{(s_1 \xi_1 / \kappa; q)_{\lambda_1} (q; q)_{\lambda_1}}{(q; q)_{\lambda_1} (s_1^2; q)_{\lambda_1}} & \text{if } \lambda_2 = \lambda_3 = \dots = 0, \\ 0 & \text{otherwise.} \end{cases}$$

After replacing Ξ, \mathbf{S} by $\tau_{\mathbf{S}}^{n-1} \Xi, \tau_{\Xi}^{n-1} \mathbf{S}$ and setting $\kappa = s_1^{(n-1)} \xi_1^{(n-1)} = s_n \xi_n$, we get

$$\mathbb{F}_{\lambda}(s_n \xi_n \mid \tau_{\mathbf{S}}^{n-1} \Xi, \tau_{\Xi}^{n-1} \mathbf{S}) = \mathbb{1}_{\lambda = \emptyset}.$$

The claim now follows from the branching rule of Proposition 5.5 for $m = n - 1$. \square

THEOREM 5.9. — *The functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ are symmetric polynomials in $\kappa_1, \dots, \kappa_n$.*

Proof. — The functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ were observed to be polynomials in Section 5.3. To prove that they are symmetric, we use Proposition 5.3 to write $\mathbb{F}_{\lambda/\mu}^{\mathbf{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ as a matrix coefficient of a composition of the row operators $\mathbb{B}(\kappa \mid \tau_{\mathbf{S}}^i \Xi, \tau_{\Xi}^i \mathbf{S})$ with shifted inhomogeneities. Proposition 4.5 shows that this composition is invariant under simple transpositions exchanging κ_i and κ_{i+1} , which implies the claim. \square

Note that, via Proposition 5.4 and the second relation from Proposition 4.5, the same argument can be applied to the functions $\tilde{\mathbb{F}}_{\lambda/\mu}(u_1, \dots, u_n \mid \Xi, \mathbf{S})$ to directly prove that they are symmetric.

For our next result we need dual functions defined by

$$\tilde{\mathbb{F}}_{\lambda'/\mu'}^*(u_1, \dots, u_n \mid \Xi, \mathbf{S}) := \frac{c_{\mathbf{S}}(\lambda)}{c_{\mathbf{S}}(\mu)} \tilde{\mathbb{F}}_{\lambda'/\mu'}(u_1, \dots, u_n \mid \Xi, \mathbf{S}),$$

$$\begin{aligned} \tilde{\mathbb{F}}_{\lambda'/\mu'}^{\mathbf{S}*}(u_1, \dots, u_n \mid \Xi, \mathbf{S}) &:= \frac{\mathbf{S}^{2\mu} c_{\mathbf{S}}(\mu)}{\mathbf{S}^{2\lambda} c_{\mathbf{S}}(\lambda)} \tilde{\mathbb{F}}_{\lambda'/\mu'}^{\mathbf{S}}(u_1, \dots, u_n \mid \Xi, \mathbf{S}) \\ &= (-1)^{|\mu| - |\lambda|} \frac{\mathbf{S}^{\mu} c_{\mathbf{S}}(\mu)}{\mathbf{S}^{\lambda} c_{\mathbf{S}}(\lambda)} \tilde{\mathbb{F}}_{\lambda'/\mu'}^*(u_1, \dots, u_n \mid \Xi, \mathbf{S}). \end{aligned} \quad (5.7)$$

In view of Proposition 5.4, we have

$$\tilde{\mathbb{F}}_{\lambda'/\mu'}^{\mathbf{S}*}(u_1, \dots, u_n \mid \Xi, \mathbf{S}) = \langle \mu \mid \tilde{\mathbb{B}}^*(u_n \mid \Xi, \mathbf{S}) \dots \tilde{\mathbb{B}}^*(u_2 \mid \Xi, \mathbf{S}) \tilde{\mathbb{B}}^*(u_1 \mid \Xi, \mathbf{S}) \mid \lambda \rangle. \quad (5.8)$$

THEOREM 5.10. — For any partitions μ, ν the following identity holds

$$\begin{aligned} & \sum_{\lambda} \widetilde{F}_{\lambda'/\nu'}^*(u_1, \dots, u_m \mid \Xi, \mathbb{S}) \mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \\ &= \prod_{i=1}^n \prod_{j=1}^m \frac{1 - u_j \kappa_i}{1 - u_j \xi_i s_i} \sum_{\lambda} \mathbb{F}_{\nu/\lambda}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \widetilde{F}_{\mu'/\lambda'}^*(u_1, \dots, u_m \mid \tau_{\Xi}^n \Xi, \tau_{\Xi}^n \mathbb{S}), \end{aligned}$$

with both sums above having finitely many nonzero terms.

Proof. — Note that by (5.2) and (5.7), Theorem 5.10 is equivalent to the following identity:

$$\begin{aligned} & \sum_{\lambda} \widetilde{F}_{\lambda'/\nu'}^{\mathbb{S}*}(u_1, \dots, u_m \mid \Xi, \mathbb{S}) \mathbb{F}_{\lambda/\mu}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \\ &= \prod_{i=1}^n \prod_{j=1}^m \frac{1 - u_j \kappa_i}{1 - u_j \xi_i s_i} \sum_{\lambda} \mathbb{F}_{\nu/\lambda}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \widetilde{F}_{\mu'/\lambda'}^{\mathbb{S}*}(u_1, \dots, u_m \mid \tau_{\Xi}^n \Xi, \tau_{\Xi}^n \mathbb{S}). \end{aligned}$$

So we will actually prove this identity instead of the identity in the theorem. The proof follows a standard outline given in [15, Theorem 7.3]: We rewrite the matrix element

$$\begin{aligned} \mathcal{E}_{\mu}^{\nu}(\kappa_1, \dots, \kappa_n; u_1, \dots, u_m \mid \Xi, \mathbb{S}) &:= \langle \nu \mid \widetilde{B}^*(u_m \mid \Xi, \mathbb{S}) \dots \widetilde{B}^*(u_1 \mid \Xi, \mathbb{S}) \\ &\quad \times \mathbb{B}(\kappa_1 \mid \tau_{\Xi} \Xi, \tau_{\Xi} \mathbb{S}) \mathbb{B}(\kappa_2 \mid \tau_{\Xi}^2 \Xi, \tau_{\Xi}^2 \mathbb{S}) \dots \mathbb{B}(\kappa_n \mid \tau_{\Xi}^n \Xi, \tau_{\Xi}^n \mathbb{S}) \mid \mu \rangle \end{aligned}$$

in two different ways, corresponding to the two sides of the claim.

For the left-hand side we insert $\sum_{\lambda} |\lambda\rangle\langle\lambda| = \text{id}$ between the operators \mathbb{C} and \widetilde{B}^* to get

$$\begin{aligned} \mathcal{E}_{\mu}^{\nu}(\kappa_1, \dots, \kappa_n; u_1, \dots, u_m \mid \Xi, \mathbb{S}) \\ &= \sum_{\lambda} \widetilde{F}_{\lambda'/\nu'}^{\mathbb{S}*}(u_1, \dots, u_m \mid \Xi, \mathbb{S}) \mathbb{F}_{\lambda/\mu}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}), \end{aligned}$$

where we have used Proposition 5.3 and relation (5.8).

On the other hand, set

$$\mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) := \mathbb{B}(\kappa_1 \mid \tau_{\Xi} \Xi, \tau_{\Xi} \mathbb{S}) \mathbb{B}(\kappa_2 \mid \tau_{\Xi}^2 \Xi, \tau_{\Xi}^2 \mathbb{S}) \dots \mathbb{B}(\kappa_n \mid \tau_{\Xi}^n \Xi, \tau_{\Xi}^n \mathbb{S})$$

and iterate the exchange relation from Proposition 4.6 to get

$$\begin{aligned} & \widetilde{B}^*(u \mid \Xi, \mathbb{S}) \mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \\ &= \prod_{i=1}^n \frac{1 - u \kappa_i}{1 - u \xi_i s_i} \mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \widetilde{B}^*(u \mid \tau_{\Xi}^n \Xi, \tau_{\Xi}^n \mathbb{S}). \end{aligned}$$

Hence,

$$\begin{aligned} \mathcal{E}_\mu^\nu(\kappa_1, \dots, \kappa_n; u_1, \dots, u_m \mid \Xi, \mathbf{S}) &= \prod_{i=1}^n \prod_{j=1}^m \frac{1 - u_j \kappa_i}{1 - u_j \xi_i s_i} \\ &\times \langle \nu \mid \mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) \tilde{B}^*(u_m \mid \tau_{\mathbf{S}}^n \Xi, \tau_{\Xi}^n \mathbf{S}) \dots \tilde{B}^*(u_1 \mid \tau_{\mathbf{S}}^n \Xi, \tau_{\Xi}^n \mathbf{S}) \mid \mu \rangle. \end{aligned}$$

Now we can again insert $\sum_\lambda |\lambda\rangle\langle\lambda| = \text{id}$ and obtain

$$\begin{aligned} \mathcal{E}_\mu^\nu(\kappa_1, \dots, \kappa_n; u_1, \dots, u_m \mid \Xi, \mathbf{S}) &= \prod_{i=1}^n \prod_{j=1}^m \frac{1 - u_j \kappa_i}{1 - u_j \xi_i s_i} \\ &\times \sum_\lambda \mathbb{F}_{\nu/\lambda}^{\mathbf{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) \tilde{\mathbb{F}}_{\mu'/\lambda'}^{\mathbf{S}^*}(u_1, \dots, u_m \mid \tau_{\mathbf{S}}^n \Xi, \tau_{\Xi}^n \mathbf{S}). \quad \square \end{aligned}$$

By setting $\mu = \emptyset$ or $(\mu, \nu) = (\emptyset, \emptyset)$ and noting that

$$\mathbb{F}_{\emptyset/\lambda}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) = \tilde{\mathbb{F}}_{\emptyset/\lambda}(u_1, \dots, u_m \mid \Xi, \mathbf{S}) = \mathbb{1}_{\lambda=\emptyset},$$

Theorem 5.10 immediately leads to the following results:

COROLLARY 5.11 (Pieri rule). — *For any partition ν we have*

$$\begin{aligned} \prod_{i=1}^n \prod_{j=1}^m \frac{1 - u_j \kappa_i}{1 - u_j \xi_i s_i} \mathbb{F}_\nu(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) \\ = \sum_\lambda \tilde{\mathbb{F}}_{\lambda'/\nu'}^{\mathbf{S}^*}(u_1, \dots, u_m \mid \Xi, \mathbf{S}) \mathbb{F}_\lambda(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}). \end{aligned}$$

COROLLARY 5.12 (Dual Cauchy identity). — *We have*

$$\sum_\lambda \tilde{\mathbb{F}}_{\lambda'}^{\mathbf{S}^*}(u_1, \dots, u_m \mid \Xi, \mathbf{S}) \mathbb{F}_\lambda(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) = \prod_{i=1}^n \prod_{j=1}^m \frac{1 - u_j \kappa_i}{1 - u_j \xi_i s_i}.$$

5.5. Integral representation

We can use the dual Cauchy identity of Corollary 5.12 and previously known results about inhomogeneous (stable) spin Hall–Littlewood functions to obtain a multivariate integral representation for the inhomogeneous spin q -Whittaker functions.

To formulate the result, set q to be a complex number such that $0 < |q| < 1$ and assume that the variables $\kappa_1, \dots, \kappa_n$ and the parameters $\Xi = (\xi_0, \xi_1, \dots)$, $\mathbf{S} = (s_0, s_1, \dots)$ are complex numbers such that there exists a positively-oriented simple contour \mathcal{C} on the complex plane satisfying the following conditions:




Figure 5.3. A possible of configuration of $\{\xi_i s_i\}_{i=1}^{\infty}$, $\{\xi_i^{-1} s_i\}_{i=1}^{\infty}$ and \mathcal{C} for Theorem 5.13.

- all points $\{0\}$ and $\{\xi_i^{-1} s_i\}_{i=1}^{\infty}$ are inside the contour \mathcal{C} ;
- all points $\{\xi_i^{-1} s_i^{-1}\}_{i=1}^{\infty}$ are outside of the contour \mathcal{C} ;
- the image $q\mathcal{C}$ of the contour \mathcal{C} under the multiplication by q is inside \mathcal{C} .

An example of such a configuration is sketched in Figure 5.3.

THEOREM 5.13. — *Under the above assumptions, we have*

$$\begin{aligned} \mathbb{F}_{\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S}) &= \oint_{\mathcal{C}} \frac{dz_1}{2\pi i z_1} \cdots \oint_{\mathcal{C}} \frac{dz_k}{2\pi i z_k} \prod_{\alpha < \beta} \frac{z_{\alpha} - z_{\beta}}{z_{\alpha} - qz_{\beta}} \\ &\times \prod_{\alpha=1}^k \left(\frac{\xi_0^{-1}}{z_{\alpha} - \xi_{\mu'_{\alpha}}^{-1} s_{\mu'_{\alpha}}} \prod_{j=1}^{\mu'_{\alpha}-1} \frac{1 - s_j \xi_j z_{\alpha}}{z_{\alpha} \xi_j - s_j} \prod_{i=1}^n \frac{1 - z_{\alpha} \kappa_i}{1 - z_{\alpha} \xi_i s_i} \right), \end{aligned}$$

where $k = \mu_1$.

Proof. — We will use the orthogonality of \tilde{F}_{λ} , cf. [10, Theorem 7.4]: for any pair of partitions λ, μ such that $l(\mu) \leq L$ we have

$$\begin{aligned} \frac{(q; q)_{L-l(\mu)} c_{\mathbf{S}}(\lambda)}{(1-q)^L} \oint_{\mathcal{C}} \frac{dz_1}{2\pi i} \cdots \oint_{\mathcal{C}} \frac{dz_L}{2\pi i} \prod_{\alpha < \beta} \frac{z_{\alpha} - z_{\beta}}{z_{\alpha} - qz_{\beta}} \tilde{F}_{\lambda}(z_1, \dots, z_L \mid \Xi, \mathbf{S}) \\ \times \prod_{\alpha=1}^L z_{\alpha}^{-1} \tilde{\varphi}_{\mu_{\alpha}}(z_{\alpha}^{-1} \mid \bar{\Xi}, \mathbf{S}) = \mathbb{1}_{\lambda=\mu}, \quad (5.9) \end{aligned}$$

where

$$\bar{\Xi} = (\xi_0^{-1}, \xi_1^{-1}, \xi_2^{-1}, \dots)$$

and the functions $\tilde{\varphi}_k(u \mid \Xi, \mathbf{S})$ were defined at the end of Section 5.2.

Fixing a partition μ , conjugating partitions in (5.9), setting $L = k = \mu_1 = l(\mu')$ and multiplying by $\mathbb{F}_\lambda(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S})$, we get

$$\begin{aligned} & \frac{1}{(1-q)^k} \oint_{\mathcal{C}} \frac{dz_1}{2\pi\mathbf{i}} \cdots \oint_{\mathcal{C}} \frac{dz_k}{2\pi\mathbf{i}} \prod_{\alpha < \beta} \frac{z_\alpha - z_\beta}{z_\alpha - qz_\beta} \\ & \times \widetilde{\mathbb{F}}_{\lambda'}^*(z_1, \dots, z_k \mid \Xi, \mathbb{S}) \mathbb{F}_\lambda(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \prod_{\alpha=1}^k z_\alpha^{-1} \widetilde{\varphi}_{\mu'_\alpha}(z_\alpha^{-1} \mid \bar{\Xi}, \mathbb{S}) \\ & = \mathbb{1}_{\lambda=\mu} \cdot \mathbb{F}_\lambda(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}). \end{aligned}$$

Summing over λ and using Corollary 5.12, we arrive at the integral representation for $\mathbb{F}_\mu(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S})$. \square

6. A Cauchy identity

In this section we use fusion of \widetilde{B}^* -operators to get a Cauchy-like identity between inhomogeneous spin q -Whittaker functions $\mathbb{F}_{\lambda/\mu}$ similar to [10, Corollary 4.10] and [15, Theorem 7.1]. However, during the proof we will lose the degrees of freedom corresponding to the parameters Ξ , proving an identity involving only functions with $\Xi = \mathbb{S}$ and $\bar{\Xi} = \bar{\mathbb{S}}$. Recall that for a sequence $\mathbb{S} = (s_0, s_1, \dots)$ we set

$$\bar{\mathbb{S}} := (s_0^{-1}, s_1^{-1}, s_2^{-1}, \dots).$$

6.1. $\Xi = \mathbb{S}$ and $\bar{\Xi} = \bar{\mathbb{S}}$ specializations

We start with the description of a simplification of functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S})$ when $\Xi = \mathbb{S}$ and $\bar{\Xi} = \bar{\mathbb{S}}$. In these two cases, the parameters of the vertex weights in the partition function $ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S})$ degenerate to $(\frac{s_{i+j}}{\sqrt{\kappa_i}}, s_{i+j})$ for $\Xi = \mathbb{S}$, and to $(\kappa_i^{-\frac{1}{2}}, s_j)$ for $\bar{\Xi} = \bar{\mathbb{S}}$, where i and j denote the row and the column of the corresponding vertex. These parameters are depicted in Figure 6.1.

Note that the action of the mixed shift operators τ_{Ξ} and $\tau_{\bar{\mathbb{S}}}$ also simplifies:

$$\tau_{\mathbb{S}}\mathbb{S} = \tau\mathbb{S}, \quad \tau_{\bar{\mathbb{S}}}\bar{\mathbb{S}} = \bar{\mathbb{S}}, \quad \tau_{\bar{\mathbb{S}}}\mathbb{S} = \mathbb{S},$$

where τ denotes the ordinary shift operator

$$\tau^i \mathbb{S} := (s_i, s_{i+1}, s_{i+2}, \dots), \quad i \geq 0.$$




Figure 6.1. Partition functions $ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \mathbf{S}, \mathbf{S})$ and $ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \bar{\mathbf{S}}, \mathbf{S})$

Then the functions $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbf{S})$ simplify to

$$\begin{aligned}
 & \mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \mathbf{S}, \mathbf{S}) \\
 &= (-1)^{|\mu| - |\lambda|} \frac{(\tau^n \mathbf{S})^\mu}{S^\lambda} \frac{c_{\tau^n \mathbf{S}}(\mu)}{c_{\mathbf{S}}(\lambda)} \mathbb{F}_{\lambda/\mu}^{\mathbf{S}}(\kappa_1, \dots, \kappa_n \mid \mathbf{S}, \mathbf{S}) \\
 &= (-1)^{|\mu| - |\lambda|} \frac{(\tau^n \mathbf{S})^\mu}{S^\lambda} \frac{c_{\tau^n \mathbf{S}}(\mu)}{c_{\mathbf{S}}(\lambda)} \\
 & \quad \times \sum_{a_1, \dots, a_n} \left(\prod_{i=1}^n \kappa_i^{a_i} \frac{(s_i^2 / \kappa_i; q)_{a_i}}{(q; q)_{a_i}} \right) ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \mathbf{S}, \mathbf{S}).
 \end{aligned}$$

$$\begin{aligned}
 & \mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \bar{\mathbf{S}}, \mathbf{S}) \\
 &= (-1)^{|\mu| - |\lambda|} \frac{S^\mu}{S^\lambda} \frac{c_{\mathbf{S}}(\mu)}{c_{\mathbf{S}}(\lambda)} \mathbb{F}_{\lambda/\mu}^{\mathbf{S}}(\kappa_1, \dots, \kappa_n \mid \bar{\mathbf{S}}, \mathbf{S}) \\
 &= (-1)^{|\mu| - |\lambda|} \frac{S^\mu}{S^\lambda} \frac{c_{\mathbf{S}}(\mu)}{c_{\mathbf{S}}(\lambda)} \\
 & \quad \times \sum_{a_1, \dots, a_n} \left(\prod_{i=1}^n (\kappa_i s_0^2)^{a_i} \frac{(\kappa_i^{-1}; q)_{a_i}}{(q; q)_{a_i}} \right) ZW_{\lambda/\mu}^{(a_1, \dots, a_n)}(\kappa_1, \dots, \kappa_n \mid \bar{\mathbf{S}}, \mathbf{S}).
 \end{aligned}$$

6.2. Cauchy identity

For any pair of partitions λ, μ define *dual inhomogeneous spin q -Whittaker polynomials* by (cf. (5.2))

$$\mathbb{F}_{\lambda/\mu}^*(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) := \frac{c_{\mathbb{S}}(\lambda)}{c_{\tau_{\Xi}^n \mathbb{S}}(\mu)} \mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}).$$

Additionally, define

$$\begin{aligned} \mathbb{F}_{\lambda/\mu}^{\mathbb{S}*}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) &:= \frac{(\tau_{\Xi}^n \mathbb{S})^{2\mu} c_{\tau_{\Xi}^n \mathbb{S}}(\mu)}{S^{2\lambda} c_{\mathbb{S}}(\lambda)} \mathbb{F}_{\lambda/\mu}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \\ &= (-1)^{|\mu| - |\lambda|} \frac{(\tau_{\Xi}^n \mathbb{S})^{\mu} c_{\tau_{\Xi}^n \mathbb{S}}(\mu)}{S^{\lambda} c_{\mathbb{S}}(\lambda)} \mathbb{F}_{\lambda/\mu}^*(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}). \end{aligned}$$

In view of Proposition 5.3, we have

$$\begin{aligned} \mathbb{F}_{\lambda/\mu}^{\mathbb{S}*}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) &= \langle \mu \mid \mathbb{B}^*(\kappa_n \mid \tau_{\mathbb{S}}^{n-1} \Xi, \tau_{\Xi}^{n-1} \mathbb{S}) \dots \mathbb{B}^*(\kappa_2 \mid \tau_{\mathbb{S}} \Xi, \tau_{\Xi} \mathbb{S}) \mathbb{B}^*(\kappa_1 \mid \Xi, \mathbb{S}) \mid \lambda \rangle. \end{aligned}$$

THEOREM 6.1. — *The following identity holds in the space of formal power series in $\kappa_1, \dots, \kappa_n, \chi_1, \dots, \chi_m, q$ with rational in \mathbb{S} coefficients:*

$$\begin{aligned} &\sum_{\lambda} \mathbb{F}_{\lambda/\nu}^*(\chi_1, \dots, \chi_m \mid \bar{\mathbb{S}}, \mathbb{S}) \mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \mathbb{S}, \mathbb{S}) \\ &= \prod_{i=1}^n \prod_{j=1}^m \frac{(\kappa_i; q)_{\infty} (s_i^2 \chi_j; q)_{\infty}}{(s_i^2; q)_{\infty} (\kappa_i \chi_j; q)_{\infty}} \\ &\quad \times \sum_{\lambda} \mathbb{F}_{\nu/\lambda}(\kappa_1, \dots, \kappa_n \mid \mathbb{S}, \mathbb{S}) \mathbb{F}_{\mu/\lambda}^*(\chi_1, \dots, \chi_m \mid \tau^n \bar{\mathbb{S}}, \tau^n \mathbb{S}). \quad (6.1) \end{aligned}$$

Proof. — Similarly to the proof of Theorem 5.10, we will actually prove an equivalent identity

$$\begin{aligned} &\sum_{\lambda} \mathbb{F}_{\lambda/\nu}^{\mathbb{S}*}(\chi_1, \dots, \chi_m \mid \bar{\mathbb{S}}, \mathbb{S}) \mathbb{F}_{\lambda/\mu}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \mathbb{S}, \mathbb{S}) \\ &= \prod_{i=1}^n \prod_{j=1}^m \frac{(\kappa_i; q)_{\infty} (s_i^2 \chi_j; q)_{\infty}}{(s_i^2; q)_{\infty} (\kappa_i \chi_j; q)_{\infty}} \\ &\quad \times \sum_{\lambda} \mathbb{F}_{\nu/\lambda}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \mathbb{S}, \mathbb{S}) \mathbb{F}_{\mu/\lambda}^{\mathbb{S}*}(\chi_1, \dots, \chi_m \mid \tau^n \bar{\mathbb{S}}, \tau^n \mathbb{S}), \quad (6.2) \end{aligned}$$

which is obtained by rescaling \mathbb{F}, \mathbb{F}^* to $\mathbb{F}^{\mathbb{S}}, \mathbb{F}^{\mathbb{S}*}$.

Following the proof of Theorem 5.10, let

$$\mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid \mathbb{S}, \mathbb{S}) = \mathbb{B}(\kappa_1 \mid \tau \mathbb{S}, \tau \mathbb{S}) \dots \mathbb{B}(\kappa_n \mid \tau^n \mathbb{S}, \tau^n \mathbb{S}).$$

Then the exchange relation of Proposition 4.6 together with the relation $\tau_S S = \tau S$ imply

$$\begin{aligned} & \tilde{B}^*(1 \mid S, S) \dots \tilde{B}^*(q^{J-1} \mid S, S) \mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid S, S) \\ &= \prod_{i=1}^n \frac{(\kappa_i; q)_J}{(s_i^2; q)_J} \mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid S, S) \tilde{B}^*(1 \mid \tau^n S, \tau^n S) \dots \tilde{B}^*(q^{J-1} \mid \tau^n S, \tau^n S). \end{aligned}$$

Now we can use Proposition 4.3 to get

$$\begin{aligned} & \mathbb{B}^*(q^J \mid \bar{S}, S) \mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid S, S) \\ &= \prod_{i=1}^n \frac{(\kappa_i; q)_\infty (s_i^2 q^J; q)_\infty}{(s_i^2; q)_\infty (\kappa_i q^J; q)_\infty} \mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid S, S) \mathbb{B}^*(q^J \mid \tau^n \bar{S}, \tau^n S). \end{aligned}$$

Iterating this equation m -times for $J = J_1, \dots, J_m$, we obtain

$$\begin{aligned} & \mathbb{B}^*(q^{J_m} \mid \bar{S}, S) \dots \mathbb{B}^*(q^{J_1} \mid \bar{S}, S) \mathbb{C}^{(n)}(\kappa_1, \dots, \kappa_n \mid S, S) \\ &= \prod_{i=1}^n \prod_{j=1}^m \frac{(\kappa_i; q)_\infty (s_i^2 q^{J_j}; q)_\infty}{(s_i^2; q)_\infty (\kappa_i q^{J_j}; q)_\infty} \mathbb{B}^{(n)}(\kappa_1, \dots, \kappa_n \mid S, S) \\ & \quad \times \mathbb{B}^*(q^{J_m} \mid \tau^n \bar{S}, \tau^n S) \dots \mathbb{B}^*(q^{J_1} \mid \tau^n \bar{S}, \tau^n S), \end{aligned}$$

which, after taking the matrix element $\langle \nu \mid \cdot \mid \mu \rangle$, inserting a summation $\sum_\lambda |\lambda\rangle \langle \lambda| = \text{id}$, and using both identities from Proposition 5.3, gives the claim of the theorem for $\chi_r = q^{J_r}$.

Up until now we have operated with finite sums of rational functions: one can see⁽⁹⁾ that for $(\chi_1, \dots, \chi_m) = (q^{J_1}, \dots, q^{J_m})$ with $\{J_r\}_{r=1}^m \in \mathbb{Z}_{\geq 1}$ both sums in (6.1) have finitely many nonzero terms. But for general parameters χ the sum in the left-hand side is infinite, so we need to be careful with the general statement.

We say that a formal power series $f \in K[[x_1, \dots, x_r]]$ over a field K is of *order at least N* if coefficients of all monomials $x_1^{a_1} \dots x_r^{a_r}$ such that $\sum_{i=1}^r a_i < N$ are equal to 0. We claim that $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, S)$ is a formal power series in $K[[\kappa_1, \dots, \kappa_n, q]]$ of order at least $\lambda_1 - \mu_1 - n$, where $K = \mathbb{Q}(\Xi, S)$ is the field of rational functions in variables $\Xi = (\xi_0, \xi_1, \dots)$ and $S = (s_0, s_1, \dots)$. To see that, note that the matrix coefficients of \mathbb{T}_a are formal power series in κ, q , that is

$$\langle \lambda \mid \mathbb{T}_a(\kappa \mid \Xi, S) \mid \mu \rangle \in K[[\kappa, q]],$$

⁽⁹⁾ for example, using Proposition 4.3 and the discussion from Section 4.3

while the coefficient of \mathbb{T}_a in the definition (4.7) of the operator $\mathbb{B}(\kappa \mid \Xi, \mathbb{S})$ has order at least $a - 1$ with respect to κ, q :

$$\kappa^a s_0^a \xi_0^{-a} \frac{(\kappa^{-1} s_0 \xi_0; q)_a}{(q; q)_a} = \frac{\prod_{i=1}^a (\kappa \xi_0^{-1} - s_0^2 q^{i-1})}{(q; q)_a}.$$

Hence, the matrix element

$$\langle \lambda \mid \mathbb{C}(\kappa \mid \Xi, \mathbb{S}) \mid \mu \rangle \in K[[\kappa, q]]$$

has order at least $\lambda_1 - \mu_1 - 1$. Therefore, by Proposition 5.3, the function $\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S})$ is a formal power series in $\kappa_1, \dots, \kappa_n, q$ of order at least $\lambda_1 - \mu_1 - n$.

Now we can finish the proof. The right-hand side of (6.1) is clearly a formal power series in χ, κ, q . For the sum on the left-hand side, the term corresponding to λ has order at least $2\lambda_1 - \mu_1 - \nu_1 - n - m$. Moreover, the summand corresponding to λ vanishes if $l(\lambda)$ is greater than either $l(\mu) + n$ or $l(\nu) + m$. Overall this means that all but finitely many summands in the left-hand side have order at least N for any $N \geq 1$. Hence both sides of (6.1) are elements of $K[[\kappa_1, \dots, \kappa_n, \chi_1, \dots, \chi_m, q]]$, which are equal after substitution $(\chi_1, \dots, \chi_m) = (q^{J_1}, \dots, q^{J_m})$ for any integer vector $(J_r)_{r=1}^m \in \mathbb{Z}_{\geq 1}^m$ ⁽¹⁰⁾. An iterative application of the following elementary statement finishes the proof. \square

LEMMA 6.2 ([34, Lemma 3.2]). — *Let $F(z, q)$ and $G(z, q)$ be formal power series satisfying $F(q^J, q) = G(q^J, q)$ for infinitely many integers $J \geq 1$. Then $F(z, q) = G(z, q)$.*

Remark 6.3. — The convergence from Theorem 6.1 also holds numerically if

$$|\kappa_1|, \dots, |\kappa_n| \leq \kappa, \quad |\chi_1|, \dots, |\chi_m| \leq \chi, \quad 0 < |q|, \kappa\chi < 1,$$

and $s_i^2 \neq q^{-m}$ for any $i, m \in \mathbb{Z}_{\geq 0}$. For the right-hand side of (6.1) the convergence is immediate, while for the left-hand side this can be verified as follows: Using Proposition 5.6, the one variable functions $\mathbb{F}_{\lambda/\mu}(\kappa_i \mid \Xi, \mathbb{S})$ can be bounded by $C|\kappa_i|^{|\lambda|} \frac{(\tau_s \Xi)^\mu}{(\Xi)^\lambda}$, where C is a positive constant not depending on λ, μ (to reach this bound it is enough to bound a finite number of possibly appearing q -Pochhammer symbols $(x, q)_a$ by a uniform constant and rearrange the remaining terms). Then, by the branching rule of Proposition 5.5, we obtain an upper bound

$$|\mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S})| \leq C|\lambda|^d \kappa^{|\lambda|} / (\Xi)^\lambda,$$

⁽¹⁰⁾ Note that such a substitution is well-defined and it behaves well with respect to the convergence of formal power series, as well as the notion of order defined earlier. Indeed, the substitution cannot reduce the order of a formal power series, hence the substitution preserves convergence.

for some $C, d > 0$ not depending on λ . This bound is sufficient for proving the convergence of the left-hand side of (6.1).

Remark 6.4. — One can note a similarity between Theorem 6.1 and the $\Xi = \mathbf{S}$ case of the Cauchy identity for inhomogeneous spin Hall–Littlewood functions [10, Corollary 4.10]. Thus, one might hope that it is possible to remove the restriction $\Xi = \mathbf{S}$, although we don’t know how to do that.

By setting μ and ν equal to \emptyset one immediately obtains a Cauchy type summation identity:

COROLLARY 6.5. — *The following identity holds in the space of formal power series in $\kappa_1, \dots, \kappa_n, \chi_1, \dots, \chi_m, q$ with rational in \mathbf{S} coefficients:*

$$\sum_{\lambda} \mathbb{F}_{\lambda}^*(\chi_1, \dots, \chi_m \mid \bar{\mathbf{S}}, \mathbf{S}) \mathbb{F}_{\lambda}(\kappa_1, \dots, \kappa_n \mid \mathbf{S}, \mathbf{S}) = \prod_{i=1}^n \prod_{j=1}^m \frac{(\kappa_i; q)_{\infty} (s_i^2 \chi_j; q)_{\infty}}{(s_i^2; q)_{\infty} (\kappa_i \chi_j; q)_{\infty}}.$$

The same identity holds numerically for $0 < |q|, |\kappa_1|, \dots, |\kappa_n|, |\chi_1|, \dots, |\chi_m| < 1$.

Remark 6.6. — As noted in [15, Section 7.2], in the simplest case when $n = m = 1$ Corollary 6.5 reads

$$\sum_{a \geq 0} (-\kappa/s_0)^a \frac{(\kappa^{-1} s_1^2; q)_a}{(s_1^2; q)_a} \cdot (-\chi s_0)^a \frac{(\chi^{-1}; q)_a}{(q; q)_a} = \frac{(\kappa; q)_{\infty} (s_1^2 \chi; q)_{\infty}}{(s_1^2; q)_{\infty} (\kappa \chi; q)_{\infty}},$$

which is equivalent to the q -Gauss summation identity. Actually, we have

$$\begin{aligned} \sum_{a \geq 0} (-\kappa/\xi_0)^a \frac{(\kappa^{-1} s_1 \xi_1; q)_a}{(s_1^2; q)_a} \cdot (-\chi \xi_0)^a \frac{(\chi^{-1} s_1 \xi_1^{-1}; q)_a}{(q; q)_a} \\ = \frac{(\kappa s_1 / \xi_1; q)_{\infty} (\chi s_1 \xi_1; q)_{\infty}}{(s_1^2; q)_{\infty} (\kappa \chi; q)_{\infty}}, \end{aligned}$$

which, along the lines of Remark 6.4, may correspond to a hypothetical more general version of Theorem 6.1 with arbitrary Ξ .

Appendix

Below we give the expressions and identities which are frequently used throughout the paper as intermediate steps for the main results. They are listed in the order they appear in the text.

Vertex weights $w_{u;s}^s, W_{t,s}^s, w_{u;s}^{s*}$ and $W_{t,s}^{s*}$, Section 2:

$$\begin{array}{cccc}
 \begin{array}{c} g \\ \uparrow \\ 0 \text{---} w_{u;s}^s \text{---} 0 \\ \downarrow \\ g \end{array} &
 \begin{array}{c} g-1 \\ \uparrow \\ 0 \text{---} w_{u;s}^s \text{---} 1 \\ \downarrow \\ g \end{array} &
 \begin{array}{c} g+1 \\ \uparrow \\ 1 \text{---} w_{u;s}^s \text{---} 0 \\ \downarrow \\ g \end{array} &
 \begin{array}{c} g \\ \uparrow \\ 1 \text{---} w_{u;s}^s \text{---} 1 \\ \downarrow \\ g \end{array} \\
 \frac{1-suq^g}{1-su} &
 \frac{(q^g-1)su}{1-su} &
 \frac{1-s^2q^g}{1-su} &
 \frac{s^2q^g-su}{1-su}
 \end{array} \tag{2.2}$$

$$\begin{array}{c} k \\ \uparrow \\ j \text{---} W_{t,s}^s \text{---} l \\ \downarrow \\ i \end{array} = \mathbb{1}_{i+j=k+l} \mathbb{1}_{i \geq l} s^{2l} t^{-2l} \frac{(s^2/t^2; q)_{i-l} (t^2; q)_l}{(s^2; q)_i} \frac{(q; q)_i}{(q; q)_{i-l} (q; q)_l}, \tag{2.9}$$

$$\begin{array}{cccc}
 \begin{array}{c} g \\ \uparrow \\ 0 \text{---} w_{u;s}^{s*} \text{---} 0 \\ \downarrow \\ g \end{array} &
 \begin{array}{c} g+1 \\ \uparrow \\ 0 \text{---} w_{u;s}^{s*} \text{---} 1 \\ \downarrow \\ g \end{array} &
 \begin{array}{c} g-1 \\ \uparrow \\ 1 \text{---} w_{u;s}^{s*} \text{---} 0 \\ \downarrow \\ g \end{array} &
 \begin{array}{c} g \\ \uparrow \\ 1 \text{---} w_{u;s}^{s*} \text{---} 1 \\ \downarrow \\ g \end{array} \\
 \frac{1-suq^g}{1-su} &
 \frac{-s^{-1}u(1-s^2q^g)}{1-su} &
 \frac{1-q^g}{1-su} &
 \frac{q^g-s^{-1}u}{1-su}
 \end{array} \tag{2.13}$$

$$\begin{array}{c} k \\ \uparrow \\ j \text{---} W_{t,s}^{s*} \text{---} l \\ \downarrow \\ i \end{array} = \mathbb{1}_{i+l=j+k} \mathbb{1}_{i \geq j} t^{2i-2j} \frac{(s^2/t^2; q)_{i-j} (t^2; q)_j}{(s^2; q)_i} \frac{(q; q)_i}{(q; q)_{i-j} (q; q)_j}. \tag{2.18}$$

Deformed Yang–Baxter equations, see Section 2 for the case $\eta = 1$ and Section 3 for the general case:

$$(3.1)$$

$$(3.2)$$

Notation for the mixed shift operator and the inverse sequence, here $\mathcal{A} = (a_0, a_1, a_2, \dots)$:

$$\tau_{\mathcal{B}}^k \mathcal{A} = (a_0^{(k)}, a_1^{(k)}, \dots), \quad a_i^{(k)} = \sqrt{a_{i+k} b_{i+k} a_i / b_i}, \quad \bar{\mathcal{A}} = (a_0^{-1}, a_1^{-1}, a_2^{-1}, \dots).$$

Row operators $\mathbb{T}_a, \mathbb{T}_a^*$ and T_a^* , Section 4: Row operators $\mathbb{T}_a, \mathbb{T}_a^*$ and T_a^* , Section 4:

$$\langle \lambda | \mathbb{T}_a(\kappa | \Xi, S) | \mu \rangle = a \dots \rightarrow 0, \quad (4.2)$$

The functions $\mathbb{F}_{\lambda/\mu}$ and $\tilde{\mathbb{F}}_{\lambda/\mu}$ are defined using vertex models in Sections 5.1 and 5.2. The renormalizations $\mathbb{F}_{\lambda/\mu}^{\mathbb{S}}$ and $\tilde{\mathbb{F}}_{\lambda/\mu}^{\mathbb{S}}$ are given by:

$$\begin{aligned} \mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) &:= (-1)^{|\mu| - |\lambda|} \frac{(\tau_{\Xi}^n \mathbb{S})^\mu}{\mathbb{S}^\lambda} \frac{c_{\tau_{\Xi}^n \mathbb{S}}(\mu)}{c_{\mathbb{S}}(\lambda)} \mathbb{F}_{\lambda/\mu}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}), \\ \tilde{\mathbb{F}}_{\lambda'/\mu'}(u_1, \dots, u_n \mid \Xi, \mathbb{S}) &:= (-1)^{|\mu| - |\lambda|} \frac{\mathbb{S}^\mu}{\mathbb{S}^\lambda} \frac{c_{\mathbb{S}}(\mu)}{c_{\mathbb{S}}(\lambda)} \tilde{\mathbb{F}}_{\lambda'/\mu'}^{\mathbb{S}}(u_1, \dots, u_n \mid \Xi, \mathbb{S}), \\ c_{\mathbb{S}}(\lambda) &:= \prod_{i \geq 1} \frac{(s_i^2; q)_{\lambda_i - \lambda_{i+1}}}{(q; q)_{\lambda_i - \lambda_{i+1}}}, \quad \mathbb{S}^\lambda := \prod_{i \geq 1} s_{i-1}^{\lambda_i}. \end{aligned}$$

The expressions for $\mathbb{F}_{\lambda/\mu}^{\mathbb{S}}$ and $\tilde{\mathbb{F}}_{\lambda/\mu}^{\mathbb{S}}$ in terms of \mathbb{B} and \tilde{B}^* , Propositions 5.3, 5.4:

$$\begin{aligned} &\mathbb{F}_{\lambda/\mu}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \\ &= \langle \lambda \mid \mathbb{B}(\kappa_1 \mid \tau_{\mathbb{S}} \Xi, \tau_{\Xi} \mathbb{S}) \mathbb{B}(\kappa_2 \mid \tau_{\mathbb{S}}^2 \Xi, \tau_{\Xi}^2 \mathbb{S}) \dots \mathbb{B}(\kappa_n \mid \tau_{\mathbb{S}}^n \Xi, \tau_{\Xi}^n \mathbb{S}) \mid \mu \rangle \\ &= \frac{\mathbb{S}^{2\lambda}}{(\tau_{\Xi} \mathbb{S})^{2\mu}} \frac{c_{\mathbb{S}}(\lambda)}{c_{\tau_{\Xi} \mathbb{S}}(\mu)} \\ &\quad \times \langle \mu \mid \mathbb{B}^*(\kappa_n \mid \tau_{\mathbb{S}}^{n-1} \Xi, \tau_{\Xi}^{n-1} \mathbb{S}) \dots \mathbb{B}^*(\kappa_2 \mid \tau_{\mathbb{S}} \Xi, \tau_{\Xi} \mathbb{S}) \mathbb{B}^*(\kappa_1 \mid \Xi, \mathbb{S}) \mid \lambda \rangle, \quad (5.4) \end{aligned}$$

$$\begin{aligned} &\tilde{\mathbb{F}}_{\lambda'/\mu'}^{\mathbb{S}}(u_1, \dots, u_n \mid \Xi, \mathbb{S}) \\ &= \frac{\mathbb{S}^{2\lambda}}{\mathbb{S}^{2\mu}} \frac{c_{\mathbb{S}}(\lambda)}{c_{\mathbb{S}}(\mu)} \langle \mu \mid \tilde{B}^*(u_n \mid \Xi, \mathbb{S}) \dots \tilde{B}^*(u_2 \mid \Xi, \mathbb{S}) \tilde{B}^*(u_1 \mid \Xi, \mathbb{S}) \mid \lambda \rangle. \quad (5.5) \end{aligned}$$

The dual functions $\mathbb{F}_{\lambda/\mu}^*$, Section 6:

$$\begin{aligned} \mathbb{F}_{\lambda/\mu}^*(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) &:= \frac{c_{\mathbb{S}}(\lambda)}{c_{\tau_{\Xi}^n \mathbb{S}}(\mu)} \mathbb{F}_{\lambda/\mu}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}). \\ \mathbb{F}_{\lambda/\mu}^{\mathbb{S}*}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) &:= \frac{(\tau_{\Xi}^n \mathbb{S})^{2\mu}}{\mathbb{S}^{2\lambda}} \frac{c_{\tau_{\Xi}^n \mathbb{S}}(\mu)}{c_{\mathbb{S}}(\lambda)} \mathbb{F}_{\lambda/\mu}^{\mathbb{S}}(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}) \\ &= (-1)^{|\mu| - |\lambda|} \frac{(\tau_{\Xi}^n \mathbb{S})^\mu}{\mathbb{S}^\lambda} \frac{c_{\tau_{\Xi}^n \mathbb{S}}(\mu)}{c_{\mathbb{S}}(\lambda)} \mathbb{F}_{\lambda/\mu}^*(\kappa_1, \dots, \kappa_n \mid \Xi, \mathbb{S}). \end{aligned}$$

Bibliography

- [1] A. AGGARWAL, “Current fluctuations of the stationary ASEP and six-vertex model”, *Duke Math. J.* **167** (2018), no. 2, p. 269-384.
- [2] A. AGGARWAL & A. BORODIN, “Phase transitions in the ASEP and stochastic six-vertex model”, *Ann. Probab.* **47** (2019), no. 2, p. 613-689.
- [3] G. BARRAQUAND & I. CORWIN, “Random walk in beta-distributed random environment”, *Probab. Theory Relat. Fields* **167** (2017), no. 3, p. 1057-1116.

- [4] A. BORODIN, “On a family of symmetric rational functions”, *Adv. Math.* **306** (2017), p. 973-1018.
- [5] ———, “Symmetric elliptic functions, IRF models, and dynamic exclusion processes”, *J. Eur. Math. Soc.* **22** (2020), no. 5, p. 1353-1421.
- [6] A. BORODIN, I. CORWIN, L. PETROV & T. SASAMOTO, “Spectral theory for interacting particle systems solvable by coordinate Bethe ansatz”, *Commun. Math. Phys.* **339** (2015), no. 3, p. 1167-1245.
- [7] ———, “Spectral theory for the q -Boson particle system”, *Compos. Math.* **151** (2015), no. 1, p. 1-67.
- [8] A. BORODIN, V. GORIN & M. WHEELER, “Shift-invariance for vertex models and polymers”, <https://arxiv.org/abs/1912.02957>, 2019.
- [9] A. BORODIN & L. PETROV, “Integrable probability: Stochastic vertex models and symmetric functions”, in *Stochastic Processes and Random Matrices: Lecture Notes of the Les Houches Summer School (Les Houches, 2015)*, vol. 104, Oxford University Press, 2017, p. 26-131.
- [10] ———, “Higher spin six vertex model and symmetric rational functions”, *Sel. Math., New Ser.* **24** (2018), no. 2, p. 751-874.
- [11] ———, “Inhomogeneous exponential jump model”, *Probab. Theory Relat. Fields* **172** (2018), no. 1-2, p. 323-385.
- [12] A. BORODIN & M. WHEELER, “Coloured stochastic vertex models and their spectral theory”, <https://arxiv.org/abs/1808.01866>, 2018.
- [13] ———, “Nonsymmetric Macdonald polynomials via integrable vertex models”, <https://arxiv.org/abs/1904.06804>, to appear in *Trans. Am. Math. Soc.*, 2019.
- [14] ———, “Observables of coloured stochastic vertex models and their polymer limits”, *Probab. Math. Phys.* **1** (2020), no. 1, p. 205-265.
- [15] ———, “Spin q -Whittaker polynomials”, *Adv. Math.* **376** (2020), article no. 107449 (51 pages).
- [16] G. BOSNJAK & V. MANGAZEEV, “Construction of R -matrices for symmetric tensor representations related to $U_q(\widehat{sl}_n)$ ”, *J. Phys. A. Math. Theor.* **49** (2016), no. 49, article no. 495204 (19 pages).
- [17] B. BRUBAKER, D. BUMP & S. FRIEDBERG, “Schur Polynomials and The Yang-Baxter Equation”, *Commun. Math. Phys.* **308** (2011), no. 2, p. 281-301.
- [18] V. BUCIUMAS & T. SCRIMSHAW, “Double Grothendieck polynomials and colored lattice models”, <https://arxiv.org/abs/2007.04533>, 2020.
- [19] A. BUFETOV & S. KOROTKIKH, “Observables of stochastic colored vertex models and local relation”, <https://arxiv.org/abs/2011.11426>, 2020.
- [20] A. BUFETOV, M. MUCCICONI & L. PETROV, “Yang–Baxter random fields and stochastic vertex models”, <https://arxiv.org/abs/1905.06815>, 2019.
- [21] L. CANTINI, J. DE GIER & M. WHEELER, “Matrix product formula for Macdonald polynomials”, *J. Phys. A. Math. Theor.* **48** (2015), no. 38, article no. 384001 (25 pages).
- [22] S. FOMIN & A. N. KIRILLOV, “The Yang-Baxter equation, symmetric functions, and Schubert polynomials”, *Discrete Math.* **153** (1996), no. 1-3, p. 123-143.
- [23] A. GARBALI, J. DE GIER & M. WHEELER, “A new generalisation of Macdonald polynomials”, *Commun. Math. Phys.* **352** (2017), no. 2, p. 773-804.
- [24] A. GARBALI & M. WHEELER, “Modified Macdonald polynomials and integrability”, *Commun. Math. Phys.* **374** (2020), no. 3, p. 1809-1876.
- [25] C. KORFF, “Cylindric versions of specialised Macdonald functions and a deformed Verlinde algebra”, *Commun. Math. Phys.* **318** (2013), no. 1, p. 173-246.
- [26] S. KOROTKIKH, “Hidden diagonal integrability of q -Hahn vertex model and Beta polymer model”, <https://arxiv.org/abs/2105.05058>, 2021.

- [27] A. KUNIBA, V. MANGAZEEV, S. MARUYAMA & M. OKADO, “Stochastic R matrix for $U_q(A_n^{(1)})$ ”, *Nucl. Phys., B* **913** (2016), p. 248-277.
- [28] I. G. MACDONALD, *Symmetric functions and Hall polynomials*, 2nd ed., Oxford Mathematical Monographs, Oxford Science Publications, 1995, x+475 pages.
- [29] V. MANGAZEEV, “On the Yang-Baxter equation for the six-vertex model”, *Nucl. Phys., B* **882** (2014), p. 70-96.
- [30] K. MOTEGI, “Combinatorial properties of symmetric polynomials from integrable vertex models in finite lattice”, *J. Math. Phys.* **58** (2017), no. 9, article no. 091703 (23 pages).
- [31] K. MOTEGI & K. SAKAI, “Vertex models, TASEP and Grothendieck polynomials”, *J. Phys. A: Math. Theor.* **46** (2013), no. 35, article no. 355201 (26 pages).
- [32] M. MUCCICONI & L. PETROV, “Spin q -Whittaker polynomials and deformed quantum Toda”, <https://arxiv.org/abs/2003.14260>, 2020.
- [33] L. PETROV, “Refined Cauchy identity for spin Hall–Littlewood symmetric rational functions”, <https://arxiv.org/abs/2007.10886>, 2007.
- [34] J. R. STEMBRIDGE, “A short proof of Macdonald’s conjecture for the root systems of type A ”, *Proc. Am. Math. Soc.* **102** (1988), no. 4, p. 777-785.
- [35] N. V. TSILEVICH, “Quantum inverse scattering method for the q -boson model and symmetric functions”, *Funct. Anal. Appl.* **40** (2006), no. 3, p. 207-217.
- [36] M. WHEELER & P. ZINN-JUSTIN, “Refined Cauchy/Littlewood identities and six-vertex model partition functions. III. Deformed bosons”, *Adv. Math.* **299** (2016), p. 543-600.
- [37] ———, “Hall polynomials, inverse Kostka polynomials and puzzles”, *J. Comb. Theory, Ser. A* **159** (2018), p. 107-163.
- [38] P. ZINN-JUSTIN, “Six-vertex, loop and tiling models: integrability and combinatorics”, Habilitation thesis, <http://www.lpthe.jussieu.fr/~pzinn/publi/hdr.pdf>, <https://arxiv.org/abs/0901.0665>.