



Annales de la Faculté des Sciences de Toulouse

MATHÉMATIQUES

CHRISTINE HUYGHE AND TOBIAS SCHMIDT

Kashiwara's theorem for twisted arithmetic differential operators

Tome XXXIII, n° 2 (2024), p. 311–347.

<https://doi.org/10.5802/afst.1772>

© les auteurs, 2024.

Les articles des *Annales de la Faculté des Sciences de Toulouse* sont mis à disposition sous la licence Creative Commons Attribution (CC-BY) 4.0
<http://creativecommons.org/licenses/by/4.0/>



Publication membre du centre
Mersenne pour l'édition scientifique ouverte
<http://www.centre-mersenne.org/>
e-ISSN : 2258-7519

Kashiwara’s theorem for twisted arithmetic differential operators ^(*)

CHRISTINE HUYGHE ⁽¹⁾ AND TOBIAS SCHMIDT ⁽²⁾

ABSTRACT. — We establish a version of Kashiwara’s theorem for twisted sheaves of Berthelot’s arithmetic differential operators for a closed immersion between smooth p -adic formal schemes. As an application, we give a geometric construction of simple modules for crystalline distribution algebras of reductive groups.

RÉSUMÉ. — On établit une version du théorème de Kashiwara (relative à une immersion fermée entre deux schémas formels p -adiques) pour les faisceaux tordus des opérateurs différentiels arithmétiques de Berthelot. Comme application de ce théorème, nous donnons une construction géométrique des modules simples sur une algèbre de distributions arithmétiques d’un groupe réductif.

1. Introduction

Let X be a smooth complex variety and $Y \subset X$ a smooth closed subvariety. A basic result in \mathcal{D} -module theory is Kashiwara’s theorem which states an equivalence of categories between the category of \mathcal{D}_Y -modules, quasi-coherent over \mathcal{O}_Y , and the category of \mathcal{D}_X -modules, quasi-coherent over \mathcal{O}_X , with support in Y . Recall that a *twisted sheaf of differential operators* on X is a sheaf of rings \mathcal{A} on X together with a ring homomorphism $\iota : \mathcal{O}_X \rightarrow \mathcal{A}$ such that the pair (ι, \mathcal{A}) is locally isomorphic to the pair $(\text{can}, \mathcal{D}_X)$ where $\text{can} : \mathcal{O}_X \rightarrow \mathcal{D}_X$ is the canonical inclusion. Originally, such twisted sheaves were introduced in the early 1980s by Beilinson–Bernstein in order to study localisations of Lie algebra representations with general infinitesimal character on complex flag varieties [1]. It is well-known that the

^(*) Reçu le 9 juin 2021, accepté le 25 avril 2022.

⁽¹⁾ IRMA, Université de Strasbourg, 7 rue René Descartes, 67084 Strasbourg cedex, France — huyghe@math.unistra.fr

⁽²⁾ IRMAR, Université de Rennes 1, Campus Beaulieu, 35042 Rennes cedex, France — tobias.schmidt@univ-rennes1.fr — toschmidt@uni-wuppertal.de

Article proposé par Damian Rössler.

right module version of Kashiwara’s theorem generalizes to twisted sheaves of differential operators [16, 4.3]. Under additional hypotheses on the twisted sheaf \mathcal{A} (e.g. an analogue of the order filtration on \mathcal{A}) one may establish side-changing functors in the general setting of \mathcal{A} -modules and then deduce a version of Kashiwara’s theorem for left modules [15, 2.3].

In the arithmetic setting, let \mathfrak{o} denote a complete discrete valuation ring of mixed characteristic $(0, p)$ with uniformizer ϖ , field of fractions K and perfect residue field. Let \mathfrak{X} be a smooth formal scheme over \mathfrak{o} and let $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ be the sheaf of arithmetic differential operators on \mathfrak{X} [2]. If $\mathfrak{Y} \subset \mathfrak{X}$ is a closed smooth formal subscheme, Berthelot’s version of Kashiwara’s theorem gives an equivalence between the category of coherent left $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -modules and the category of coherent left $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ -modules with support in \mathfrak{Y} . Berthelot gives a proof of the theorem in his course on arithmetic \mathcal{D} -modules 1997 at the Centre Emile Borel, which, however, is not included in the corresponding reference [4]. In the appendix of [7] Caro establishes a version of the theorem for log structures and coefficients.

Our goal in this paper is to establish a version of Kashiwara’s theorem for twisted sheaves of arithmetic differential operators. Similar to the complex analytic setting, such twisted sheaves appear naturally in the context of the localization theory of crystalline distribution algebras of reductive groups, when varying the infinitesimal character of representations [13, 18]. Following Beilinson–Bernstein, we define a *twisted sheaf of arithmetic differential operators* to be a sheaf of rings \mathcal{A} on \mathfrak{X} together with a ring homomorphism $\iota : \mathcal{O}_{\mathfrak{X}, \mathbb{Q}} \rightarrow \mathcal{A}$ such that the pair (ι, \mathcal{A}) is locally isomorphic to the pair $(\text{can}, \mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger)$ where $\text{can} : \mathcal{O}_{\mathfrak{X}, \mathbb{Q}} \rightarrow \mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ is the canonical inclusion. At this level of generality, as we have explained above, there are no side-changing functors and one may only hope for a right module version of Kashiwara’s theorem.

To formulate our main result, let \mathcal{A} be a twisted sheaf of differential operators on \mathfrak{X} and let $i : \mathfrak{Y} \rightarrow \mathfrak{X}$ be the inclusion of a closed smooth formal subscheme defined by the ideal $\mathcal{I} \subset \mathcal{O}_{\mathfrak{X}}$. Let $\mathcal{N}_{\mathcal{A}}(\mathcal{I}\mathcal{A})$ be the normalizer of the right ideal generated by \mathcal{I} in \mathcal{A} . We show that

$$\mathcal{A}_{\mathfrak{Y}} := i^{-1}(\mathcal{N}_{\mathcal{A}}(\mathcal{I}\mathcal{A})/\mathcal{I}\mathcal{A}) \quad \text{resp.} \quad \mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}} := i^*(\mathcal{A})$$

are a twisted sheaf of arithmetic differential operators on \mathfrak{Y} resp. a $(\mathcal{A}_{\mathfrak{Y}}, i^{-1}\mathcal{A})$ -bimodule. We obtain an adjoint pair of functors $(i_{\mathcal{A}, +}, i_{\mathcal{A}}^{\natural})$ between the categories of right modules over $\mathcal{A}_{\mathfrak{Y}}$ and \mathcal{A} respectively: the direct image

$$i_{\mathcal{A}, +}\mathcal{N} := i_*(\mathcal{N} \otimes_{\mathcal{A}_{\mathfrak{Y}}} \mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}})$$

from right $\mathcal{A}_{\mathfrak{y}}$ -modules to right \mathcal{A} -modules and, in the opposite direction, the restriction functor

$$i_{\mathcal{A}}^{\natural} \mathcal{M} := \mathcal{H}om_{i^{-1}\mathcal{A}}(\mathcal{A}_{\mathfrak{y}} \rightarrow \mathfrak{x}, i^{-1}\mathcal{M}).$$

Let $\text{Coh}^r(\mathcal{A}_{\mathfrak{y}})$ and $\text{Coh}^{r,\mathfrak{y}}(\mathcal{A})$ be the categories of coherent right $\mathcal{A}_{\mathfrak{y}}$ -modules and coherent right \mathcal{A} -modules with support in \mathfrak{y} , respectively. Our main result is the following.

THEOREM (cf. Theorem 4.16). — *The functors $i_{\mathcal{A},+}$, $i_{\mathcal{A}}^{\natural}$ induce quasi-inverse equivalences of categories*

$$\text{Coh}^r(\mathcal{A}_{\mathfrak{y}}) \begin{array}{c} \xrightarrow{i_{\mathcal{A},+}} \\ \xleftarrow[\simeq]{i_{\mathcal{A}}^{\natural}} \end{array} \text{Coh}^{r,\mathfrak{y}}(\mathcal{A}).$$

For the proof, we proceed as follows. We first establish some complements on arithmetic differential operators, notably the normalizer description for operators on closed subspaces. We then give a full and self-contained proof of the Berthelot–Kashiwara theorem for left $\mathcal{D}_{\mathfrak{x},\mathbb{Q}}^{\dagger}$ -modules. Note that Caro's logarithmic Kashiwara theorem for coefficients [7] contains this result as a special case. However, this special case is buried under a lot of additional notation. We therefore believe that it is instructive, and a useful basis for our future work, to have a clear direct proof in this special case, using only the tools of the basic reference [2]. As in [7], the key ingredient is a lemma of Berthelot on a certain matrix identity involving matrices over arithmetic differential operators. We give a full proof of this lemma in Section 2.5 (in [7] only the rank 1-case is really considered). We then use side-changing functors to obtain Berthelot–Kashiwara for right $\mathcal{D}_{\mathfrak{x},\mathbb{Q}}^{\dagger}$ -modules. Finally, we prove sufficiently many properties and compatibilities of the adjoint pair $(i_{\mathcal{A},+}, i_{\mathcal{A}}^{\natural})$ to reduce the proof of the main theorem to a local situation. This allows us to undo the twist and then conclude via right Berthelot–Kashiwara.

In the final section, we give an application to the representation theory of crystalline distributions algebras. We fix a connected split reductive group G over \mathfrak{o} and denote its ϖ -adic completion by \mathcal{G} . Let $D^{\dagger}(\mathcal{G})_{\mathbb{Q}}$ be its crystalline distribution algebra, as introduced and studied in [12]. Irreducible modules over $D^{\dagger}(\mathcal{G})_{\mathbb{Q}}$ can be considered as local data for certain admissible locally analytic $G(K)$ -representations and thus, are of interest in the so-called p -adic local Langlands programme. We let \mathfrak{X} be the formal flag variety of G . In [18] Sarrazola–Alzate generalizes a classical construction of Beilinson–Bernstein [1] and Borho–Brylinski [5] to the arithmetic setting and constructs a family of twisted sheaves of arithmetic differential operators $\mathcal{D}_{\mathfrak{x},\mathbb{Q},\lambda}^{\dagger}$ on \mathfrak{X} , indexed by certain characters λ of a Cartan subalgebra of $\text{Lie}(G) \otimes \mathbb{Q}$. We

apply our Berthelot–Kashiwara theorem to the sheaves $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger$. For algebraic λ (i.e. when λ lifts to a character of a maximal split torus in G) this leads to a class of simple $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger$ -modules, parametrized by closed smooth subschemes $\mathfrak{Y} \subset \mathfrak{X}$. By the arithmetic localisation theorem [13], their global sections give rise to simple $D^\dagger(\mathcal{G})_{\mathbb{Q}}$ -modules.

Acknowledgements

We thank Daniel Caro for having made available to us a very preliminary version of the article [7] and for helpful discussions regarding certain points in this article. We also thank the anonymous referee, whose comments have improved the exposition of the material at several places.

Notations and Conventions

Throughout the article, \mathfrak{o} denotes a complete discrete valuation ring with mixed characteristic $(0, p)$. We denote by K its fraction field and by k its residue field, which is assumed to be perfect. We let ϖ be a uniformizer of \mathfrak{o} . A formal scheme \mathfrak{X} over \mathfrak{o} which is locally noetherian and such that $\varpi \mathcal{O}_{\mathfrak{X}}$ is an ideal of definition is called an \mathfrak{o} -formal scheme. We denote its special fibre by \mathfrak{X}_s .

2. Complements on arithmetic differential operators

2.1. Arithmetic differential operators

In this subsection we assume a certain familiarity with the basic notions of divided powers and divided power envelopes. Our basic references are [2, 3].

Let \mathfrak{X} be an \mathfrak{o} -formal scheme, which is smooth over $\mathfrak{S} := \mathrm{Spf}(\mathfrak{o})$, with structure sheaf $\mathcal{O}_{\mathfrak{X}}$.

Let $m \geq 0$. Let $\mathcal{P}_{\mathfrak{X}/\mathfrak{S}, (m)}^n$, $n \geq 0$ be the projective system of sheaves of principal parts of level m and order n of \mathfrak{X} relative to \mathfrak{S} . There are two morphisms $p_0, p_1 : \mathcal{P}_{\mathfrak{X}/\mathfrak{S}, (m)}^n \rightarrow \mathcal{O}_{\mathfrak{X}}$, induced from the two projection morphisms $\mathfrak{X} \times \mathfrak{X} \rightarrow \mathfrak{X}$, making $\mathcal{P}_{\mathfrak{X}/\mathfrak{S}, (m)}^n$ a commutative $\mathcal{O}_{\mathfrak{X}}$ -algebra in two ways, on the left (via p_0) and on the right (via p_1). The two structures are denoted by $d_i : \mathcal{O}_{\mathfrak{X}} \rightarrow \mathcal{P}_{\mathfrak{X}/\mathfrak{S}, (m)}^n$ for $i = 0, 1$. If \mathfrak{X} has étale coordinates

t_1, \dots, t_d and $\tau_i := p_1^*(t_i) - p_0^*(t_i) \in \mathcal{O}_{\mathfrak{X} \times \mathfrak{X}}$, then $\mathcal{P}_{\mathfrak{X}/\mathfrak{S},(m)}^n$ is a free $\mathcal{O}_{\mathfrak{X}}$ -module (for both its left and right structure) on the basis $\underline{\tau}^{\{k\}} := \tau_1^{\{k_1\}} \dots \tau_d^{\{k_d\}}$ for $|k| \leq n$.

The sheaf of arithmetic differential operators on \mathfrak{X} of level m and order n is the $\mathcal{O}_{\mathfrak{X}}$ -dual $\mathcal{D}_{\mathfrak{X},n}^{(m)} := \mathcal{H}om_{\mathcal{O}_{\mathfrak{X}}}(\mathcal{P}_{\mathfrak{X}/\mathfrak{S},(m)}^n, \mathcal{O}_{\mathfrak{X}})$. An element $P \in \mathcal{D}_{\mathfrak{X},n}^{(m)}$ acts on $\mathcal{O}_{\mathfrak{X}}$ via the composition

$$\mathcal{O}_{\mathfrak{X}} \xrightarrow{d_1} \mathcal{P}_{\mathfrak{X}/\mathfrak{S},(m)}^n \xrightarrow{P} \mathcal{O}_{\mathfrak{X}}.$$

The union $\mathcal{D}_{\mathfrak{X}}^{(m)} := \bigcup_n \mathcal{D}_{\mathfrak{X},n}^{(m)}$ is a ring and there is a natural ring homomorphism $\mathcal{D}_{\mathfrak{X}}^{(m)} \rightarrow \mathcal{D}_{\mathfrak{X}}^{(m+1)}$. We denote by $\widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)} = \varprojlim_i \mathcal{D}_{\mathfrak{X}}^{(m)}/\varpi^i$ the ϖ -adic completion and let

$$\mathcal{D}_{\mathfrak{X}}^{\dagger} := \varinjlim_m \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)} \quad \text{and} \quad \mathcal{D}_{\mathfrak{X},\mathbb{Q}}^{\dagger} := \mathcal{D}_{\mathfrak{X}}^{\dagger} \otimes_{\mathbb{Z}} \mathbb{Q}.$$

We shall also need to consider the usual (i.e. with divided powers) ring of algebraic differential operators $\mathcal{D}_{\mathfrak{X}} := \varinjlim_m \mathcal{D}_{\mathfrak{X}}^{(m)}$ on the \mathfrak{o} -formal scheme \mathfrak{X} , cf. [9, 16.8]. We denote by $\widehat{\mathcal{D}}_{\mathfrak{X}}$ its ϖ -adic completion.

It will be useful to make the following definition.

DEFINITION 2.1. — *An $\mathcal{O}_{\mathfrak{X}}$ -ring is a pair $(\mathcal{A}, \iota_{\mathcal{A}})$ consisting of a sheaf of rings \mathcal{A} on \mathfrak{X} , together with a ring morphism $\iota_{\mathcal{A}} : \mathcal{O}_{\mathfrak{X}} \rightarrow \mathcal{A}$. A morphism between two $\mathcal{O}_{\mathfrak{X}}$ -rings $(\mathcal{A}, \iota_{\mathcal{A}})$ and $(\mathcal{A}', \iota_{\mathcal{A}'})$ is a morphism $f : \mathcal{A} \rightarrow \mathcal{A}'$ of sheaves of rings satisfying $f \circ \iota_{\mathcal{A}} = \iota_{\mathcal{A}'}$.*

One obtains thereby a category of $\mathcal{O}_{\mathfrak{X}}$ -rings. In situations where the morphism of rings $\iota_{\mathcal{A}}$ is understood, we will often drop it from the notation and just write \mathcal{A} instead of $(\mathcal{A}, \iota_{\mathcal{A}})$.

There are obvious variants of this category when the structure sheaf $\mathcal{O}_{\mathfrak{X}}$ is replaced by another sheaf associated with the formal scheme \mathfrak{X} , such as $\mathcal{O}_{\mathfrak{X},\mathbb{Q}}$ or $\mathcal{O}_{\mathfrak{X},\mathbb{Q}}/\mathcal{I}$ (for an ideal sheaf $\mathcal{I} \subseteq \mathcal{O}_{\mathfrak{X},\mathbb{Q}}$).

All the rings $\mathcal{D}_{\mathfrak{X}}^{(m)}$, $\mathcal{D}_{\mathfrak{X}}$, $\widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)}$, $\widehat{\mathcal{D}}_{\mathfrak{X}}$, $\mathcal{D}_{\mathfrak{X}}^{\dagger}$ are $\mathcal{O}_{\mathfrak{X}}$ -rings, and $\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^{\dagger}$ is even an $\mathcal{O}_{\mathfrak{X},\mathbb{Q}}$ -ring.

2.2. Side-changing functors

Let \mathfrak{X} be an \mathfrak{o} -formal scheme of relative dimension d , which is smooth over $\mathfrak{S} := \mathrm{Spf}(\mathfrak{o})$. Let

$$\omega_{\mathfrak{X}} := \bigwedge^d \Omega_{\mathfrak{X}/\mathfrak{S}}^1$$

be the module of differentials of highest degree, with its natural right $\mathcal{D}_{\mathfrak{X}}$ -action [3, 1.2.1]. As in the classical setting, the functors $\omega_{\mathfrak{X}} \otimes_{\mathcal{O}_{\mathfrak{X}}} (-)$ and $\mathcal{H}\text{om}_{\mathcal{O}_{\mathfrak{X}}}(\omega_{\mathfrak{X}}, -)$ induce mutually inverse equivalences of categories between left and right $\mathcal{D}_{\mathfrak{X}}^{(m)}$ -modules, for any $m \geq 0$ [3, 1.2.7(c)]. The following proposition is certainly well-known, we record it for a lack of reference.

PROPOSITION 2.2. — *Let $Y \subseteq \mathfrak{X}_s$ be a closed subset. The functors $\omega_{\mathfrak{X}} \otimes_{\mathcal{O}_{\mathfrak{X}}} (-)$ and $\mathcal{H}\text{om}_{\mathcal{O}_{\mathfrak{X}}}(\omega_{\mathfrak{X}}, -)$ induce mutually inverse equivalences of categories between left and right $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^{\dagger}$ -modules supported on Y .*

Proof. — Since $\omega_{\mathfrak{X}}$ is a coherent $\mathcal{O}_{\mathfrak{X}}$ -module, its spaces of local sections over affine opens in \mathfrak{X} are ϖ -adically complete. Its $\mathcal{D}_{\mathfrak{X}}$ -action extends therefore to a $\widehat{\mathcal{D}}_{\mathfrak{X}}$ -action. The functors $\omega_{\mathfrak{X}} \otimes_{\mathcal{O}_{\mathfrak{X}}} (-)$ and $\mathcal{H}\text{om}_{\mathcal{O}_{\mathfrak{X}}}(\omega_{\mathfrak{X}}, -)$ descend therefore to equivalences between left and right $\widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)}$ -modules. Inverting p and passing to the inductive limit over all $m \geq 0$ yields the proposition in the case $Y = \mathfrak{X}_s$. The general case follows from the fact that both functors $\omega_{\mathfrak{X}} \otimes_{\mathcal{O}_{\mathfrak{X}}} (-)$ and $\mathcal{H}\text{om}_{\mathcal{O}_{\mathfrak{X}}}(\omega_{\mathfrak{X}}, -) = (-) \otimes_{\mathcal{O}_{\mathfrak{X}}} \omega_{\mathfrak{X}}^{-1}$ preserve the support condition. \square

2.3. Operators on closed subspaces

Let

$$i : \mathfrak{Y} \longrightarrow \mathfrak{X}$$

be a closed immersion between two \mathfrak{o} -formal schemes, which are smooth over $\mathfrak{S} := \text{Spf}(\mathfrak{o})$. Let $r := \dim \mathfrak{Y}$ and $d := \dim \mathfrak{X}$.

It is well-known that the adjoint pair of functors (i_*, i^{-1}) induces an equivalence of categories between abelian sheaves on \mathfrak{Y} and abelian sheaves on \mathfrak{X} with support in \mathfrak{Y} . We denote by i^* the functor $\mathcal{O}_{\mathfrak{Y}} \otimes_{i^{-1}\mathcal{O}_{\mathfrak{X}}} i^{-1}(-)$ from $\mathcal{O}_{\mathfrak{X}}$ -modules to $\mathcal{O}_{\mathfrak{Y}}$ -modules.

Let $\mathcal{I} \subseteq \mathcal{O}_{\mathfrak{X}}$ be the ideal sheaf defining the closed immersion i . There is the Hom sheaf $\mathcal{H}\text{om}_{\mathfrak{o}}(i^{-1}\mathcal{I}, \mathcal{O}_{\mathfrak{Y}})$ on \mathfrak{Y} , which is an $\mathcal{O}_{\mathfrak{Y}}$ -module via multiplication on the target, i.e. $(sf)(a) := s(f(a))$ for local sections $s \in \mathcal{O}_{\mathfrak{Y}}$, $f \in \mathcal{H}\text{om}_{\mathfrak{o}}(i^{-1}\mathcal{I}, \mathcal{O}_{\mathfrak{Y}})$, and $a \in i^{-1}\mathcal{I}$. Similarly, the sheaf $\mathcal{H}\text{om}_K(i^{-1}\mathcal{I}_{\mathbb{Q}}, \mathcal{O}_{\mathfrak{Y}, \mathbb{Q}})$ is an $\mathcal{O}_{\mathfrak{Y}, \mathbb{Q}}$ -module.

LEMMA 2.3. — *Let $\mathcal{A} \in \{\mathcal{D}_{\mathfrak{X}}^{(m)}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)}, \mathcal{D}_{\mathfrak{X}}^{\dagger}\}$. The restriction map $\mathcal{A} \rightarrow \mathcal{H}\text{om}_{\mathfrak{o}}(\mathcal{I}, \mathcal{O}_{\mathfrak{X}})$, $P \mapsto P|_{\mathcal{I}}$ induces a $\mathcal{O}_{\mathfrak{Y}}$ -linear morphism*

$$i^* \mathcal{A} \longrightarrow \mathcal{H}\text{om}_{\mathfrak{o}}(i^{-1}\mathcal{I}, \mathcal{O}_{\mathfrak{Y}}).$$

Proof. — The morphism $i^{\sharp} : i^{-1}\mathcal{O}_{\mathfrak{X}} \rightarrow \mathcal{O}_{\mathfrak{Y}}$ induces a morphism

$$\mathcal{H}\text{om}_{\mathfrak{o}}(i^{-1}\mathcal{I}, i^{-1}\mathcal{O}_{\mathfrak{X}}) \longrightarrow \mathcal{H}\text{om}_{\mathfrak{o}}(i^{-1}\mathcal{I}, \mathcal{O}_{\mathfrak{Y}}).$$

Let $\mathcal{A} \in \{\mathcal{D}_{\mathfrak{X}}^{(m)}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)}, \mathcal{D}_{\mathfrak{X}}^{\dagger}\}$. Applying i^{-1} to the restriction morphism $\mathcal{A} \rightarrow \mathcal{H}\text{om}_{\mathfrak{o}}(\mathcal{I}, \mathcal{O}_{\mathfrak{X}})$ and composing with the above morphism yields a morphism $i^{-1}\mathcal{A} \rightarrow \mathcal{H}\text{om}_{\mathfrak{o}}(i^{-1}\mathcal{I}, \mathcal{O}_{\mathfrak{Y}})$. Since the latter is $i^{-1}\mathcal{O}_{\mathfrak{X}}$ -linear and the target a $\mathcal{O}_{\mathfrak{Y}}$ -module, it extends to a $\mathcal{O}_{\mathfrak{Y}}$ -linear morphism $i^*\mathcal{A} \rightarrow \mathcal{H}\text{om}_{\mathfrak{o}}(i^{-1}\mathcal{I}, \mathcal{O}_{\mathfrak{Y}})$, as claimed. \square

We call the morphism appearing in the lemma $\theta^{(m)}, \widehat{\theta}^{(m)}, \theta^{\dagger}$ in the cases $\mathcal{D}_{\mathfrak{X}}^{(m)}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)}, \mathcal{D}_{\mathfrak{X}}^{\dagger}$ respectively.

According to [2, 2.1.4.3], for any $n \geq 0$, there is a canonical $\mathcal{O}_{\mathfrak{Y}}$ -linear morphism

$$\mathcal{P}_{(m)}^n(i) : i^*\mathcal{P}_{\mathfrak{X}/\mathfrak{S},(m)}^n \longrightarrow \mathcal{P}_{\mathfrak{Y}/\mathfrak{S},(m)}^n.$$

Dualising and taking the union over all n yields a $\mathcal{O}_{\mathfrak{Y}}$ -linear morphism $\mathcal{D}_{\mathfrak{Y}}^{(m)} \rightarrow i^*\mathcal{D}_{\mathfrak{X}}^{(m)}$. Completing ϖ -adically, taking the union over all $m \geq 0$ and finally inverting p yields a $\mathcal{O}_{\mathfrak{Y},\mathbb{Q}}$ -linear morphism $\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^{\dagger} \rightarrow i^*\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^{\dagger}$.

PROPOSITION 2.4. — *There is an exact sequence of $\mathcal{O}_{\mathfrak{Y},\mathbb{Q}}$ -modules*

$$0 \longrightarrow \mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^{\dagger} \longrightarrow i^*\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^{\dagger} \xrightarrow{\theta_{\mathbb{Q}}^{\dagger}} \mathcal{H}\text{om}_K(i^{-1}\mathcal{I}_{\mathbb{Q}}, \mathcal{O}_{\mathfrak{Y},\mathbb{Q}}).$$

Proof. — The exactness is a local question and we may assume that \mathfrak{X} is affine. We let $A := \mathcal{O}(\mathfrak{X})$ and $I := \mathcal{O}(\mathcal{I})$. By the Jacobi criterion, e.g. [6, Prop. 3.5], we may even assume that \mathfrak{X} has étale coordinates $t_1, \dots, t_d \in A$ such that

- the images of t_1, \dots, t_r in $\bar{A} = A/I$ are étale coordinates for \mathfrak{Y} ,
- the ideal $I \subseteq A$ is generated by t_{r+1}, \dots, t_d .

Fix $m \geq 0$ and let $P_{A/\mathfrak{o},(m)}^n = \mathcal{O}(\mathcal{P}_{\mathfrak{X}/\mathfrak{S},(m)}^n)$ and $\mathcal{D}_A^{(m)} = \mathcal{O}(\mathcal{D}_{\mathfrak{X}}^{(m)})$. There is a natural morphism of left \bar{A} -modules

$$0 \longrightarrow \ker(\delta) \longrightarrow \bar{A} \otimes_A P_{A/\mathfrak{o},(m)}^n \xrightarrow{\delta} P_{A/\mathfrak{o},(m)}^n \longrightarrow 0$$

where the map δ equals the global sections of the morphism $i^*\mathcal{P}_{\mathfrak{X}/\mathfrak{S},(m)}^n \rightarrow \mathcal{P}_{\mathfrak{Y}/\mathfrak{S},(m)}^n$. Since δ is A -linear on the right, we have $\bar{A} \otimes_A (P_{A/\mathfrak{o},(m)}^n I) \subseteq \ker(\delta)$. The \bar{A} -linear morphism δ maps the elements $1 \otimes (\tau_1^{\{k_1\}} \dots \tau_r^{\{k_r\}})$ with $k_1 + \dots + k_r \leq n$ bijectively to a \bar{A} -basis of $P_{A/\mathfrak{o},(m)}^n$ and so is indeed surjective. Moreover, we have

$$\ker(\delta) = \left\{ \sum_{\underline{k} \in \mathbb{N}^d \setminus \mathbb{N}^r} \bar{A} \tau^{\{\underline{k}\}} \right\},$$

where $\underline{k} \notin \mathbb{N}^r$ means that $k_j > 0$ for some $j > r$. In particular, all \bar{A} -modules in the above short exact sequence are free and dualizing gives the

exact sequence

$$0 \longrightarrow \mathrm{Hom}_{\bar{A}}(P_{\bar{A}/\mathfrak{o},(m)}^n, \bar{A}) \longrightarrow \mathrm{Hom}_{\bar{A}}(\bar{A} \otimes_A P_{A/\mathfrak{o},(m)}^n, \bar{A}) \xrightarrow{\delta^*} \mathrm{Hom}_{\bar{A}}(\ker(\delta), \bar{A}) \longrightarrow 0.$$

Since \bar{A} is flat over \mathbb{Z} , the localization map $\mathrm{Hom}_{\bar{A}}(\ker(\delta), \bar{A}) \rightarrow \mathrm{Hom}_{\bar{A}_{\mathbb{Q}}}(\ker(\delta)_{\mathbb{Q}}, \bar{A}_{\mathbb{Q}})$ is injective and we have the exact sequence

$$0 \longrightarrow \mathrm{Hom}_{\bar{A}}(P_{\bar{A}/\mathfrak{o},(m)}^n, \bar{A}) \longrightarrow \mathrm{Hom}_{\bar{A}}(\bar{A} \otimes_A P_{A/\mathfrak{o},(m)}^n, \bar{A}) \xrightarrow{\delta^*} \mathrm{Hom}_{\bar{A}_{\mathbb{Q}}}(\ker(\delta)_{\mathbb{Q}}, \bar{A}_{\mathbb{Q}}).$$

The formula $q! \tau^{\{k\}} = \tau^k$, where q is the euclidian division of k by p^m [2, 1.3.5.2], shows that

$$\ker(\delta)_{\mathbb{Q}} = \left\{ \sum_{k \in \mathbb{N}^d \setminus \mathbb{N}^r} \bar{A}_{\mathbb{Q}} \underline{\tau}^k \right\}.$$

On the other hand, given $t_i \in I$ (i.e. $i > r$), the image of $\tau_i^k = (1 \otimes t_i - t_i \otimes 1)^k \in A \otimes_{\mathfrak{o}} A$ in the quotient $\bar{A} \otimes_{\mathfrak{o}} A$ equals $\bar{1} \otimes t_i^k$. Hence, the image of the set $\{d_1(f) \mid f \in I\} \subset P_{A/\mathfrak{o},(m)}^n$ in $\bar{A}_{\mathbb{Q}} \otimes_A P_{A/\mathfrak{o},(m)}^n$ generates the $\bar{A}_{\mathbb{Q}}$ -module $\ker(\delta)_{\mathbb{Q}}$. For a given element

$$\begin{aligned} \bar{P} &= P + I \mathcal{D}_{A,n}^{(m)} \in \mathrm{Hom}_{\bar{A}}(\bar{A} \otimes_A P_{A/\mathfrak{o},(m)}^n, \bar{A}) \\ &= \bar{A} \otimes_A \mathrm{Hom}_A(P_{A/\mathfrak{o},(m)}^n, A) = \mathcal{D}_{A,n}^{(m)} / I \mathcal{D}_{A,n}^{(m)} \end{aligned}$$

we therefore have

$$\delta^*(\bar{P}) = 0 \quad \text{if and only if} \quad P \circ d_1(I) \subseteq I_{\mathbb{Q}}.$$

Since A/I is p -torsionfree, one has $A \cap I_{\mathbb{Q}} = I$ and so the condition $P \circ d_1(I) \subseteq I_{\mathbb{Q}}$ is equivalent to $P \circ d_1(I) \subset A \cap I_{\mathbb{Q}} = I$. In other words, the sequence

$$0 \longrightarrow \mathrm{Hom}_{\bar{A}}(P_{\bar{A}/\mathfrak{o},(m)}^n, \bar{A}) \longrightarrow \mathrm{Hom}_{\bar{A}}(\bar{A} \otimes_A P_{A/\mathfrak{o},(m)}^n, \bar{A}) \longrightarrow \mathrm{Hom}_{\mathfrak{o}}(I, \bar{A})$$

is exact. Taking the union over $n \geq 0$ yields the exact sequence

$$0 \longrightarrow \mathcal{D}_{\bar{A}}^{(m)} \longrightarrow \mathcal{D}_A^{(m)} / I \mathcal{D}_A^{(m)} \xrightarrow{\theta^{(m)}} \mathrm{Hom}_{\mathfrak{o}}(I, \bar{A}),$$

where $\theta^{(m)}$ is the global sections of the map appearing in Lemma 2.3 for $\mathcal{A} = \mathcal{D}_{\mathfrak{x}}^{(m)}$. By left-exactness of ϖ -adic completion, we obtain the exact sequence

$$0 \longrightarrow \widehat{\mathcal{D}_{\bar{A}}^{(m)}} \longrightarrow \widehat{\mathcal{D}_A^{(m)} / I \mathcal{D}_A^{(m)}} \xrightarrow{\widehat{\theta^{(m)}}} \widehat{\mathrm{Hom}_{\mathfrak{o}}(I, \bar{A})}.$$

Since ϖ -adic completion is exact on finitely generated modules over the noetherian ring $\mathcal{D}_A^{(m)}$, the completion of $\mathcal{D}_A^{(m)}/I\mathcal{D}_A^{(m)}$ equals $\widehat{\mathcal{D}}_A^{(m)}/I\widehat{\mathcal{D}}_A^{(m)}$. Therefore, $\widehat{\theta}^{(m)}$ admits the factorisation

$$\widehat{\mathcal{D}}_A^{(m)}/I\widehat{\mathcal{D}}_A^{(m)} \xrightarrow{\widehat{\theta}^{(m)}} \mathrm{Hom}_{\mathfrak{o}}(I, \bar{A}) \longrightarrow \widehat{\mathrm{Hom}_{\mathfrak{o}}(I, \bar{A})}$$

where $\widehat{\theta}^{(m)}$ is the global sections of the map appearing in Lemma 2.3 for $\mathcal{A} = \widehat{\mathcal{D}}_{\mathfrak{x}}^{(m)}$. Since \bar{A} is ϖ -adically complete, the \mathfrak{o} -module $\mathrm{Hom}_{\mathfrak{o}}(I, \bar{A})$ is ϖ -adically separated and hence injects into its ϖ -adic completion. Therefore the induced sequence

$$0 \longrightarrow \widehat{\mathcal{D}}_A^{(m)} \longrightarrow \widehat{\mathcal{D}}_A^{(m)}/I\widehat{\mathcal{D}}_A^{(m)} \xrightarrow{\widehat{\theta}^{(m)}} \mathrm{Hom}_{\mathfrak{o}}(I, \bar{A})$$

remains exact. Taking an inductive limit over $m \geq 0$ and inverting p yields the claim. \square

DEFINITION 2.5. — *Let $\mathcal{I} \subseteq \mathcal{O}_{\mathfrak{x}, \mathbb{Q}}$ be an ideal. The normalizer of the ideal $\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}$ is the subset of $\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}$ equal to*

$$\mathcal{N}(\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}) := \mathcal{N}_{\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}}(\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}) := \{P \in \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger} \mid P \mathcal{I} \subseteq \mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}\}.$$

LEMMA 2.6. — *One has the following basic properties:*

- (i) $\mathcal{N}(\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger})$ is a sub- $\mathcal{O}_{\mathfrak{x}, \mathbb{Q}}$ -ring of $\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}$.
- (ii) $\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger} \subseteq \mathcal{N}(\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger})$ is a two-sided ideal.
- (iii) The quotient $\mathcal{N}(\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger})/\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}$ is an $\mathcal{O}_{\mathfrak{x}, \mathbb{Q}}/\mathcal{I}$ -ring.

Proof. — Let $\mathcal{N} := \mathcal{N}(\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger})$. It is clear that \mathcal{N} contains $\mathcal{O}_{\mathfrak{x}, \mathbb{Q}}$ and is stable under addition. Given $P, Q \in \mathcal{N}$, one has

$$(PQ) \mathcal{I} \subseteq P(Q \mathcal{I}) \subseteq P(\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}) \subseteq (P \mathcal{I}) \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger} \subseteq \mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger},$$

and so $PQ \in \mathcal{N}$. This shows (i). For (ii), note that $\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}$ is certainly contained in \mathcal{N} as a right ideal. If $P \in \mathcal{N}$, then $P \mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger} \subseteq \mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}$, so that $\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}$ is also a left ideal in \mathcal{N} . This gives (ii). The point (iii) is a consequence of (i) and (ii) by observing that the induced ring homomorphism

$$\mathcal{O}_{\mathfrak{x}, \mathbb{Q}} \xrightarrow{\subseteq} \mathcal{N} \longrightarrow \mathcal{N}/\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}$$

factors through the quotient morphism $\mathcal{O}_{\mathfrak{x}, \mathbb{Q}} \rightarrow \mathcal{O}_{\mathfrak{x}, \mathbb{Q}}/\mathcal{I}$. \square

COROLLARY 2.7. — *Let $i : \mathfrak{Y} \rightarrow \mathfrak{X}$ be a closed immersion between two smooth \mathfrak{o} -formal schemes, given by the ideal $\mathcal{I} \subseteq \mathcal{O}_{\mathfrak{x}}$. There is a canonical isomorphism of $\mathcal{O}_{\mathfrak{y}, \mathbb{Q}}$ -rings*

$$\mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^{\dagger} \simeq i^{-1}(\mathcal{N}(\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger})/\mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}).$$

Proof. — According to Proposition 2.4, one has

$$i_* \mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^\dagger \simeq \frac{\{P \in \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^\dagger \mid P(\mathcal{I}_{\mathbb{Q}}) \subseteq \mathcal{I}_{\mathbb{Q}}\}}{\mathcal{I}_{\mathbb{Q}} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^\dagger}$$

as $\mathcal{O}_{\mathfrak{x}, \mathbb{Q}}/\mathcal{I}$ -modules. We claim that

$$\mathcal{N} := \mathcal{N}(\mathcal{I}_{\mathbb{Q}} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^\dagger) = \{P \in \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^\dagger \mid P(\mathcal{I}_{\mathbb{Q}}) \subseteq \mathcal{I}_{\mathbb{Q}}\}.$$

If $P \in \mathcal{N}$ and $f \in \mathcal{I}_{\mathbb{Q}}$, then there is $Q \in \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^\dagger$ and $h \in \mathcal{I}_{\mathbb{Q}}$ such that $Pf = hQ$. It follows $P(f) = Pf(1) = hQ(1) \in h\mathcal{O}_{\mathfrak{x}, \mathbb{Q}} \subseteq \mathcal{I}_{\mathbb{Q}}$. This gives the forward inclusion. The inclusion being an equality may be checked locally. We may therefore assume that \mathfrak{X} is affine. Let $A = \mathcal{O}(\mathfrak{X})$, $I = \mathcal{O}(\mathcal{I})$ and $\bar{A} = A/I$. One has

$$N := \mathcal{O}(\mathcal{N}) = \{P \in \mathcal{D}_{A, \mathbb{Q}}^\dagger \mid PI_{\mathbb{Q}} \subseteq I_{\mathbb{Q}} \mathcal{D}_{A, \mathbb{Q}}^\dagger\}.$$

The above isomorphism says that the restriction map $P \mapsto \bar{P}$, where $\bar{P}(\bar{a}) := P(a) + I_{\mathbb{Q}}$ for $\bar{a} \in \bar{A}_{\mathbb{Q}}$ induces a surjective $A_{\mathbb{Q}}$ -linear morphism

$$\text{res} : \{P \in \mathcal{D}_{A, \mathbb{Q}}^\dagger \mid P(I_{\mathbb{Q}}) \subseteq I_{\mathbb{Q}}\} \longrightarrow \mathcal{D}_{\bar{A}, \mathbb{Q}}^\dagger$$

with kernel equal to $I_{\mathbb{Q}} \mathcal{D}_{A, \mathbb{Q}}^\dagger$. Let $P \in \mathcal{D}_{A, \mathbb{Q}}^\dagger$ with $P(I_{\mathbb{Q}}) \subseteq I_{\mathbb{Q}}$ and take $f \in I_{\mathbb{Q}}$. Given $h \in I_{\mathbb{Q}}$, we have $Pf(h) = P(fh) \subseteq \mathcal{I}_{\mathbb{Q}}$, since $fh \in I_{\mathbb{Q}}$, and hence $Pf(I_{\mathbb{Q}}) \subseteq I_{\mathbb{Q}}$. To calculate $\text{res}(Pf)$, we observe that

$$Pf(A_{\mathbb{Q}}) = P(fA_{\mathbb{Q}}) \subseteq P(I_{\mathbb{Q}}) \subseteq I_{\mathbb{Q}}.$$

Hence $\text{res}(Pf) = 0$, whence $Pf \in I_{\mathbb{Q}} \mathcal{D}_{A, \mathbb{Q}}^\dagger$. This shows $P \in N$ and establishes the isomorphism of $\mathcal{O}_{\mathfrak{x}, \mathbb{Q}}/\mathcal{I}_{\mathbb{Q}}$ -modules

$$i_* \mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^\dagger \simeq \frac{\mathcal{N}(\mathcal{I}_{\mathbb{Q}} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^\dagger)}{\mathcal{I}_{\mathbb{Q}} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^\dagger}.$$

However, since the map res is a ring homomorphism, this isomorphism is in fact an isomorphism of $\mathcal{O}_{\mathfrak{x}, \mathbb{Q}}/\mathcal{I}_{\mathbb{Q}}$ -rings. It remains to apply the functor i^{-1} . \square

2.4. β -bounded operators

In this subsection, \mathfrak{X} is a smooth affine \mathfrak{o} -formal scheme endowed with local coordinates x_1, \dots, x_M . Let $A := \mathcal{O}(\mathfrak{X})$.

The following basic result for local sections of $\mathcal{D}_{\mathfrak{x}}^\dagger$ is [2, 2.4.4]. Since its proof contains many arguments which we will refer to in the following, we will recall the proof here. This also allows us to fix some notations.

PROPOSITION 2.8. — *Let $\|\cdot\|$ be a Banach norm on $A_{\mathbb{Q}}$. For any operator*

$$P = \sum_{\underline{\nu}} a_{\underline{\nu}} \underline{\partial}^{[\underline{\nu}]} = \sum_{\underline{\nu}} a_{\underline{\nu}} \underline{\partial}^{\underline{\nu}} / \underline{\nu}! \in \Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}})$$

we denote by $\overline{P}_i \in \Gamma(X_i, \mathcal{D}_{X_i, k})$ the reduction of P modulo ϖ^{i+1} . The following three conditions are equivalent:

- (i) $P \in \Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}}^{\dagger})$
- (ii) *There are constants $\alpha, \beta \in \mathbb{R}, \alpha > 0$ such that for any $i \geq 0$*

$$\text{ord}(\overline{P}_i) \leq \alpha i + \beta$$
- (iii) *There are constants $c, \eta \in \mathbb{R}, \eta < 1$, such that for any $\underline{\nu}$*

$$\|a_{\underline{\nu}}\| \leq c\eta^{|\underline{\nu}|}.$$

Proof. —

(i) \Rightarrow (ii). — Since \mathfrak{X} is quasi-compact and quasi-separated, the functor $\Gamma(\mathfrak{X}, \cdot)$ commutes with direct limits and we have $\Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}}^{\dagger}) = \varinjlim_m \Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)})$. We may therefore choose m large enough and write P as

$$P = \sum_{\underline{\nu}} b_{\underline{\nu}} \underline{\partial}^{(\underline{\nu})} = \sum_{\underline{\nu}} \underline{q}! b_{\underline{\nu}} \underline{\partial}^{[\underline{\nu}]}$$

where $b_{\underline{\nu}} \in A$ tends to zero for $|\underline{\nu}| \rightarrow \infty$, and where, for any $\nu \in \mathbb{N}$, we have written $\nu = p^m q_{\nu} + r, 0 \leq r < p^m$. Let v be the normalized valuation of \mathfrak{o} . According to [2, Lem. 2.4.3], there exists $\alpha', \beta' \in \mathbb{R}, \alpha' > 0$ such that $v(q_{\nu}!) = ev_p(q_{\nu}!) \geq \alpha' \nu + \beta'$ for any $\nu \in \mathbb{N}$. Summing over all entries of $\underline{\nu}$, we therefore find $\alpha, \beta'' \in \mathbb{R}, \alpha > 0$ such that $v(\underline{q}!) \geq \alpha^{-1} |\underline{\nu}| + \beta''$ for any $\underline{\nu} \in \mathbb{N}^M$. Fix i . For any $\underline{\nu}$ with $|\underline{\nu}| \geq \alpha(i - \beta'' + 1)$, the inequality $|\underline{\nu}| \geq \alpha(i - \beta'' + 1)$ gives therefore

$$v(\underline{q}!) \geq i + 1.$$

The latter means $\underline{q}! b_{\underline{\nu}} \in \varpi^{i+1} A$ and so this coefficient of P reduces to zero in \overline{P}_i . It follows

$$\text{ord}(\overline{P}_i) \leq \alpha(i - \beta'' + 1) = \alpha i + \beta$$

where $\beta := \alpha(1 - \beta'')$.

(ii) \Rightarrow (iii). — To prove (iii), we may take for $\|\cdot\|$ the Banach norm of $A_{\mathbb{Q}}$, coming from the p -adic filtration on A . It satisfies $\|b\| \leq 1$ for any $b \in A$. Now suppose that $\text{ord}(\overline{P}_i) \leq \alpha i + \beta$ for all i . Fix $\underline{\nu}$. If $|\underline{\nu}| > \alpha i + \beta$, one has $a_{\underline{\nu}} \in \varpi^{i+1} A$, and hence $\|a_{\underline{\nu}} / \varpi^{i+1}\| \leq 1$. So for any $i \geq 0$ such that $|\underline{\nu}| > \alpha i + \beta$, i.e. $\alpha^{-1}(|\underline{\nu}| - \beta) > i$, one has $\|a_{\underline{\nu}}\| \leq |\varpi|^{i+1}$. Now take i such that

$$i + 1 \geq \alpha^{-1}(|\underline{\nu}| - \beta) > i.$$

Then, by what we have just shown,

$$\|a_{\underline{\nu}}\| \leq |\varpi|^{\alpha^{-1}(|\underline{\nu}|-\beta)}$$

for any $\underline{\nu}$. With $\eta := |\varpi|^{\alpha^{-1}}$ and $c := |\varpi|^{-\beta\alpha^{-1}}$ we therefore have $\|a_{\underline{\nu}}\| \leq c\eta^{|\underline{\nu}|}$ for any $\underline{\nu}$ where $\eta < 1$.

(iii) \Rightarrow (i). — Suppose that $\|a_{\underline{\nu}}\| \leq c\eta^{|\underline{\nu}|}$ with $\eta < 1$. Here, we may again assume that $\|\cdot\|$ is the Banach norm on $A_{\mathbb{Q}}$ coming from the p -adic filtration on A . We first show that, for $m \geq 0$ sufficiently big, the elements $b_{\underline{\nu}} := a_{\underline{\nu}}/\underline{q}!$ tend to zero for $|\underline{\nu}| \rightarrow \infty$ in the Banach algebra $A_{\mathbb{Q}}$. By hypothesis we may write $\eta = p^{-\frac{\alpha}{e}}$ with $\alpha > 0$ and obtain

$$v_p(a_{\underline{\nu}}) := -\log_p \|a_{\underline{\nu}}\| \geq |\underline{\nu}|(-\log_p \eta) - \log_p c = |\underline{\nu}| \left(\frac{\alpha}{e} \right) + \mu$$

où $\mu = -\log_p c$. An upper bound for the p -adic valuation of $\underline{q}!$ is given by $|\underline{\nu}|/p^m(p-1)$ [2, Lem. 2.4.3]. Hence,

$$v_p(b_{\underline{\nu}}) = v_p(a_{\underline{\nu}}) - v_p(\underline{q}!) \geq |\underline{\nu}| \left(\frac{\alpha}{e} - \frac{1}{p^m(p-1)} \right) + \mu.$$

Thus, for sufficiently big m , we obtain indeed $v_p(b_{\underline{\nu}}) \rightarrow \infty$ for $|\underline{\nu}| \rightarrow \infty$. For the remaining statement, let $\lambda := \alpha/e$. If

$$|\underline{\nu}| \geq -\mu(\lambda - 1/p^m(p-1))^{-1},$$

then $v_p(b_{\underline{\nu}}) \geq 0$ and hence $\|b_{\underline{\nu}}\| \leq 1$. This implies $b_{\underline{\nu}} \in A$. Suppose therefore that $\mu < 0$. Adjusting m we may suppose that $p^m > \lambda^{-1}(-\mu + 1/p - 1)$. Then

$$p^m > -\mu(\lambda - 1/p^m(p-1))^{-1}.$$

So for all

$$|\underline{\nu}| < -\mu(\lambda - 1/p^m(p-1))^{-1}$$

we obtain $\underline{q} = 0$, which implies $b_{\underline{\nu}} = a_{\underline{\nu}} \in A$. □

Let $n \geq 1$ and let $P \in M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{O}}_{\mathfrak{X}}))$ be a given $n \times n$ -matrix, with entries in the ring $\Gamma(\mathfrak{X}, \widehat{\mathcal{O}}_{\mathfrak{X}})$. It will be convenient to write $P = \sum_{\underline{\nu}} a_{\underline{\nu}} \underline{\partial}^{[\underline{\nu}]}$, with coefficients $a_{\underline{\nu}} \in M_n(\Gamma(\mathfrak{X}, \mathcal{O}_{\mathfrak{X}}))$. In particular, we may speak of the order $\text{ord}(P)$ of P , whenever the sum is finite. In general, we write for any integer $\ell \geq 0$

$$[P]_{\ell} := \varpi^{-\ell} \sum_{\underline{\nu} \in E_{\ell}} a_{\underline{\nu}} \underline{\partial}^{[\underline{\nu}]} \quad \text{and} \quad \sigma_{\ell}(P) := \sum_{\ell' \leq \ell} \varpi^{\ell'} [P]_{\ell'}$$

where $E_{\ell} := \{\underline{\nu} \in \mathbb{N}^M, v_p(a_{\underline{\nu}}) = \ell\}$ is a finite set. Here, $v_p(\cdot) := -\log_p \|\cdot\|$ where $\|\cdot\|$ denotes the Banach norm on $M_n(A_{\mathbb{Q}})$ coming from the p -adic filtration on $M_n(A)$. By definition one has

$$\text{ord}(\sigma_{\ell}(P)) = \text{ord}(\overline{P}_{\ell})$$

where \overline{P}_ℓ is the reduction of P modulo $\varpi^{\ell+1}$. This implies

$$\text{ord}(\sigma_\ell(P + Q)) \leq \max(\text{ord}(\sigma_\ell(P)), \text{ord}(\sigma_\ell(Q)))$$

for two operators P, Q .

LEMMA 2.9. — *Let $\alpha > 0$. One has the equivalence*

$$\text{ord}(\sigma_\ell(P)) \leq \alpha\ell + \beta \quad \text{for all } \ell \iff \text{ord}([P]_\ell) \leq \alpha\ell + \beta \quad \text{for all } \ell.$$

Proof. — If $\ell' \neq \ell$, then E_ℓ is disjoint from $E_{\ell'}$. This implies the general identity

$$\text{ord}(\sigma_\ell(P)) = \max_{\ell' \leq \ell} \text{ord}(\varpi^{\ell'} [P]_{\ell'}).$$

In particular, the implication \Rightarrow is clear. On the other hand, if $[P]_{\ell'} \leq \alpha\ell' + \beta$ for any ℓ' , then the right hand side in the above identity is bounded by $\alpha\ell + \beta$, since $\alpha > 0$. \square

DEFINITION 2.10. — *Let $P \in M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_\mathfrak{X}))$ and $\beta > 0$. The operator P is called β -bounded, if for all $\ell \geq 0$*

$$\text{ord}(\sigma_\ell(P)) \leq \beta(\ell + 1).$$

LEMMA 2.11. — *Let $\beta > 0$ and let P_ℓ be a ϖ -adically convergent sequence of β -bounded elements in $M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_\mathfrak{X}))$. Then $P = \lim_\ell P_\ell$ is β -bounded.*

Proof. — Fix $n \geq 0$. By the inequality before Lemma 2.9, we have $\text{ord}(\sigma_n(P)) \leq \max(\text{ord}(\sigma_n(P - P_\ell)), \text{ord}(\sigma_n(P_\ell)))$. Choose ℓ sufficiently large such that $P - P_\ell$ is divisible by ϖ^{n+1} . Then $\sigma_n(P - P_\ell) = 0$ and the claim follows. \square

The interest in this notion comes from the following result.

PROPOSITION 2.12. — *Any $P \in M_n(\Gamma(\mathfrak{X}, \mathcal{D}_\mathfrak{X}^\dagger))$ is β -bounded for some $\beta > 0$. Conversely, any β -bounded $P \in M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_\mathfrak{X}))$ belongs to $M_n(\Gamma(\mathfrak{X}, \mathcal{D}_\mathfrak{X}^\dagger))$.*

Proof. — According to the above discussion, P is β -bounded iff $\text{ord}([P]_\ell) \leq \beta(\ell + 1)$ for all ℓ or, equivalently, if $\text{ord}(\overline{P}_\ell) \leq \beta(\ell + 1)$ for all ℓ . Thus, Proposition 2.8 implies the claim. \square

2.5. A key lemma

In this subsection, \mathfrak{X} is a smooth affine \mathfrak{o} -formal scheme endowed with local coordinates x_1, \dots, x_M . Let $A := \mathcal{O}(\mathfrak{X})$. The main steps in the proof of the key lemma presented here are extracted from Caro's discussion in [7].

Let $\mathbb{1} \in M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}))$ be the identity in the matrix ring $M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}))$. We consider $\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}})$ to be a subring of $M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}))$ via the injective ring homomorphism

$$\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}) \longrightarrow M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}})), P \longmapsto P\mathbb{1}.$$

Let $m \geq 1$ and let $R \in M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)}))$ be a given $n \times n$ -matrix.

By Proposition 2.12, R is α -bounded for some $\alpha > 0$, so that $\text{ord}(\sigma_\ell(R)) \leq \alpha(\ell + 1)$ for all ℓ . In the following we fix $\beta > 0$ sufficiently large, such that

$$2\alpha + p^m \leq \beta. \quad (\text{HYP})$$

We start with two auxiliary lemmas. Let $t := x_M$.

LEMMA 2.13. — *For any $U \in M_n(\Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}}))$ there is an operator $Q \in M_n(\Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}}))$ such that*

$$[t^{p^m}, Q] \equiv U \pmod{\varpi} \quad \text{and} \quad \text{ord}(Q) \leq \text{ord}(U) + p^m.$$

Proof. — Soit $\partial_t := \partial_{x_M}$. We first establish the general identity

$$t^{p^m} \partial_t^{[N+p^m]} - \partial_t^{[N+p^m]} t^{p^m} = -\partial_t^{[N]} \pmod{\varpi}$$

for any integer N . Indeed, one has

$$\partial_t^{[N+p^m]} t^{p^m} = \sum_{\nu+\nu'=N+p^m} \partial_t^{[\nu]} (t^{p^m}) \partial_t^{[\nu']}.$$

Since $\partial_t^{[\nu]}(t^{p^m}) = 0$ for $\nu > p^m$ and $\equiv 0 \pmod{\varpi}$ for $0 < \nu < p^m$, only the terms indexed by $(0, N + p^m)$ and (p^m, N) survive in the sum. This yields the claimed identity. Now write $U = \sum_{\underline{\nu}} a_{\underline{\nu}} \partial^{[\underline{\nu}]}$ and take $Q = -\sum_{\underline{\nu}} a_{\underline{\nu}} \partial^{[\underline{\nu}+(0, \dots, 0, p^m)]}$. Then $\text{ord}(Q) \leq \text{ord}(U) + p^m$ and

$$\begin{aligned} [t^{p^m}, Q] &= \sum_{\underline{\nu}} (-a_{\underline{\nu}}) \partial^{[(\nu_1, \dots, \nu_{M-1})]} (t^{p^m} \partial_t^{[\nu_M+p^m]} - \partial_t^{[\nu_M+p^m]} t^{p^m}) \\ &\equiv \sum_{\underline{\nu}} (-a_{\underline{\nu}}) \partial^{[(\nu_1, \dots, \nu_{M-1})]} (-\partial_t^{[\nu_M]}) \pmod{\varpi} \\ &\equiv U. \end{aligned} \quad \square$$

LEMMA 2.14. — *Let $Q_\ell \in M_n(\Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}}))$ such that $\text{ord}(Q_\ell) \leq \beta(\ell + 2)$. Then $\varpi^{\ell+1} Q_\ell$ is β -bounded.*

Proof. — Let $P = \varpi^{\ell+1} Q_\ell$. If $i \leq \ell$, one has $\bar{P}_i = 0$ and therefore $\text{ord}(\bar{P}_i) \leq \beta(i + 1)$ trivially. If $i > \ell$, then

$$\text{ord}(\bar{P}_i) \leq \text{ord}(P) \leq \beta(\ell + 2) \leq \beta(i + 1). \quad \square$$

We now construct by induction a sequence $P_\ell \in M_n(\Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}}))$ (depending on the matrix R) such that

- (i) $P_0 = 1, \sigma_\ell(P_\ell) = P_\ell$ and $P_{\ell+1} \equiv P_\ell \pmod{\varpi^{\ell+1}}$
- (ii) P_ℓ is β -bounded
- (iii) $t^{p^m} P_\ell \equiv P_\ell(t^{p^m} - \varpi\sigma_{\ell-1}(R)) \pmod{\varpi^{\ell+1}}$.

Suppose that P_0, \dots, P_ℓ are already constructed. We will construct $P_{\ell+1}$ in the following.

Since $\varpi\sigma_\ell \equiv \varpi\sigma_{\ell-1} \pmod{\varpi^{\ell+1}}$, the property (iii) implies

$$\begin{aligned} [t^{p^m}, P_\ell] + \varpi P_\ell \sigma_\ell(R) &= t^{p^m} P_\ell - P_\ell(t^{p^m} + \varpi\sigma_\ell(R)) \\ &\equiv t^{p^m} P_\ell - P_\ell(t^{p^m} + \varpi\sigma_{\ell-1}(R)) \pmod{\varpi^{\ell+1}} \\ &\equiv 0 \pmod{\varpi^{\ell+1}}. \end{aligned}$$

On the other hand, since $\sigma_\ell(P_\ell) = P_\ell$ by (i), one has $P_\ell = \sum_{\ell_1 \leq \ell} \varpi^{\ell_1} [P_\ell]_{\ell_1}$ and hence

$$\begin{aligned} P_\ell \sigma_\ell(R) &= \sum_{\ell_1, \ell_2 \leq \ell} \varpi^{\ell_1 + \ell_2} [P_\ell]_{\ell_1} [R]_{\ell_2} \\ &\equiv \sum_{\ell_1, \ell_2 \leq \ell, \ell_1 + \ell_2 \leq \ell + 1} \varpi^{\ell_1 + \ell_2} [P_\ell]_{\ell_1} [R]_{\ell_2} \pmod{\varpi^{\ell+2}}. \end{aligned}$$

So altogether one obtains

$$[t^{p^m}, P_\ell] + \varpi \sum_{\ell_1, \ell_2 \leq \ell, \ell_1 + \ell_2 \leq \ell + 1} \varpi^{\ell_1 + \ell_2} [P_\ell]_{\ell_1} [R]_{\ell_2} \equiv 0 \pmod{\varpi^{\ell+1}}.$$

So there is an operator $U_\ell \in M_n(\Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}}))$ such that

$$-\varpi^{\ell+1} U_\ell = [t^{p^m}, P_\ell] + \varpi \sum_{\ell_1, \ell_2 \leq \ell, \ell_1 + \ell_2 \leq \ell + 1} \varpi^{\ell_1 + \ell_2} [P_\ell]_{\ell_1} [R]_{\ell_2}.$$

By the above discussion

$$-\varpi^{\ell+1} U_\ell \equiv [t^{p^m}, P_\ell] + \varpi P_\ell \sigma_\ell(R) \pmod{\varpi^{\ell+2}}.$$

ASSERTION 1. — *One has $\text{ord}(U_\ell) \leq \beta(\ell + 1) + 2\alpha$.*

To prove the assertion, we use that P_ℓ is β -bounded by (ii). In particular, $\text{ord}(P_\ell) = \text{ord}(\sigma_\ell(P_\ell)) \leq \beta(\ell + 1)$ which gives $\text{ord}([t^{p^m}, P_\ell]) \leq \beta(\ell + 1)$.

Again, by (ii), we have $\text{ord}([P_\ell]_{\ell_1}) \leq \beta(\ell_1 + 1)$ for all ℓ_1 . This gives

$$\begin{aligned} \text{ord}([P_\ell]_{\ell_1} [R]_{\ell_2}) &\leq \text{ord}([P_\ell]_{\ell_1}) + \text{ord}([R]_{\ell_2}) \\ &\leq \beta(\ell_1 + 1) + \alpha(\ell_2 + 1) \leq \beta(\ell + 1) + 2\alpha. \end{aligned}$$

Note that the last inequality follows from $\alpha(\ell_2 - 1) \leq \beta(\ell - \ell_1)$ which in turn follows directly from $\ell_1 + \ell_2 \leq \ell + 1$ and $\alpha \leq \beta$. This implies the assertion.

We now use Lemma 2.13 to find an operator $Q_\ell \in M_n(\Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}}))$ such that

$$[t^{p^m}, Q_\ell] \equiv U_\ell \pmod{\varpi} \quad \text{and} \quad \text{ord}(Q_\ell) \leq \text{ord}(U_\ell) + p^m. \quad (*)$$

We now set

$$P_{\ell+1} := \sigma_{\ell+1}(P_\ell + \varpi^{\ell+1}Q_\ell) \in M_n(\Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}})).$$

ASSERTION 2. — *The operator $P_{\ell+1}$ satisfies (i), (ii), (iii) above.*

We start with (iii). Modulo $\varpi^{\ell+2}$ one certainly has the two congruences

$$t^{p^m} P_{\ell+1} \equiv t^{p^m} (P_\ell + \varpi^{\ell+1}Q_\ell)$$

and

$$P_{\ell+1}(t^{p^m} - \varpi\sigma_\ell(R)) \equiv (P_\ell + \varpi^{\ell+1}Q_\ell)(t^{p^m} - \varpi\sigma_\ell(R))$$

so it suffices to show that the two right-hand sides are congruent. But modulo $\varpi^{\ell+2}$, one has

$$\begin{aligned} t^{p^m} (P_\ell + \varpi^{\ell+1}Q_\ell) &= t^{p^m} P_\ell + \varpi^{\ell+1}t^{p^m} Q_\ell \\ &\equiv t^{p^m} P_\ell + \varpi^{\ell+1}(U_\ell + Q_\ell t^{p^m}) \\ &\equiv t^{p^m} P_\ell + \varpi^{\ell+1}Q_\ell t^{p^m} - ([t^{p^m}, P_\ell] + \varpi P_\ell \sigma_\ell(R)) \\ &= (P_\ell + \varpi^{\ell+1}Q_\ell)t^{p^m} - \varpi P_\ell \sigma_\ell(R) \\ &\equiv (P_\ell + \varpi^{\ell+1}Q_\ell)(t^{p^m} - \varpi\sigma_\ell(R)) \end{aligned}$$

where the first congruence is the property (*) and the middle congruence is the congruence before Assertion 1.

To see (ii), we just note that our hypothesis (HYP) implies $\text{ord}(Q_\ell) \leq \beta(\ell + 2)$ by Assertion 1. Hence, $\varpi^{\ell+1}Q_\ell$ is β -bounded by Lemma 2.14. Let $\ell' \leq \ell + 1$. Since $\sigma_{\ell'} \circ \sigma_{\ell+1} = \sigma_{\ell'}$ we find

$$\begin{aligned} \text{ord}(\sigma_{\ell'}(P_{\ell+1})) &= \text{ord}(\sigma_{\ell'}(P_\ell + \varpi^{\ell+1}Q_\ell)) \\ &\leq \max(\text{ord}(\sigma_{\ell'}(P_\ell)), \text{ord}(\sigma_{\ell'}(\varpi^{\ell+1}Q_\ell))) \leq \beta(\ell' + 1) \end{aligned}$$

where we have used the inequality before Lemma 2.9 and the fact that P_ℓ and $\varpi^{\ell+1}Q_\ell$ are β -bounded. Let $\ell' \geq \ell + 1$. Since $\sigma_{\ell'} \circ \sigma_{\ell+1} = \sigma_{\ell+1}$ we find

$$\begin{aligned} \text{ord}(\sigma_{\ell'}(P_{\ell+1})) &= \text{ord}(P_{\ell+1}) \leq \max(\text{ord}(\sigma_{\ell+1}(P_\ell)), \text{ord}(\sigma_{\ell+1}(\varpi^{\ell+1}Q_\ell))) \\ &\leq \beta(\ell + 1) \leq \beta(\ell' + 1). \end{aligned}$$

Hence, $P_{\ell+1}$ is β -bounded.

It remains to see (i). The identity $P_{\ell+1} = \sigma_{\ell+1}(P_{\ell+1})$ is trivial. In particular, we may write

$$P_{\ell+1} = \sum_{v_p(a_{\underline{\nu}}^{\ell+1}) \leq \ell+1} a_{\underline{\nu}}^{\ell+1} \underline{\partial}^{[\underline{\nu}]}$$

where $a_{\underline{\nu}}^{\ell+1} \in M_n(\Gamma(\mathfrak{X}, \mathcal{O}_{\mathfrak{X}}))$ are the coefficients of $P_{\ell+1}$. By definition, one has

$$a_{\underline{\nu}}^{\ell+1} = a_{\underline{\nu}}^{\ell} + \varpi^{\ell+1} \tilde{a}_{\underline{\nu}}$$

where $\tilde{a}_{\underline{\nu}}$ are the coefficients of Q_{ℓ} . Because of $\sigma_{\ell}(P_{\ell}) = P_{\ell}$, one has $a_{\underline{\nu}}^{\ell} = 0$ for all coefficients $a_{\underline{\nu}}^{\ell}$ with $v_p(a_{\underline{\nu}}^{\ell}) \geq \ell + 1$. In turn, the inequality $v_p(a_{\underline{\nu}}^{\ell}) \leq \ell$ implies $v_p(a_{\underline{\nu}}^{\ell+1}) = v_p(a_{\underline{\nu}}^{\ell}) \leq \ell$ by the ultrametric inequality for v_p .

This means

$$P_{\ell+1} \pmod{\varpi^{\ell+1}} \equiv \sum_{v_p(a_{\underline{\nu}}^{\ell+1}) \leq \ell} a_{\underline{\nu}}^{\ell} \underline{\partial}^{[\underline{\nu}]} = \sum_{v_p(a_{\underline{\nu}}^{\ell}) \leq \ell} a_{\underline{\nu}}^{\ell} \underline{\partial}^{[\underline{\nu}]} = \sigma_{\ell}(P_{\ell}) = P_{\ell}$$

which completes (i).

So there is indeed a sequence $(P_{\ell})_{\ell}$ with the properties (i)–(iii) as claimed. Choose $m' \geq m$ such that $P_{\ell} \in M_n(\Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}}^{(m')}))$ for all ℓ by Proposition 2.12. We may consider its limit

$$P = \lim_{\ell} P_{\ell} \in M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m')})).$$

Then we have inside $M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m')}))$

- (1) $P \equiv 1 \pmod{\varpi}$
- (2) $t^{p^m} P = P(t^{p^m} - \varpi R)$.

Thus we have proved the following lemma.

LEMMA 2.15 (Berthelot's key lemma). — *Let $t := x_M$ and let $R \in M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)}))$ be a given $n \times n$ -matrix. There exist $m' \geq m$ and $P \in M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m')}))$, such that*

- (1) $P \equiv 1 \pmod{\varpi}$
- (2) $t^{p^m} P = P(t^{p^m} - \varpi R)$ in $M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m')}))$.

Note that, as a consequence of (1), the matrix P appearing in the lemma is invertible in the ϖ -adically complete ring $M_n(\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m')}))$.

COROLLARY 2.16. — *Let M be a finitely generated $\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)})$ -module with generators e_1, \dots, e_n . Suppose that $t^{p^m} e_i \equiv 0 \pmod{\varpi}$ for all i . Then*

there is $m' \geq m$ and a set of generators e'_1, \dots, e'_n for the $\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m')})$ -module $\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m')}) \otimes_{\Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)})} M$ with the property $t^{p^m} e'_i = 0$ for all i .

Proof. — Write $\widehat{D}_{\mathfrak{X}}^{(m)} := \Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)})$ and \underline{e} be the column vector $T(e_1, \dots, e_n)$. By assumption, there is $R \in M_n(\widehat{D}_{\mathfrak{X}}^{(m)})$ such that

$$t^{p^m} \underline{e} = \varpi R \underline{e}.$$

By the key lemma, there are $m' \geq m$ and $P \in M_n(\widehat{D}_{\mathfrak{X}}^{(m')})$, such that

- (1) $P \equiv 1 \pmod{\varpi}$
- (2) $t^{p^m} P = P(t^{p^m} - \varpi R)$ in $M_n(\widehat{D}_{\mathfrak{X}}^{(m')})$.

Let

$$M' := \widehat{D}_{\mathfrak{X}}^{(m')} \otimes_{\widehat{D}_{\mathfrak{X}}^{(m)}} M \quad \text{and} \quad \underline{e}' := P(1 \otimes \underline{e}).$$

Here, $1 \otimes \underline{e}$ is the column vector $T(1 \otimes e_1, \dots, 1 \otimes e_n) \in (M')^n$, so that

$$e'_i = \sum_j P_{ij} \otimes e_j \in M' \quad \text{for } i = 1, \dots, n.$$

The e'_i are generators for the left $\widehat{D}_{\mathfrak{X}}^{(m')}$ -module M' . Indeed, given $y \in M'$ with $y = \sum \lambda_i \otimes e_i$, then $y = \sum \lambda'_i \otimes e'_i$ with $(\lambda'_1, \dots, \lambda'_n) := (\lambda_1, \dots, \lambda_n) \cdot P^{-1}$. Moreover, (2) implies

$$\begin{aligned} t^{p^m} \underline{e}' &= t^{p^m} P(1 \otimes \underline{e}) = P(t^{p^m} - \varpi R)(1 \otimes \underline{e}) \\ &= P(1 \otimes (t^{p^m} \underline{e} - \varpi R \underline{e})) = 0. \end{aligned} \quad \square$$

3. The Berthelot–Kashiwara theorem

Let

$$i : \mathfrak{Y} \longrightarrow \mathfrak{X}$$

be a closed immersion between smooth formal \mathfrak{o} -schemes given by an ideal sheaf $\mathcal{I} \subseteq \mathcal{O}_{\mathfrak{X}}$.

3.1. Direct image and adjointness

Let $\mathcal{D}_{\mathfrak{X} \leftarrow \mathfrak{Y}, \mathbb{Q}}^\dagger$ be the associated transfer module, a $(i^{-1} \mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger, \mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger)$ -bimodule, cf. [3, 3.4]. Let \mathcal{N} be a left $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -module. Its direct image along i is the left $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ -module

$$i_+ \mathcal{N} := i_* (\mathcal{D}_{\mathfrak{X} \leftarrow \mathfrak{Y}, \mathbb{Q}}^\dagger \otimes_{\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger} \mathcal{N}).$$

This yields a functor i_+ from left $\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger$ -modules to left $\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger$ -modules, cf. [4, 4.3.7], which preserves coherence [4, 4.3.8]. If $\mathfrak{Z} \xrightarrow{k} \mathfrak{Y}$ is a second closed immersion of smooth formal \mathfrak{o} -schemes and if we restrict to *coherent* modules, then there is a natural isomorphism $(i \circ k)_+ \simeq i_+ \circ k_+$ of functors [4, 3.5.2].

PROPOSITION 3.1. —

- (i) *The right $\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger$ -module $\mathcal{D}_{\mathfrak{X} \leftarrow \mathfrak{Y},\mathbb{Q}}^\dagger$ is flat.*
- (ii) *The functor i_+ is exact.*
- (iii) *If $S \subseteq \mathfrak{Y}$ is a closed subset and \mathcal{N} is a left $\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger$ -module supported on S , then $i_+ \mathcal{N}$ is supported on S .*

Proof. — (i) maybe proved by adapting the proof in the classical setting [10, 1.3.5] as follows. Fix a level $m \geq 0$. By definition [3, 3.4.1],

$$\mathcal{D}_{\mathfrak{Y} \leftarrow \mathfrak{X}}^{(m)} = i^{-1}(\mathcal{D}_{\mathfrak{X}}^{(m)} \otimes_{\mathcal{O}_{\mathfrak{X}}} \omega_{\mathfrak{X}}^{-1}) \otimes_{i^{-1}\mathcal{O}_{\mathfrak{X}}} \omega_{\mathfrak{Y}}$$

where $\omega_{\mathfrak{X}}$ and $\omega_{\mathfrak{Y}}$ are the modules of differentials of highest order on \mathfrak{X} and \mathfrak{Y} respectively. Since (i) is a local question, we may from now on assume that \mathfrak{X} is affine equipped with local coordinates $t_1, \dots, t_d \in \mathcal{O}_{\mathfrak{X}}$, that $\mathcal{I} = (t_{r+1}, \dots, t_d)$ for some $0 \leq r < d$ and that $\tilde{t}_1, \dots, \tilde{t}_r$ are local coordinates for \mathfrak{Y} . Let ∂_i be the derivation relative to t_i . We identify $i^{-1}\omega_{\mathfrak{X}}^{-1} \otimes_{i^{-1}\mathcal{O}_{\mathfrak{X}}} \omega_{\mathfrak{Y}}$ with $\mathcal{O}_{\mathfrak{Y}}$ via the section

$$(dt_1 \wedge \dots \wedge dt_d)^{\otimes -1} \otimes (dt_1 \wedge \dots \wedge dt_r).$$

Note that $D' := \bigoplus_{\nu} \partial_1^{(\nu_1)} \dots \partial_r^{(\nu_r)} \mathcal{O}_{\mathfrak{X}} \subset \mathcal{D}_{\mathfrak{X}}^{(m)}$ is a subring of $\mathcal{D}_{\mathfrak{X}}^{(m)}$. It is clear that $\mathcal{D}_{\mathfrak{X}}^{(m)} \simeq \mathfrak{o}[\partial]^{(m)} \otimes_{\mathfrak{o}} D'$ as a right D' -module, where $\mathfrak{o}[\partial]^{(m)}$ equals the free \mathfrak{o} -module on the basis $\partial_{r+1}^{(\nu_{r+1})} \dots \partial_d^{(\nu_d)}$. It follows that

$$\mathcal{D}_{\mathfrak{Y} \leftarrow \mathfrak{X}}^{(m)} \simeq \mathfrak{o}[\partial]^{(m)} \otimes_{\mathfrak{o}} (i^{-1}D' \otimes_{i^{-1}\mathcal{O}_{\mathfrak{X}}} \mathcal{O}_{\mathfrak{Y}}).$$

It is easy to see that $i^{-1}D' \otimes_{i^{-1}\mathcal{O}_{\mathfrak{X}}} \mathcal{O}_{\mathfrak{Y}} \simeq \mathcal{D}_{\mathfrak{Y}}^{(m)}$ as a right $\mathcal{D}_{\mathfrak{Y}}^{(m)}$ -module. This means

$$\mathcal{D}_{\mathfrak{X} \leftarrow \mathfrak{Y}}^{(m)} \simeq \mathfrak{o}[\partial]^{(m)} \otimes_{\mathfrak{o}} \mathcal{D}_{\mathfrak{Y}}^{(m)}$$

as right $\mathcal{D}_{\mathfrak{Y}}^{(m)}$ -modules. In particular, $\mathcal{D}_{\mathfrak{X} \leftarrow \mathfrak{Y}}^{(m)}$ is free, and hence flat, as a right $\mathcal{D}_{\mathfrak{Y}}^{(m)}$ -module. According to [11, 2.2.2] the sheaf of rings $\mathcal{D}_{\mathfrak{Y}}^{(m)}$ is locally noetherian, hence [2, 3.2.4] implies that $\widehat{\mathcal{D}}_{\mathfrak{X} \leftarrow \mathfrak{Y}}^{(m)}$ is flat as a right $\widehat{\mathcal{D}}_{\mathfrak{Y}}^{(m)}$ -module. Passing to the limit and inverting p , we finally see that $\mathcal{D}_{\mathfrak{X} \leftarrow \mathfrak{Y}}^\dagger$ is flat as a right $\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger$ -module, as claimed. The point (ii) follows from (i). Finally (iii) follows from the fact that, for any abelian sheaf \mathcal{K} on \mathfrak{Y} , the stalk of $i_* \mathcal{K}$ at $x \in \mathfrak{X}$ equals \mathcal{K}_x if $x \in \mathfrak{Y}$ and is zero otherwise. Hence, if $S \subset \mathfrak{Y}$ is closed and \mathcal{N} is supported on S and $x \in \mathfrak{Y} \setminus S$, then

$$(i_+ \mathcal{N})_x = (\mathcal{D}_{\mathfrak{X} \leftarrow \mathfrak{Y},\mathbb{Q}}^\dagger \otimes_{\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger} \mathcal{N})_x = \mathcal{D}_{\mathfrak{X} \leftarrow \mathfrak{Y},\mathbb{Q},x}^\dagger \otimes_{\mathcal{D}_{\mathfrak{Y},\mathbb{Q},x}^\dagger} \mathcal{N}_x = 0. \quad \square$$

We define the following functor from left $\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger$ -modules to left $\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger$ -modules:

$$i^\natural \mathcal{M} := \mathcal{H}\mathrm{om}_{i^{-1}\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger}(\mathcal{D}_{\mathfrak{X}\leftarrow\mathfrak{Y},\mathbb{Q}}^\dagger, i^{-1}\mathcal{M}).$$

PROPOSITION 3.2. —

- (i) *The functor i^\natural is right adjoint to i_+ . In particular, i^\natural is left-exact.*
- (ii) *If $\mathfrak{Z} \xrightarrow{k} \mathfrak{Y}$ is a second closed immersion of smooth formal \mathfrak{o} -schemes, and if k^\natural, i^\natural preserve coherence, then there is a natural isomorphism $(i \circ k)^\natural \simeq k^\natural \circ i^\natural$.*
- (iii) *If $S \subseteq \mathfrak{X}$ is a closed subset and \mathcal{M} is a coherent left $\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger$ -module supported on S , then $i^\natural \mathcal{M}$ is supported on $S \cap \mathfrak{Y}$.*

Proof. — Since i is a closed immersion, i_* has the right adjoint i^{-1} . Therefore, for any coherent left $\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger$ -module \mathcal{N} and any coherent left $\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger$ -module \mathcal{M} , one has

$$\mathcal{H}\mathrm{om}_{\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger}(i_+\mathcal{N}, \mathcal{M}) = \mathcal{H}\mathrm{om}_{i^{-1}\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger}(\mathcal{D}_{\mathfrak{X}\leftarrow\mathfrak{Y},\mathbb{Q}}^\dagger \otimes_{\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger} \mathcal{N}, i^{-1}\mathcal{M}).$$

One obtains (i) by combining this with the standard tensor-hom adjunction

$$\begin{aligned} \mathcal{H}\mathrm{om}_{i^{-1}\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger}(\mathcal{D}_{\mathfrak{X}\leftarrow\mathfrak{Y},\mathbb{Q}}^\dagger \otimes_{\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger} \mathcal{N}, i^{-1}\mathcal{M}) \\ = \mathcal{H}\mathrm{om}_{\mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger}(\mathcal{N}, \mathcal{H}\mathrm{om}_{i^{-1}\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger}(\mathcal{D}_{\mathfrak{X}\leftarrow\mathfrak{Y},\mathbb{Q}}^\dagger, i^{-1}\mathcal{M})). \end{aligned}$$

(ii) follows from uniqueness of adjoint functors and the fact that for coherent modules we have $(i \circ k)_+ \simeq i_+ \circ k_+$. For (iii), for any abelian sheaf \mathcal{K} on \mathfrak{X} , the stalk of $i^{-1}\mathcal{K}$ at $x \in \mathfrak{Y}$ equals \mathcal{K}_x . Hence, if $S \subset \mathfrak{X}$ is a closed subset and \mathcal{M} is supported on S and $x \in \mathfrak{Y} \setminus S$, then $(i^\natural \mathcal{M})_x = 0$. Indeed, this is a local statement, and we may suppose that the coherent module \mathcal{M} has a global finite presentation. This means that $i^{-1}\mathcal{M}$ can be written as $\mathrm{coker}(f)$ for some $i^{-1}\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger$ -linear morphism $(i^{-1}\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger)^{\oplus s} \xrightarrow{f} (i^{-1}\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger)^{\oplus t}$. Take an open $\mathfrak{U} \subseteq \mathfrak{Y}$ containing x with $\mathfrak{U} \cap S = \emptyset$. For any $y \in \mathfrak{U}$, one has $(i^{-1}\mathcal{M})_y = \mathcal{M}_y = 0$. Hence f_y is surjective for any $y \in \mathfrak{U}$ and so $f|_{\mathfrak{U}}$ is surjective, i.e. $i^{-1}\mathcal{M}|_{\mathfrak{U}} = 0$. Taking the limit over all open neighbourhoods $\mathfrak{V} \subset \mathfrak{U}$ of x , one finds

$$(i^\natural \mathcal{M})_x = \varinjlim_{x \in \mathfrak{V} \subset \mathfrak{U}} \mathcal{H}\mathrm{om}_{i^{-1}\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger|_{\mathfrak{V}}}(\mathcal{D}_{\mathfrak{X}\leftarrow\mathfrak{Y},\mathbb{Q}}^\dagger|_{\mathfrak{V}}, i^{-1}\mathcal{M}|_{\mathfrak{V}}) = 0. \quad \square$$

3.2. Berthelot–Kashiwara for left modules

We start with an auxiliary lemma.

LEMMA 3.3. — *Let N be a K -vector space and $s \geq 1$. Let $N[[\partial^{[\underline{\nu}}]]$ be the K -subspace of all formal infinite sums $\sum_{\underline{\nu} \in \mathbb{N}^s} n_{\underline{\nu}} \partial^{[\underline{\nu}]}$ with $n_{\underline{\nu}} \in N$ on formal symbols $\partial^{[\underline{\nu}]}$. We regard N as a K -subspace of $N[[\partial^{[\underline{\nu}}]]$ via $n \mapsto n \partial^{[0]}$. Define for $i = 1, \dots, s$ a linear operator t_i on $N[[\partial^{[\underline{\nu}}]]$ through $t_i \cdot (n_{\underline{\nu}} \partial^{[\underline{\nu}]}) := n_{\underline{\nu}} (\partial^{[\underline{\nu} - e_i]})$ when $\nu_i > 0$ and zero else. Then*

$$\bigcap_{i=1, \dots, s} \ker(N[[\partial^{[\underline{\nu}}]] \xrightarrow{t_i} N[[\partial^{[\underline{\nu}}]]) = N.$$

Proof. — It suffices to check the forward inclusion, the reverse inclusion being clear. By induction on s it suffices to treat the case $s = 1$. Writing $\sum_{\nu \in \mathbb{N}} m_{\nu} \partial^{[\nu]} := t \cdot \sum_{\nu \in \mathbb{N}} n_{\nu} \partial^{[\nu]}$, one has $m_{\nu} = n_{\nu+1}$. If $\sum_{\nu \in \mathbb{N}} n_{\nu} \partial^{[\nu]} \in \ker(t)$, then $n_{\nu+1} = m_{\nu} = 0$ for all ν . \square

Let $\text{Coh}(\mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^{\dagger})$ and $\text{Coh}^{\mathfrak{y}}(\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger})$ be the categories of coherent left $\mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^{\dagger}$ -modules and coherent left $\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}$ -modules with support in \mathfrak{y} , respectively.

PROPOSITION 3.4. —

- (i) *The functor i_+ restricts to a functor $i_+ : \text{Coh}(\mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^{\dagger}) \rightarrow \text{Coh}^{\mathfrak{y}}(\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger})$.*
- (ii) *The unit $\eta_{\mathcal{N}} : \mathcal{N} \xrightarrow{\sim} (i^{\sharp} \circ i_+) \mathcal{N}$ is an isomorphism for any $\mathcal{N} \in \text{Coh}(\mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^{\dagger})$.*
- (iii) *The functor $i_+ : \text{Coh}(\mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^{\dagger}) \rightarrow \text{Coh}^{\mathfrak{y}}(\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger})$ is fully faithful.*

Proof. — (i) is [4, 4.3.8], as we already stated above. Now (ii) is a local question and we may assume that \mathfrak{X} is affine equipped with local coordinates $t_1, \dots, t_d \in \mathcal{O}_{\mathfrak{X}}$ and that $\mathcal{I} = (t_{r+1}, \dots, t_d)$ for some $0 \leq r < d$. Let ∂_i be the derivation relative to t_i . We identify $\omega_{\mathfrak{X}}$ with $\mathcal{O}_{\mathfrak{X}}$ via the section $dt_1 \wedge \dots \wedge dt_d$ and similarly for $\omega_{\mathfrak{y}}$. It then follows from the existence of the adjoint operator [3, 1.2.2] and the fact that

$$\mathcal{D}_{\mathfrak{x} \rightarrow \mathfrak{y}, \mathbb{Q}}^{\dagger} = i^* \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger} = i^{-1}(\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger} / \mathcal{I} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}),$$

that we have an isomorphism of $(i^{-1} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}, \mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^{\dagger})$ -bimodules

$$\mathcal{D}_{\mathfrak{x} \leftarrow \mathfrak{y}, \mathbb{Q}}^{\dagger} \simeq i^{-1}(\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger} / \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger} \mathcal{I}).$$

It follows that $\varphi \mapsto \varphi(1)$ gives a natural identification

$$i^{\sharp} \mathcal{M} = \text{Hom}_{i^{-1} \mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger}}(\mathcal{D}_{\mathfrak{x} \leftarrow \mathfrak{y}, \mathbb{Q}}^{\dagger}, i^{-1} \mathcal{M}) \simeq \bigcap_{i=r+1, \dots, d} \ker(i^{-1} \mathcal{M} \xrightarrow{t_i} i^{-1} \mathcal{M}).$$

Suppose now that $\mathcal{M} = i_+ \mathcal{N}$. According to [2, 3.4.5] and [2, 3.6.2], there is an inductive system of coherent $\widehat{\mathcal{D}}_{\mathfrak{y}}^{(m)}$ -submodules $\mathcal{N}^{(m)} \subset \mathcal{N}$, such that

$\varinjlim_m \mathcal{N}^{(m)} = \mathcal{N}$. We have seen in the proof of Proposition 3.1, that there is a natural isomorphism

$$\mathcal{D}_{\mathfrak{x} \leftarrow \mathfrak{y}}^{(m)} \simeq \mathfrak{o}[\partial]^{(m)} \otimes_{\mathfrak{o}} \mathcal{D}_{\mathfrak{y}}^{(m)}$$

as right $\mathcal{D}_{\mathfrak{y}}^{(m)}$ -modules for any m . If we define an (injective) transition map $\mathfrak{o}[\partial]^{(m)} \rightarrow \mathfrak{o}[\partial]^{(m+1)}$ as in [2, 2.2.3.1], then the isomorphism is compatible with the transition maps $\mathcal{D}_{\mathfrak{x} \leftarrow \mathfrak{y}}^{(m)} \rightarrow \mathcal{D}_{\mathfrak{x} \leftarrow \mathfrak{y}}^{(m+1)}$ and $\mathcal{D}_{\mathfrak{y}}^{(m)} \rightarrow \mathcal{D}_{\mathfrak{y}}^{(m+1)}$. Hence, if $M = \mathcal{O}(\mathcal{M})$, $N = \mathcal{O}(\mathcal{N})$ and $N^{(m)} = \mathcal{O}(\mathcal{N}^{(m)})$ and if $\mathfrak{o}[\partial]^{(m)} \widehat{\otimes}_{\mathfrak{o}} N^{(m)}$ denotes the ϖ -adic completion of the \mathfrak{o} -module $\mathfrak{o}[\partial]^{(m)} \otimes_{\mathfrak{o}} N^{(m)}$, then

$$M \simeq \varinjlim_m (\mathfrak{o}[\partial]^{(m)} \widehat{\otimes}_{\mathfrak{o}} N^{(m)})_{\mathbb{Q}}.$$

Moreover, for $i = r + 1, \dots, d$, the action of $t_i \in \mathcal{O}_{\mathfrak{x}}$ on M is given on the right-hand side by the action on the left-hand factor $\mathfrak{o}[\partial]^{(m)}$. Since $\mathfrak{o}[\partial]^{(m)}$ is a free \mathfrak{o} -module, the tensor product $\mathfrak{o}[\partial]^{(m)} \otimes_{\mathfrak{o}} N^{(m)}$ is canonically isomorphic to the \mathfrak{o} -module of all finite formal sums $\sum_{\underline{\nu} \in \mathbb{N}^{d-r}} n_{\underline{\nu}} \partial^{(\underline{\nu})}$ with $n_{\underline{\nu}} \in N^{(m)}$. Its completion $\mathfrak{o}[\partial]^{(m)} \widehat{\otimes}_{\mathfrak{o}} N^{(m)}$ is therefore isomorphic to the \mathfrak{o} -module given by all formal infinite sums $\sum_{\underline{\nu} \in \mathbb{N}^{d-r}} n_{\underline{\nu}} \partial^{(\underline{\nu})}$ with $n_{\underline{\nu}} \in N^{(m)}$ and $n_{\underline{\nu}} \rightarrow 0$ in the ϖ -adic topology of $N^{(m)}$. Using the notation of Lemma 3.3 with $s = d - r$, we obtain thus an injective K -linear map from $(\mathfrak{o}[\partial]^{(m)} \widehat{\otimes}_{\mathfrak{o}} N^{(m)})_{\mathbb{Q}}$ into $N[[\partial^{[\underline{L}] }]]$. It is equivariant for the action of t_i on $N[[\partial^{[\underline{L}] }]]$ given by $t_i \cdot (n_{\underline{\nu}} \partial^{[\underline{L}]}) := n_{\underline{\nu}} (\partial^{[\underline{L} - e_i]})$ when $\nu_i > 0$ and zero else. Passing to the limit over m yields K -linear injection

$$M \longrightarrow N[[\partial^{[\underline{L}] }]],$$

which is equivariant for the action of t_i for all $i = r + 1, \dots, d$. According to Lemma 3.3 we obtain

$$N \subseteq \bigcap_{i=r+1, \dots, d} \ker(M \xrightarrow{t_i} M) \subseteq \bigcap_{i=r+1, \dots, d} \ker(N[[\partial^{[\underline{L}] }]] \xrightarrow{t_i} N[[\partial^{[\underline{L}] }]]) = N.$$

This implies $i^{\sharp} \mathcal{M} \simeq \mathcal{N}$. Hence the unit of the adjunction is an isomorphism. This shows (i). The statement (i) implies immediately that i_+ is faithful. For the fullness, let $\gamma : i_+(\mathcal{N}) \rightarrow i_+(\mathcal{N}')$ be a morphism. A preimage is given by the morphism

$$\eta_{\mathcal{N}'}^{-1} \circ i^{\sharp} (\varepsilon_{i_+(\mathcal{N}')} \circ i_+(\eta_{\mathcal{N}'}) \circ \gamma) \circ \eta_{\mathcal{N}} : \mathcal{N} \longrightarrow \mathcal{N}'$$

where $\varepsilon : i_+ \circ i^{\sharp} \rightarrow \text{id}$ is the counit of the adjunction, cf. the proof of [19, Tag 07RB]. \square

We now work towards the essential surjectivity of $i_+ : \text{Coh}(\mathcal{D}_{\mathfrak{y}, \mathbb{Q}}^{\dagger}) \rightarrow \text{Coh}^{\mathfrak{y}}(\mathcal{D}_{\mathfrak{x}, \mathbb{Q}}^{\dagger})$. Note that if i_+ is essentially surjective, then i^{\sharp} preserves coherence, by part (ii) of the preceding proposition.

LEMMA 3.5. — *Let $\mathfrak{Y} \subset \mathfrak{X}$ be of codimension 1 and let $\mathcal{M} \in \text{Coh}^{\mathfrak{Y}}(\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^{\dagger})$. The counit of the adjunction $\varepsilon_{\mathcal{M}} : (i_+ \circ i^{\natural})_{*}\mathcal{M} \rightarrow \mathcal{M}$ is surjective.*

Proof. — The surjectivity of $\varepsilon_{\mathcal{M}}$ is a local problem. We may therefore assume that \mathfrak{X} is an open affine with local coordinates x_1, \dots, x_M such that \mathfrak{S} is generated by $t := x_M$. According to [2, 3.6], we can assume that there is $m \geq 0$ such that

$$\mathcal{M} \simeq \mathcal{D}_{\mathfrak{X},\mathbb{Q}}^{\dagger} \otimes_{\widehat{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^{(m)}} \mathcal{M}_m$$

with a coherent $\widehat{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^{(m)}$ -module \mathcal{M}_m supported on \mathfrak{Y} . Let $\widehat{D}_{\mathfrak{X}}^{(m)} := \Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X}}^{(m)})$ and $\widehat{D}_{\mathfrak{X},\mathbb{Q}}^{(m)} := \Gamma(\mathfrak{X}, \widehat{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^{(m)})$. Let $M_m = \Gamma(\mathfrak{X}, \mathcal{M}_m)$. Let $\mathring{M}_m \subset M_m$ be a finitely generated $\widehat{D}_{\mathfrak{X}}^{(m)}$ -submodule such that $\widehat{D}_{\mathfrak{X},\mathbb{Q}}^{(m)} \mathring{M}_m = M_m$. Let $e_1, \dots, e_n \in \mathring{M}_m$ such that $\mathring{M}_m = \sum_i \widehat{D}_{\mathfrak{X}}^{(m)} e_i$. The module $\mathring{M}_m / \varpi \mathring{M}_m$ has support contained in \mathfrak{Y} and hence $t^{p^m} \bar{e}_i = 0$ for all i (increasing m if necessary), where $\bar{e}_i = e_i \bmod \varpi \mathring{M}_m$. By Corollary 2.16, there are generators e'_1, \dots, e'_n for the $\widehat{D}_{\mathfrak{X}}^{(m')}$ -module $\widehat{D}_{\mathfrak{X}}^{(m')} \otimes_{\widehat{D}_{\mathfrak{X}}^{(m)}} \mathring{M}_m$ with the property $t^{p^m} e'_i = 0$ for all i .

Now

$$\mathcal{M} \simeq \mathcal{D}_{\mathfrak{X},\mathbb{Q}}^{\dagger} \otimes_{\widehat{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^{(m')}} \mathcal{M}_{m'} \quad \text{with} \quad \mathcal{M}_{m'} := \widehat{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^{(m')} \otimes_{\widehat{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^{(m)}} \mathcal{M}_m.$$

Let $D_{\mathfrak{X},\mathbb{Q}}^{\dagger} = \Gamma(\mathfrak{X}, \mathcal{D}_{\mathfrak{X},\mathbb{Q}}^{\dagger})$, $M := \Gamma(\mathfrak{X}, \mathcal{M})$ and $M_0 := \Gamma(\mathfrak{X}, i^{\natural} \mathcal{M})$. Since $\Gamma(\mathfrak{X}, \mathcal{M}_{m'}) = (\mathring{M}_{m'})_{\mathbb{Q}}$, it is clear that the e'_i are generators for the $D_{\mathfrak{X},\mathbb{Q}}^{\dagger}$ -module M . As we have seen in the proof of Proposition 3.4, we have $M_0 = \ker(t) \subseteq M$. The counit $\varepsilon_{\mathcal{M}}$ is therefore surjective, if and only if $D_{\mathfrak{X},\mathbb{Q}}^{\dagger} M_0 = M$. This is the case if $e'_i \in D_{\mathfrak{X},\mathbb{Q}}^{\dagger} M_0$ for all i .

Since $t^{p^m} e'_i = 0$ for all i , it suffices to show the following claim: given an element $u \in M$ with $t^j u = 0$ for some $1 \leq j \leq p^m$, then $u \in D_{\mathfrak{X},\mathbb{Q}}^{\dagger} M_0$. To prove the claim, we use a finite induction on j , the case $j = 1$ being clear. So suppose $j > 1$ and that the statement holds for $j - 1$. We have

$$t^{j-1}(ju + t\partial u) = \partial(t^j)u + t^j \partial u = \partial(t^j)u = \partial(0) = 0$$

and so, by induction hypothesis, $ju + t\partial u \in D_{\mathfrak{X},\mathbb{Q}}^{\dagger} M_0$. Similarly, $t^{j-1}(tu) = 0$ implies $tu \in D_{\mathfrak{X},\mathbb{Q}}^{\dagger} M_0$ and hence also $-\partial(tu) \in D_{\mathfrak{X},\mathbb{Q}}^{\dagger} M_0$. Altogether,

$$(j-1)u = ju - u = ju + (t\partial - \partial t)u = ju + t\partial u - \partial(tu) \in D_{\mathfrak{X},\mathbb{Q}}^{\dagger} M_0.$$

and it remains to divide by $j - 1$. This completes the induction step and establishes the equality $D_{\mathfrak{X},\mathbb{Q}}^{\dagger} M_0 = M$. Hence the lemma is proved. \square

THEOREM 3.6 (Berthelot–Kashiwara theorem, left version). — *The functors i_+, i^\natural induce mutually inverse equivalences of categories*

$$\mathrm{Coh}(\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger) \begin{array}{c} \xrightarrow{i_+} \\ \xleftarrow[\simeq]{i^\natural} \end{array} \mathrm{Coh}^{\mathfrak{Y}}(\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger).$$

Proof. — We first suppose that $\mathfrak{Y} \subset \mathfrak{X}$ be of codimension 1. It suffices to show that the counit of the adjunction $\varepsilon_{\mathcal{M}} : (i_+ \circ i^\natural)\mathcal{M} \rightarrow \mathcal{M}$ is an isomorphism for any \mathcal{M} . This is a local question and we may assume that \mathfrak{X} is affine with coordinates x_1, \dots, x_d and that \mathcal{S} is generated by $t := x_d$. Since $(i_+ \circ i^\natural)\mathcal{M} \rightarrow \mathcal{M}$ is surjective according to Lemma 3.5 we may also assume that \mathcal{M} is globally generated, as $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ -module, by finitely many sections $e_1, \dots, e_n \in i^\natural\mathcal{M} = \ker(t) \subseteq \mathcal{M}$. Hence there is a free $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -module \mathcal{L} of rank n and a linear surjection

$$i_+\mathcal{L} \rightarrow \mathcal{M}.$$

Let \mathcal{K} be the kernel of this morphism, a coherent $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ -module with support in \mathfrak{Y} . The morphism $i_+i^\natural\mathcal{K} \rightarrow \mathcal{K}$ is surjective, again by Lemma 3.5. Similarly to the above, \mathcal{K} is therefore globally generated, as $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ -module, by finitely many sections $f_1, \dots, f_m \in i^\natural\mathcal{K} = \ker(t) \subseteq \mathcal{K}$. Consider the $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -submodule

$$\mathcal{V} := \sum_j \mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger f_j \subseteq i^\natural\mathcal{K}.$$

By construction, the composite map $i_+\mathcal{V} \rightarrow i_+i^\natural\mathcal{K} \rightarrow \mathcal{K}$ is a linear surjection

$$i_+\mathcal{V} \rightarrow \mathcal{K}.$$

Moreover,

$$\mathcal{V} \subseteq i^\natural\mathcal{K} \subseteq i^\natural i_+\mathcal{L} \simeq \mathcal{L},$$

where the second inclusion holds by left-exactness of i^\natural , cf. Proposition 3.2, and the final isomorphism holds by Proposition 3.4. Hence, $i_+\mathcal{V} \hookrightarrow i_+\mathcal{L}$ is injective with image \mathcal{K} . All in all,

$$i_+(\mathcal{L}/\mathcal{V}) \simeq i_+\mathcal{L}/i_+\mathcal{V} \simeq i_+\mathcal{L}/\mathcal{K} \simeq \mathcal{M}.$$

The $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -module \mathcal{L}/\mathcal{V} is finitely presented and hence coherent. So i_+ is essentially surjective. Moreover, $\mathcal{L}/\mathcal{V} \simeq (i^\natural \circ i_+)(\mathcal{L}/\mathcal{V}) \simeq i^\natural\mathcal{M}$. So the functor i^\natural takes $\mathrm{Coh}^{\mathfrak{Y}}(\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger)$ into $\mathrm{Coh}(\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger)$ and is a quasi-inverse to i_+ . This proves the theorem in case of codimension 1.

In the general case, we again reduce to the case where \mathfrak{X} is affine with coordinates x_1, \dots, x_d and that \mathcal{S} is generated by x_{r+1}, \dots, x_d . Define a

series of closed subschemes of \mathfrak{X} by $\mathfrak{Y}_k := V(x_{r+1}, \dots, x_{d-k-1})$ for $k = 1, \dots, d-r$, i.e.

$$\mathfrak{Y} = \mathfrak{Y}_1 \subset \mathfrak{Y}_2 \subset \dots \subset \mathfrak{Y}_{d-r-1} \subset \mathfrak{Y}_{d-r} \subset \mathfrak{X}$$

and $i_k : \mathfrak{Y}_k \subset \mathfrak{Y}_{k+1}$ is a closed immersion between smooth formal schemes of codimension 1. We use a finite induction on k . We call (S_k) the following statement : $(i_k)_+ \circ \dots \circ (i_1)_+$ induces an equivalence of categories between $\text{Coh}(\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger)$ and $\text{Coh}^{\mathfrak{Y}}(\mathcal{D}_{\mathfrak{Y}_k, \mathbb{Q}}^\dagger)$ with quasi-inverse $(i_1)^\sharp \circ \dots \circ (i_k)^\sharp$.

By the codimension 1 case, the functor $(i_{k+1})_+$ is an equivalence of categories between $\text{Coh}(\mathcal{D}_{\mathfrak{Y}_k, \mathbb{Q}}^\dagger)$ and $\text{Coh}^{\mathfrak{Y}_k}(\mathcal{D}_{\mathfrak{Y}_{k+1}, \mathbb{Q}}^\dagger)$ with quasi-inverse $(i_{k+1})^\sharp$. In particular, (S_1) is true. Suppose that (S_k) is true. According to Propositions 3.1 (iii) and 3.2 (iii), the functor $(i_{k+1})_+$ restricts to an equivalence between objects supported on \mathfrak{Y} , i.e. to an equivalence $\text{Coh}^{\mathfrak{Y}}(\mathcal{D}_{\mathfrak{Y}_k, \mathbb{Q}}^\dagger)$ and $\text{Coh}^{\mathfrak{Y}}(\mathcal{D}_{\mathfrak{Y}_{k+1}, \mathbb{Q}}^\dagger)$ with quasi-inverse $(i_{k+1})^\sharp$. This establishes (S_{k+1}) . The statement (S_{d-r}) then establishes the theorem, since $i_+ = (i_{d-r})_+ \circ \dots \circ (i_1)_+$ and $i^\sharp = (i_1)^\sharp \circ \dots \circ (i_{d-r})^\sharp$. To see the latter equality, note that all functors $i_+, (i_{d-r})_+, \dots, (i_1)_+$ are essentially surjective. According to part (ii) of Proposition 3.4, all functors $i^\sharp, (i_1)^\sharp, \dots, (i_{d-r})^\sharp$ therefore preserve coherent modules. But then $i^\sharp = (i_1)^\sharp \circ \dots \circ (i_{d-r})^\sharp$ by part (ii) of Proposition 3.2, as claimed. \square

3.3. Side-changing

We deduce the right version by using the side-changing functors $\omega_{\mathfrak{X}} \otimes_{\mathcal{O}_{\mathfrak{X}}} (-)$ and $\mathcal{H}\text{om}_{\mathcal{O}_{\mathfrak{X}}}(\omega_{\mathfrak{X}}, -)$, cf. Section 2.2. We consider the $(\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger, i^{-1}\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger)$ -bimodule $\mathcal{D}_{\mathfrak{X} \rightarrow \mathfrak{Y}, \mathbb{Q}}^\dagger := i^*\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$.

Denote by $i_{r,+}$ the functor from right $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -modules to right $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ -modules given by

$$i_{r,+}(\mathcal{N}) := \mathcal{N} \otimes_{\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger} \mathcal{D}_{\mathfrak{X} \rightarrow \mathfrak{Y}, \mathbb{Q}}^\dagger$$

as well as the functor from right $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ -modules to right $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -modules given by

$$i_r^\sharp(\mathcal{M}) := \mathcal{H}\text{om}_{i^{-1}\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger}(\mathcal{D}_{\mathfrak{X} \rightarrow \mathfrak{Y}, \mathbb{Q}}^\dagger, i^{-1}\mathcal{M}).$$

LEMMA 3.7. —

- (i) *The left $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -module $\mathcal{D}_{\mathfrak{X} \rightarrow \mathfrak{Y}, \mathbb{Q}}^\dagger$ is flat.*
- (ii) *One has a natural isomorphism*

$$i_{r,+}(\mathcal{N}) \simeq \omega_{\mathfrak{X}} \otimes_{\mathcal{O}_{\mathfrak{X}}} i_+(\mathcal{H}\text{om}_{\mathcal{O}_{\mathfrak{Y}}}(\omega_{\mathfrak{Y}}, \mathcal{N})).$$

(iii) *One has a natural isomorphism*

$$i_r^\natural(\mathcal{M}) \simeq \omega_{\mathfrak{Y}} \otimes_{\mathcal{O}_{\mathfrak{Y}}} i^\natural(\mathrm{Hom}_{\mathcal{O}_{\mathfrak{X}}}(\omega_{\mathfrak{X}}, \mathcal{M})).$$

Proof. — Given the definition of the transfer module [3, 3.4.1]

$$\mathcal{D}_{\mathfrak{X} \leftarrow \mathfrak{Y}, \mathbb{Q}}^\dagger = i^{-1}(\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger \otimes_{\mathcal{O}_{\mathfrak{X}}} \omega_{\mathfrak{X}}^{-1}) \otimes_{i^{-1}\mathcal{O}_{\mathfrak{X}}} \omega_{\mathfrak{Y}},$$

part (ii) follows formally exactly as in the classical case [10, Lem. 1.3.4]. Since i_+ is exact by Proposition 3.1, so is $i_{r,+}$, and then (ii) implies the flatness of the left $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -module $\mathcal{D}_{\mathfrak{X} \rightarrow \mathfrak{Y}, \mathbb{Q}}^\dagger$ (one may also give a direct argument along the lines of the proof of part (i) of Proposition 3.1). This shows (i). Finally, as in the proof of part (i) of Proposition 3.2, one verifies that $\mathrm{Hom}_{i^{-1}\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger}(\mathcal{D}_{\mathfrak{X} \rightarrow \mathfrak{Y}, \mathbb{Q}}^\dagger, i^{-1}(-))$ is right adjoint to $(-) \otimes_{\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger} \mathcal{D}_{\mathfrak{X} \rightarrow \mathfrak{Y}, \mathbb{Q}}^\dagger$ and so (iii) follows from unicity of adjoint functors. \square

Let $\mathrm{Coh}^r(\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger)$ and $\mathrm{Coh}^{r, \mathfrak{Y}}(\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger)$ be the categories of coherent right $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger$ -modules and coherent right $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$ -modules with support in \mathfrak{Y} , respectively.

THEOREM 3.8 (Berthelot–Kashiwara theorem, right version). — *The functors $i_{r,+}$, i_r^\natural induce mutually inverse equivalences of categories*

$$\mathrm{Coh}^r(\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^\dagger) \begin{array}{c} \xrightarrow{i_{r,+}} \\ \simeq \\ \xleftarrow{i_r^\natural} \end{array} \mathrm{Coh}^{r, \mathfrak{Y}}(\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger).$$

Proof. — Taking into account Proposition 2.2, this is a consequence of Theorem 3.6 and Lemma 3.7. \square

4. The Berthelot–Kashiwara theorem in the twisted case

4.1. Twisted sheaves

The following definition, adapted to arithmetic differential operators, is taken from [1]. Note that, given an $\mathcal{O}_{\mathfrak{X}, \mathbb{Q}}$ -ring $(\mathcal{A}, \iota_{\mathcal{A}})$, and an open subset $\mathfrak{U} \subset \mathfrak{X}$, the sheaf \mathcal{A} and the morphism $\iota_{\mathcal{A}}$ may be restricted to \mathfrak{U} and yield the $\mathcal{O}_{\mathfrak{U}, \mathbb{Q}}$ -ring $(\mathcal{A}|_{\mathfrak{U}}, \iota_{\mathcal{A}}|_{\mathfrak{U}})$. We say that two $\mathcal{O}_{\mathfrak{X}, \mathbb{Q}}$ -rings $(\mathcal{A}, \iota_{\mathcal{A}})$ and $(\mathcal{A}', \iota_{\mathcal{A}'})$ are *locally isomorphic*, if there is a (Zariski) open covering of \mathfrak{X} by open subsets \mathfrak{U} together with isomorphisms $(\mathcal{A}|_{\mathfrak{U}}, \iota_{\mathcal{A}}|_{\mathfrak{U}}) \simeq (\mathcal{A}'|_{\mathfrak{U}}, \iota_{\mathcal{A}'}|_{\mathfrak{U}})$ (in the category of $\mathcal{O}_{\mathfrak{U}, \mathbb{Q}}$ -rings, cf. Definition 2.1) for all \mathfrak{U} .

DEFINITION 4.1. — *A twisted sheaf of arithmetic differential operators on \mathfrak{X} is an $\mathcal{O}_{\mathfrak{X}, \mathbb{Q}}$ -ring \mathcal{A} , which is locally isomorphic to the $\mathcal{O}_{\mathfrak{X}, \mathbb{Q}}$ -ring $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^\dagger$.*

Let in the following \mathcal{A} be a twisted sheaf of arithmetic differential operators on \mathfrak{X} .

LEMMA 4.2. — *Let \mathcal{A}^{opp} be the opposite ring, i.e. the order of multiplication is reversed. Then \mathcal{A}^{opp} is an $\mathcal{O}_{\mathfrak{X},\mathbb{Q}}$ -ring.*

Proof. — Being a local statement, we can assume that \mathfrak{X} has étale coordinates. The existence of the adjoint operator says that $(\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger)^{opp}$ is isomorphic to $\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger$ [3, 1.2.2/3]. Since this holds even as $\mathcal{O}_{\mathfrak{X},\mathbb{Q}}$ -rings, the lemma follows. \square

DEFINITION 4.3. — *Let $\mathcal{J} \subseteq \mathcal{O}_{\mathfrak{X},\mathbb{Q}}$ be an ideal. The normalizer of the ideal $\mathcal{J}\mathcal{A}$ is the subset of \mathcal{A} equal to*

$$\mathcal{N}_{\mathcal{A}}(\mathcal{J}\mathcal{A}) := \{P \in \mathcal{A} \mid P\mathcal{J} \subseteq \mathcal{J}\mathcal{A}\}.$$

LEMMA 4.4. — *One has the following basic properties:*

- (i) $\mathcal{N}_{\mathcal{A}}(\mathcal{J}\mathcal{A})$ is a sub- $\mathcal{O}_{\mathfrak{X},\mathbb{Q}}$ -ring of \mathcal{A} .
- (ii) $\mathcal{J}\mathcal{A} \subseteq \mathcal{N}_{\mathcal{A}}(\mathcal{J}\mathcal{A})$ is a two-sided ideal.
- (iii) The quotient $\mathcal{N}_{\mathcal{A}}(\mathcal{J}\mathcal{A})/\mathcal{J}\mathcal{A}$ is an $\mathcal{O}_{\mathfrak{X},\mathbb{Q}}/\mathcal{J}$ -ring.
- (iv) $\mathcal{A}/\mathcal{J}\mathcal{A}$ is a $(\mathcal{N}_{\mathcal{A}}(\mathcal{J}\mathcal{A})/\mathcal{J}\mathcal{A}, \mathcal{A})$ -bimodule.

Proof. — The proof of (i)–(iii) is identical to the proof of Lemma 2.6. The point (iv) is easy to check. \square

Now let $i : \mathfrak{Y} \rightarrow \mathfrak{X}$ be a smooth closed formal subscheme defined by some coherent ideal $\mathcal{J} \subseteq \mathcal{O}_{\mathfrak{X}}$. According to Lemma 4.4 the sheaf on \mathfrak{Y}

$$\mathcal{A}_{\mathfrak{Y}} := i^{-1}(\mathcal{N}_{\mathcal{A}}(\mathcal{J}_{\mathbb{Q}}\mathcal{A})/\mathcal{J}_{\mathbb{Q}}\mathcal{A})$$

is an $\mathcal{O}_{\mathfrak{Y},\mathbb{Q}}$ -ring.

LEMMA 4.5. — *If $\mathcal{A} = \mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger$, then $\mathcal{A}_{\mathfrak{Y}} = \mathcal{D}_{\mathfrak{Y},\mathbb{Q}}^\dagger$.*

Proof. — This is Corollary 2.7. \square

COROLLARY 4.6. — $\mathcal{A}_{\mathfrak{Y}}$ is a twisted sheaf of arithmetic differential operators on \mathfrak{Y} .

Before we proceed, we recall an elementary lemma.

LEMMA 4.7. — *Let X be a topological space and $\mathcal{F}_1, \mathcal{F}_2 \subseteq \mathcal{G}$ abelian sheaves. If $\mathcal{F}_{1,x} = \mathcal{F}_{2,x}$ inside the stalk \mathcal{G}_x for all $x \in X$, then $\mathcal{F}_1 = \mathcal{F}_2$.*

Proof. — By symmetry, it suffices to see $\mathcal{F}_1 \subseteq \mathcal{F}_2$. The stalk of the sheaf $(\mathcal{F}_1 + \mathcal{F}_2)/\mathcal{F}_2$ vanishes at every $x \in X$, so that $(\mathcal{F}_1 + \mathcal{F}_2)/\mathcal{F}_2 = 0$, i.e. $\mathcal{F}_1 \subseteq \mathcal{F}_2$. \square

PROPOSITION 4.8. — *Let $\mathfrak{Z} \xrightarrow{k} \mathfrak{Y} \xrightarrow{i} \mathfrak{X}$ be closed immersions of smooth formal \mathfrak{o} -schemes. There is a canonical isomorphism $(\mathcal{A}_{\mathfrak{Y}})_{\mathfrak{Z}} \simeq \mathcal{A}_{\mathfrak{Z}}$ as $\mathcal{O}_{\mathfrak{Z}, \mathbb{Q}}$ -rings.*

Proof. — Let $\mathcal{I} \subseteq \mathcal{O}_{\mathfrak{X}}$ be the ideal defining the closed immersion $i \circ k : \mathfrak{Z} \rightarrow \mathfrak{X}$. In particular, $\mathcal{I} \subseteq \mathcal{J}$. Let $\overline{\mathcal{J}}$ be the image of $i^{-1}(\mathcal{J}/\mathcal{I})$ in $\mathcal{O}_{\mathfrak{Y}, \mathbb{Q}}$. By construction, the rings $(\mathcal{A}_{\mathfrak{Y}})_{\mathfrak{Z}}$ and $\mathcal{A}_{\mathfrak{Y}}$ are contained in $k^*(\mathcal{A}_{\mathfrak{Y}})$ and $i^*(\mathcal{A})$ respectively. Hence there is an injective ring homomorphism

$$\varphi : (\mathcal{A}_{\mathfrak{Y}})_{\mathfrak{Z}} \longrightarrow (i \circ k)^*(\mathcal{A})$$

compatible with the $\mathcal{O}_{\mathfrak{Z}}$ -structures. It suffices to see $\text{im}(\varphi) = \mathcal{A}_{\mathfrak{Z}}$. This is a local question, by Lemma 4.7, and so we may assume $\mathcal{A} \simeq \mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^{\dagger}$. Then $\mathcal{A}_{\mathfrak{Y}} \simeq \mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^{\dagger}$ and $\mathcal{A}_{\mathfrak{Z}} \simeq \mathcal{D}_{\mathfrak{Z}, \mathbb{Q}}^{\dagger}$ by Lemma 4.5 and φ becomes the canonical injective morphism

$$k^{-1}(\mathcal{N}_{\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^{\dagger}}(\overline{\mathcal{J}} \mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^{\dagger}) / \overline{\mathcal{J}} \mathcal{D}_{\mathfrak{Y}, \mathbb{Q}}^{\dagger}) \longrightarrow (i \circ k)^*(\mathcal{D}_{\mathfrak{X}, \mathbb{Q}}^{\dagger}).$$

Its image equals $\mathcal{D}_{\mathfrak{Z}, \mathbb{Q}}^{\dagger}$, according to Corollary 2.7. □

In the following we take the canonical isomorphism $(\mathcal{A}_{\mathfrak{Y}})_{\mathfrak{Z}} \simeq \mathcal{A}_{\mathfrak{Z}}$ as an identification.

4.2. Direct image and the main theorem

Let

$$i : \mathfrak{Y} \longrightarrow \mathfrak{X}$$

be a closed immersion between smooth formal \mathfrak{o} -schemes defined by some ideal $\mathcal{I} \subset \mathcal{O}_{\mathfrak{X}}$. Let \mathcal{A} be a twisted sheaf of arithmetic differential operators on \mathfrak{X} . We have the $(\mathcal{N}_{\mathcal{A}}(\mathcal{I}\mathcal{A})/\mathcal{I}\mathcal{A}, \mathcal{A})$ -bimodule $\mathcal{A}/\mathcal{I}\mathcal{A}$, cf. Lemma 4.4.

DEFINITION 4.9. — *The transfer bimodule along i is the $(\mathcal{A}_{\mathfrak{Y}}, i^{-1}\mathcal{A})$ -bimodule*

$$\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}} := i^*(\mathcal{A}) = i^{-1}(\mathcal{A}/\mathcal{I}\mathcal{A}).$$

PROPOSITION 4.10. — *Let $\mathfrak{Z} \xrightarrow{k} \mathfrak{Y} \xrightarrow{i} \mathfrak{X}$ be closed immersions of smooth formal \mathfrak{o} -schemes. There is a natural isomorphism as $(\mathcal{A}_{\mathfrak{Z}}, (i \circ k)^{-1}\mathcal{A})$ -bimodules*

$$\mathcal{A}_{\mathfrak{Z} \rightarrow \mathfrak{Y}} \otimes_{k^{-1}\mathcal{A}_{\mathfrak{Y}}} k^{-1}\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}} \simeq \mathcal{A}_{\mathfrak{Z} \rightarrow \mathfrak{X}}.$$

Proof. — One has

$$\begin{aligned}
 \mathcal{A}_{\mathfrak{Z} \rightarrow \mathfrak{Y}} \otimes_{k^{-1}\mathcal{A}_{\mathfrak{Y}}} k^{-1}\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}} \\
 &\simeq (\mathcal{O}_{\mathfrak{Z}} \otimes_{k^{-1}\mathcal{O}_{\mathfrak{Y}}} k^{-1}\mathcal{A}_{\mathfrak{Y}}) \otimes_{k^{-1}\mathcal{A}_{\mathfrak{Y}}} k^{-1}(\mathcal{O}_{\mathfrak{Y}} \otimes_{i^{-1}\mathcal{O}_{\mathfrak{X}}} i^{-1}\mathcal{A}) \\
 &\simeq (\mathcal{O}_{\mathfrak{Z}} \otimes_{k^{-1}\mathcal{O}_{\mathfrak{Y}}} k^{-1}\mathcal{A}_{\mathfrak{Y}}) \otimes_{k^{-1}\mathcal{A}_{\mathfrak{Y}}} (k^{-1}\mathcal{O}_{\mathfrak{Y}} \otimes_{(i \circ k)^{-1}\mathcal{O}_{\mathfrak{X}}} (i \circ k)^{-1}\mathcal{A}) \\
 &\simeq \mathcal{O}_{\mathfrak{Z}} \otimes_{(i \circ k)^{-1}\mathcal{O}_{\mathfrak{X}}} (i \circ k)^{-1}\mathcal{A} = \mathcal{A}_{\mathfrak{Z} \rightarrow \mathfrak{X}}.
 \end{aligned}$$

In the middle isomorphism we have used the compatibility of inverse images with tensor products, e.g. [10, C.1.11 (i)]. \square

DEFINITION 4.11. — Let \mathcal{M} be a right $\mathcal{A}_{\mathfrak{Y}}$ -module. Its direct image along i is the right \mathcal{A} -module

$$i_{\mathcal{A},+}\mathcal{M} := i_*(\mathcal{M} \otimes_{\mathcal{A}_{\mathfrak{Y}}} \mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}}).$$

It is clear that this yields a functor $i_{\mathcal{A},+}$ from right $\mathcal{A}_{\mathfrak{Y}}$ -modules to right \mathcal{A} -modules.

LEMMA 4.12. —

- (i) The left $\mathcal{A}_{\mathfrak{Y}}$ -module $\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}}$ is flat.
- (ii) The functor $i_{\mathcal{A},+}$ is exact.

Proof. — Part (i) is a local question and so we may assume $\mathcal{A} = \mathcal{D}_{\mathfrak{X}}^{\dagger}$ and $\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}} = \mathcal{D}_{\mathfrak{Y} \rightarrow \mathfrak{X}}^{\dagger}$. The claim follows then from part (i) of Lemma 3.7. Since i_* is exact, (i) implies (ii). \square

PROPOSITION 4.13. — Let $\mathfrak{Z} \xrightarrow{k} \mathfrak{Y} \xrightarrow{i} \mathfrak{X}$ be closed immersions of smooth formal \mathfrak{o} -schemes. There is a natural isomorphism $(i \circ k)_{\mathcal{A},+} = i_{\mathcal{A},+} \circ k_{\mathcal{A},+}$ as functors from right $\mathcal{A}_{\mathfrak{Z}}$ -modules to right \mathcal{A} -modules.

Proof. — Using the Proposition 4.10 one finds

$$\begin{aligned}
 (i \circ k)_{\mathcal{A},+}(\mathcal{M}) &\simeq (i \circ k)_*(\mathcal{M} \otimes_{\mathcal{A}_{\mathfrak{Z}}} \mathcal{A}_{\mathfrak{Z} \rightarrow \mathfrak{X}}) \\
 &\simeq i_*(k_*(\mathcal{M} \otimes_{\mathcal{A}_{\mathfrak{Z}}} (\mathcal{A}_{\mathfrak{Z} \rightarrow \mathfrak{Y}} \otimes_{k^{-1}\mathcal{A}_{\mathfrak{Y}}} k^{-1}\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}}))) \\
 &\simeq i_*(k_*(\mathcal{M} \otimes_{\mathcal{A}_{\mathfrak{Z}}} \mathcal{A}_{\mathfrak{Z} \rightarrow \mathfrak{Y}}) \otimes_{\mathcal{A}_{\mathfrak{Y}}} \mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}}) \\
 &\simeq i_{\mathcal{A},+}(k_{\mathcal{A},+}(\mathcal{M})).
 \end{aligned}$$

In the third isomorphism, we have used the projection formula [10, C.1.11 (iii)]. For this, note that $k! = k_*$ and that $\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}}$ is flat as left $\mathcal{A}_{\mathfrak{Y}}$ -module Lemma 4.12. \square

We define the following functor from right \mathcal{A} -modules to right $\mathcal{A}_{\mathfrak{Y}}$ -modules:

$$i_{\mathcal{A}}^{\natural}\mathcal{M} := \mathcal{H}om_{i^{-1}\mathcal{A}}(\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}}, i^{-1}\mathcal{M}).$$

PROPOSITION 4.14. — *The functor $i_{\mathcal{A},+}$ has a right adjoint, given by the functor $i_{\mathcal{A}}^{\natural}$.*

Proof. — This is a sort of right version of the argument given in part (i) of Proposition 3.2. Since i is a closed immersion, i_* has the right adjoint i^{-1} . Therefore, for any right $\mathcal{A}_{\mathfrak{Y}}$ -module \mathcal{N} and any right \mathcal{A} -module \mathcal{M} , one has

$$\mathrm{Hom}_{\mathcal{A}}(i_{\mathcal{A},+}\mathcal{N}, \mathcal{M}) = \mathrm{Hom}_{i^{-1}\mathcal{A}}(\mathcal{N} \otimes_{\mathcal{A}_{\mathfrak{Y}}} \mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}}, i^{-1}\mathcal{M}).$$

One obtains (i) by combining this with the standard tensor-hom adjunction

$$\begin{aligned} \mathrm{Hom}_{i^{-1}\mathcal{A}}(\mathcal{N} \otimes_{\mathcal{A}_{\mathfrak{Y}}} \mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}}, i^{-1}\mathcal{M}) \\ = \mathrm{Hom}_{\mathcal{A}_{\mathfrak{Y}}}(\mathcal{N}, \mathrm{Hom}_{i^{-1}\mathcal{A}}(\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}}, i^{-1}\mathcal{M})). \quad \square \end{aligned}$$

COROLLARY 4.15. — *Let $\mathfrak{Z} \xrightarrow{k} \mathfrak{Y} \xrightarrow{i} \mathfrak{X}$ be closed immersions of smooth formal \mathfrak{o} -schemes. Then $(i \circ k)_{\mathcal{A}}^{\natural} = k_{\mathcal{A}}^{\natural} \circ i_{\mathcal{A}}^{\natural}$ as functors from right \mathcal{A} -modules to right $\mathcal{A}_{\mathfrak{Z}}$ -modules.*

Proof. — This follows from and Proposition 4.13 and Proposition 4.14 by uniqueness of adjoint functors. \square

We come to the main result of this paper. Let $\mathrm{Coh}^r(\mathcal{A}_{\mathfrak{Y}})$ and $\mathrm{Coh}^{r,\mathfrak{Y}}(\mathcal{A})$ be the categories of coherent right $\mathcal{A}_{\mathfrak{Y}}$ -modules and coherent right \mathcal{A} -modules with support in \mathfrak{Y} , respectively.

THEOREM 4.16 (Berthelot–Kashiwara for twisted sheaves). — *The functors $i_{\mathcal{A},+}, i_{\mathcal{A}}^{\natural}$ induces mutually inverse equivalences of categories*

$$\mathrm{Coh}^r(\mathcal{A}_{\mathfrak{Y}}) \begin{array}{c} \xrightarrow{i_{\mathcal{A},+}} \\ \xleftarrow[\simeq]{} \\ \xleftarrow{i_{\mathcal{A}}^{\natural}} \end{array} \mathrm{Coh}^{r,\mathfrak{Y}}(\mathcal{A}).$$

Proof. — Let \mathcal{N} be a coherent right $\mathcal{A}_{\mathfrak{Y}}$ -module and \mathcal{M} be a coherent right \mathcal{A} -module supported on \mathfrak{Y} . It suffices to see that the counit $i_{\mathcal{A},+} \circ i_{\mathcal{A}}^{\natural}(\mathcal{M}) \rightarrow \mathcal{M}$ and the unit $\mathcal{N} \rightarrow i_{\mathcal{A}}^{\natural} \circ i_{\mathcal{A},+}(\mathcal{N})$ of the adjunction are isomorphisms. These are local problems and we may therefore suppose that $\mathcal{A} = \mathcal{D}_{\mathfrak{X},\mathbb{Q}}^{\dagger}$. Then $\mathcal{A}_{\mathfrak{Y} \rightarrow \mathfrak{X}} = \mathcal{D}_{\mathfrak{Y} \rightarrow \mathfrak{X},\mathbb{Q}}^{\dagger}$ and the pair $(i_{\mathcal{A},+}, i_{\mathcal{A}}^{\natural})$ becomes the adjoint pair $(i_{r,+}, i_r^{\natural})$ studied in Section 3.3. Hence Theorem 3.8 implies the assertions. \square

5. Applications to crystalline distribution algebras

Let G be a connected split reductive group scheme over \mathfrak{o} .

5.1. The crystalline distribution algebra

We briefly review the construction of the crystalline distribution algebra of G , as introduced in [12]. Let I be the kernel of the morphism of \mathfrak{o} -algebras $\varepsilon_G : \mathfrak{o}[G] \rightarrow \mathfrak{o}$ which represents $1 \in G$. Then I/I^2 is a free $\mathfrak{o} = \mathfrak{o}[G]/I$ -module of finite rank. Let $t_1, \dots, t_N \in I$ whose classes modulo I^2 form a base of I/I^2 . Let $m \geq 0$ be an integer. The m -PD-envelope of I is denoted by $P_{(m)}(G)$. This algebra is a free \mathfrak{o} -module with basis

$$\underline{t}^{\{k\}} = t_1^{\{k_1\}} \dots t_N^{\{k_N\}},$$

where $q_i!t_i^{\{k_i\}} = t_i^{k_i}$ with $i = p^m q_i + r$ et $r < p^m$ [2, 1.5]. The algebra $P_{(m)}(G)$ has a descending filtration by the ideals $I^{\{n\}} = \bigoplus_{|k| \geq n} \mathfrak{o} \cdot \underline{t}^{\{k\}}$. The quotients $P_{(m)}^n(G) := P_{(m)}(G)/I^{\{n+1\}}$ are generated, as \mathfrak{o} -module, by the elements $\underline{t}^{\{k\}}$ where $|k| \leq n$ and there is an isomorphism $P_{(m)}^n(G) \simeq \bigoplus_{|k| \leq n} \mathfrak{o} \underline{t}^{\{k\}}$ as \mathfrak{o} -modules. There are canonical surjections $pr^{n+1, n} : P_{(m)}^{n+1}(G) \twoheadrightarrow P_{(m)}^n(G)$.

We note

$$\text{Lie}(G) := \text{Hom}_{\mathfrak{o}}(I/I^2, \mathfrak{o}).$$

The Lie-algebra $\text{Lie}(G)$ is a free \mathfrak{o} -module with basis ξ_1, \dots, ξ_N dual to t_1, \dots, t_N . For $m' \geq m$, the universal property of divided power algebras of level m gives homomorphisms of filtered algebras $\psi_{m, m'} : P_{(m')}^n(G) \rightarrow P_{(m)}^n(G)$ which induce on quotients homomorphisms of algebras $\psi_{m, m'}^n : P_{(m')}^n(G) \rightarrow P_{(m)}^n(G)$. *The module of distributions of level m and order n is $D_n^{(m)}(G) := \text{Hom}_{\mathfrak{o}}(P_{(m)}^n(G), \mathfrak{o})$ The algebra of distributions of level m is defined to be*

$$D^{(m)}(G) := \varinjlim_n D_n^{(m)}(G)$$

where the limit is taken with respect to the maps $\text{Hom}_{\mathfrak{o}}(pr^{n+1, n}, \mathfrak{o})$.

For $m' \geq m$, the algebra homomorphisms $\psi_{m, m'}^n$ give dually linear maps $\Phi_{m, m'}^n : D_n^{(m)}(G) \rightarrow D_n^{(m')}(G)$ and finally a morphism of filtered algebras $\Phi_{m, m'} : D^{(m)}(G) \rightarrow D^{(m')}(G)$. The direct limit $\text{Dist}(G) = \varinjlim_m D^{(m)}(G)$ equals the classical distribution algebra of the group scheme G [8, II.§4.6.1]. Instead of passing to this limit, we let \mathcal{G} be the completion of G along its special fibre. We write $G_i = \text{Spec } \mathfrak{o}[G]/\pi^{i+1}$. The morphism $G_{i+1} \hookrightarrow G_i$ induces $D^{(m)}(G_{i+1}) \rightarrow D^{(m)}(G_i)$. We put

$$\widehat{D}^{(m)}(\mathcal{G}) := \varprojlim_i D^{(m)}(G_i).$$

If $m' \geq m$, one has the morphisms $\widehat{\Phi}_{m,m'} : \widehat{D}^{(m)}(\mathcal{G}) \rightarrow \widehat{D}^{(m')}(\mathcal{G})$ and one can define the *crystalline distribution algebra* of \mathcal{G} as

$$D^\dagger(\mathcal{G})_{\mathbb{Q}} := \varinjlim_m \widehat{D}^{(m)}(\mathcal{G}) \otimes \mathbb{Q}.$$

5.2. Twisted sheaves on the flag variety

We let $B \subset G$ be a Borel subgroup scheme containing a maximal split torus T , with unipotent radical N . Let $X := G/B$ be the flag scheme and let $\widetilde{X} := G/N$. The right translation action of T on \widetilde{X} makes the projection

$$\xi : \widetilde{X} \longrightarrow X$$

a T -torsor over X .

We exhibit a certain class of twisted sheaves of arithmetic differential operators on the p -adic completion \mathfrak{X} of X . This construction goes back to the fundamental work of Beilinson–Bernstein [1] and Borho–Brylinski [5]. It was adapted to the setting of arithmetic differential operators by Sarrazola–Alzate [18].⁽¹⁾

Let \mathcal{T} and $\widetilde{\mathfrak{X}}$ be the completion of T and \widetilde{X} respectively. Then \mathcal{T} acts from the right on $\widetilde{\mathfrak{X}}$. We also write ξ for the projection morphism $\widetilde{\mathfrak{X}} \rightarrow \mathfrak{X}$ arising from ξ by completion. We put

$$\widetilde{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^\dagger := (\xi_*(\mathcal{D}_{\widetilde{\mathfrak{X},\mathbb{Q}}}^\dagger))^{\mathcal{T}}.$$

The right \mathcal{T} -action on $\widetilde{\mathfrak{X}}$ leads to a central embedding $D^\dagger(\mathcal{T})_{\mathbb{Q}} \rightarrow \widetilde{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^\dagger$.

Now we fix a character of the crystalline distribution algebra $D^\dagger(\mathcal{T})_{\mathbb{Q}}$ of \mathcal{T} , i.e. a homomorphism

$$\lambda : D^\dagger(\mathcal{T})_{\mathbb{Q}} \longrightarrow K.$$

Note that, by restriction, λ may be viewed as a weight, i.e. a linear form of the Cartan subalgebra

$$\mathfrak{t} := \mathrm{Lie}(T) \otimes \mathbb{Q} \subset \mathfrak{g} = \mathrm{Lie}(G) \otimes \mathbb{Q},$$

but not every weight extends to a character of $D^\dagger(\mathcal{T})_{\mathbb{Q}}$. If $\lambda \in X(T)$, i.e. λ lifts to an algebraic character of T , then we denote the corresponding \mathcal{G} -equivariant line bundle on \mathfrak{X} (with p inverted) by $\mathcal{O}_{\mathfrak{X},\mathbb{Q},\lambda}$. Finally, we denote the trivial character (which restricts to zero on $\mathrm{Lie}(T)$) by $\lambda = 0$.

⁽¹⁾ Note that [18] assumes $\mathfrak{o} = \mathbb{Z}_p$, however, a large part of the results and constructions are valid in full generality.

We define

$$\mathcal{D}_{\mathfrak{X},\mathbb{Q},\lambda}^\dagger := \widetilde{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^\dagger \otimes_{D^\dagger(\mathcal{T})_{\mathbb{Q},\lambda}} K,$$

compare [18, Def. 5.0.1]. Since $(\mathcal{D}_{\mathcal{T},\mathbb{Q}}^\dagger)^\mathcal{T} = D^\dagger(\mathcal{T})_{\mathbb{Q}}$ according to [12, Thm. 4.4.9.2], the ring $\widetilde{\mathcal{D}}_{\mathfrak{X},\mathbb{Q}}^\dagger$ is locally, on an open subset trivializing the torsor, isomorphic to $\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger \otimes_K D^\dagger(\mathcal{T})_{\mathbb{Q}}$. It follows that $\mathcal{D}_{\mathfrak{X},\mathbb{Q},\lambda}^\dagger$ is indeed locally isomorphic to $\mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger$, i.e. $\mathcal{D}_{\mathfrak{X},\mathbb{Q},\lambda}^\dagger$ is a twisted sheaf of arithmetic differential operators on \mathfrak{X} depending on the character λ . If $\lambda \in X(T)$, then there is a natural left action of $\mathcal{D}_{\mathfrak{X},\mathbb{Q},\lambda}^\dagger$ on $\mathcal{O}_{\mathfrak{X},\mathbb{Q},\lambda}$. For $\lambda = 0$ one recovers $\mathcal{D}_{\mathfrak{X},\mathbb{Q},0}^\dagger = \mathcal{D}_{\mathfrak{X},\mathbb{Q}}^\dagger$.

We need to determine the opposite ring of $\mathcal{D}_{\mathfrak{X},\mathbb{Q},\lambda}^\dagger$ in terms of the weight λ . To do this, for any ring A , we denote by A^{opp} its opposite ring, i.e. the same underlying abelian group, but where the order of multiplication reversed. For any \mathbb{Z}_p -algebra A , we denote by \widehat{A} its p -adic completion and let $\widehat{A}_{\mathbb{Q}} := \widehat{A} \otimes_{\mathbb{Q}}$.

LEMMA 5.1. — *There are ring isomorphisms $(\widehat{A})^{opp} \simeq \widehat{A^{opp}}$ and $(A^{opp})_{\mathbb{Q}} = (A_{\mathbb{Q}})^{opp}$.*

Proof. — The canonical ring homomorphism $A \rightarrow (\widehat{A^{opp}})^{opp}$ extends to a bijection $\widehat{A} \simeq (\widehat{A^{opp}})^{opp}$. Passing to opposite rings gives the first claim. The argument in the second case is similar. \square

Let $\rho = \frac{1}{2} \sum_{\Phi^+} \alpha$ be half the sum over the positive roots of (G, T) with respect to B .

PROPOSITION 5.2. — *There is a ring isomorphism $(\mathcal{D}_{\mathfrak{X},\mathbb{Q},\lambda}^\dagger)^{opp} \simeq \mathcal{D}_{\mathfrak{X},\mathbb{Q},2\rho-\lambda}^\dagger$.*

Proof. — By construction, one has $\mathcal{D}_{\mathfrak{X},\mathbb{Q},\lambda}^\dagger = \varinjlim_{m \geq 0} \widehat{\mathcal{D}}_{\mathfrak{X},\mathbb{Q},\lambda}^{(m)}$ with twisted sheaves $\mathcal{D}_{X,\lambda}^{(m)}$, their p -adic completions $\widehat{\mathcal{D}}_{\mathfrak{X},\lambda}^{(m)}$ and their corresponding \mathbb{Q} -algebras $\widehat{\mathcal{D}}_{\mathfrak{X},\mathbb{Q},\lambda}^{(m)}$. By the preceding lemma, it suffices to fix m and to show isomorphisms $(\mathcal{D}_{X,\lambda}^{(m)})^{opp} \simeq \mathcal{D}_{X,2\rho-\lambda}^{(m)}$ compatible with variation in m . Let $\mathcal{A}_X^{(m)} := \mathcal{O}_X \otimes_{\circ} D^{(m)}(G)$. According to [18, 3.5.13], there is a surjective morphism of \mathcal{O}_X -rings

$$\Phi_{X,\lambda}^{(m)} : \mathcal{A}_X^{(m)} \longrightarrow \mathcal{D}_{X,\lambda}^{(m)}$$

which gives, upon inversion of p and restriction to the generic fibre $\iota : X_{\mathbb{Q}} \hookrightarrow X$, the classical morphism $\Phi_{X_{\mathbb{Q}},\lambda} : \mathcal{O}_{X_{\mathbb{Q}}} \otimes_K U(\mathfrak{g}) \rightarrow D_{X_{\mathbb{Q}},\lambda}$. Following [15, 4.15.1], we denote by φ the anti-isomorphism of the $\mathcal{O}_{X_{\mathbb{Q}}}$ -ring $\mathcal{O}_{X_{\mathbb{Q}}} \otimes_K U(\mathfrak{g})$ induced by $x \mapsto -x$ on \mathfrak{g} . It restricts to an anti-isomorphism of $\mathcal{A}_X^{(m)}$, which we also denote by φ . Let $\mathcal{J}_{\lambda}^{(m)} := \ker \Phi_{X,\lambda}^{(m)}$ and $\mathcal{J}_{\lambda} := \ker \Phi_{X_{\mathbb{Q}},\lambda}$.

Then $\mathcal{I}_\lambda^{(m)} = \mathcal{A}_X^{(m)} \cap \iota_* \mathcal{I}_\lambda$. Following [1], we denote by \mathfrak{b}° the kernel of the canonical morphism $\alpha : \mathcal{O}_{X_{\mathbb{Q}}} \otimes_K \mathfrak{g} \rightarrow D_{X_{\mathbb{Q}}}$ and by λ° the morphism $\mathfrak{b}^\circ \rightarrow \mathcal{O}_{X_{\mathbb{Q}}}$ induced by λ (using that $\mathfrak{b}^\circ / [\mathfrak{b}^\circ, \mathfrak{b}^\circ] \simeq \mathcal{O}_{X_{\mathbb{Q}}} \otimes_K \mathfrak{t}$). The two-sided ideal \mathcal{I} then equals the right-ideal generated by $\ker \lambda^\circ$. By [15, 4.15.1] the anti-isomorphism φ maps $\ker \lambda^\circ$ to $\ker(2\rho - \lambda)^\circ$ (note that Kashiwara writes $\tilde{\mathfrak{g}}$ for \mathfrak{b}° , cf. [15, 4.3]). It follows $\varphi(\mathcal{I}_\lambda) = \mathcal{I}_{2\rho - \lambda}$ and this implies $\varphi(\mathcal{I}_\lambda^{(m)}) = \mathcal{I}_{2\rho - \lambda}^{(m)}$. This gives an isomorphism $(\mathcal{D}_{X, \lambda}^{(m)})^{opp} \simeq \mathcal{D}_{X, 2\rho - \lambda}^{(m)}$. Compatibility with m may be checked after inverting p and restricting to the generic fibre $X_{\mathbb{Q}}$. But then, by construction, the isomorphism coincides with the classical isomorphism $D_{X_{\mathbb{Q}}, \lambda} \simeq D_{X_{\mathbb{Q}}, 2\rho - \lambda}$ [15, 4.15.1]. \square

Let in the following

$$i : \mathfrak{Y} \longrightarrow \mathfrak{X}$$

be a smooth closed formal subscheme defined by some coherent ideal $\mathcal{I} \subseteq \mathcal{O}_{\mathfrak{X}}$. According to Corollary 4.6 and Lemma 4.2 the sheaf

$$\mathcal{O}_{\mathfrak{Y}, \mathbb{Q}, \lambda}^\dagger := i^{-1}(\mathcal{N}_{\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger}(\mathcal{I} \mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger) / \mathcal{I} \mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger)^{opp}$$

is a twisted sheaf of arithmetic differential operators on \mathfrak{Y} . If $\lambda \in X(T)$, we let

$$\mathcal{O}_{\mathfrak{Y}, \mathbb{Q}, \lambda} := i^* \mathcal{O}_{\mathfrak{X}, \mathbb{Q}, \lambda} = i^{-1}(\mathcal{O}_{\mathfrak{X}, \mathbb{Q}, \lambda} / \mathcal{I} \mathcal{O}_{\mathfrak{X}, \mathbb{Q}, \lambda}).$$

It is a line bundle on \mathfrak{Y} .

PROPOSITION 5.3. — *Let $\lambda \in X(T)$.*

- (i) *The line bundle $\mathcal{O}_{\mathfrak{Y}, \mathbb{Q}, \lambda}$ is naturally a left $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}, \lambda}^\dagger$ -module.*
- (ii) *Assume that the special fibre and the rigid-analytic generic fibre of \mathfrak{Y} are connected. Then $\mathcal{O}_{\mathfrak{Y}, \mathbb{Q}, \lambda}$ is a simple left $\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}, \lambda}^\dagger$ -module.*

Proof. — We recall that $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger$ naturally acts from the left on $\mathcal{O}_{\mathfrak{X}, \lambda, \mathbb{Q}}$. Let $f \in \mathcal{O}_{\mathfrak{X}, \lambda, \mathbb{Q}}$ and $P \in \mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger$ be local sections. Identifying the $\mathcal{O}_{\mathfrak{X}, \mathbb{Q}}$ -ring $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger$ with $(\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger)^{opp}$, there is a well-defined local section $P(f) \in \mathcal{O}_{\mathfrak{X}, \mathbb{Q}, \lambda}$. Whether or not the subset $\mathcal{N}_{\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger}(\mathcal{I} \mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger)$ stabilizes the submodule $\mathcal{I} \mathcal{O}_{\mathfrak{X}, \mathbb{Q}, \lambda} \subseteq \mathcal{O}_{\mathfrak{X}, \mathbb{Q}, \lambda}$ is a local question. We may hence fix a local $\mathcal{O}_{\mathfrak{X}, \mathbb{Q}}$ -linear isomorphism between $\mathcal{O}_{\mathfrak{X}, \mathbb{Q}, \lambda}$ and $\mathcal{O}_{\mathfrak{X}, \mathbb{Q}}$. If then $P \in \mathcal{N}_{\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger}(\mathcal{I} \mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger)$ and $f \in \mathcal{I}_{\mathbb{Q}}$, then there is $Q \in \mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger$ and $h \in \mathcal{I}_{\mathbb{Q}}$ such that $Pf = hQ$. It follows

$$P(f) = Pf(1) = hQ(1) \in h\mathcal{O}_{\mathfrak{X}, \mathbb{Q}} \subseteq \mathcal{I}_{\mathbb{Q}}.$$

In this way, the right module structure of $\mathcal{O}_{\mathfrak{X}, \mathbb{Q}, \lambda}$ over $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger = (\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger)^{opp}$ makes

$\mathcal{O}_{\mathfrak{X},\mathbb{Q},\lambda}/\mathcal{I}_{\mathbb{Q}}\mathcal{O}_{\mathfrak{X},\mathbb{Q},\lambda}$ a right module over the ring

$$\mathcal{N}_{\mathcal{D}_{\mathfrak{X},\mathbb{Q},2\rho-\lambda}^{\dagger}}(\mathcal{I}\mathcal{D}_{\mathfrak{X},\mathbb{Q},2\rho-\lambda}^{\dagger})/\mathcal{I}\mathcal{D}_{\mathfrak{X},\mathbb{Q},2\rho-\lambda}^{\dagger}.$$

This implies that $\mathcal{O}_{\mathfrak{Y},\mathbb{Q},\lambda} = i^{-1}(\mathcal{O}_{\mathfrak{X},\mathbb{Q},\lambda}/\mathcal{I}_{\mathbb{Q}}\mathcal{O}_{\mathfrak{X},\mathbb{Q},\lambda})$ is a left module over $\mathcal{D}_{\mathfrak{Y},\mathbb{Q},\lambda}^{\dagger}$, as claimed.

For (ii) we assume that the special fibre \mathfrak{Y}_s and the rigid-analytic generic fibre \mathfrak{Y}_K of \mathfrak{Y} are connected. Let $\mathcal{I} \subseteq \mathcal{O}_{\mathfrak{Y},\mathbb{Q},\lambda}$ be a $\mathcal{D}_{\mathfrak{Y},\mathbb{Q},\lambda}^{\dagger}$ -stable submodule. Then \mathcal{I} is a coherent $\mathcal{O}_{\mathfrak{Y},\mathbb{Q}}$ -module. We claim that the intersection $\text{Supp}(\mathcal{I}) \cap \text{Supp}(\mathcal{O}_{\mathfrak{Y},\mathbb{Q},\lambda}/\mathcal{I})$ is empty. Indeed, let us assume for a contradiction that

$$y \in \text{Supp}(\mathcal{I}) \cap \text{Supp}(\mathcal{O}_{\mathfrak{Y},\mathbb{Q},\lambda}/\mathcal{I}).$$

Choose an open affine rig-connected $\mathfrak{U} \subset \mathfrak{Y}$ containing the point y . Making \mathfrak{U} smaller if necessary, we may undo the twist and assume that the restrictions of $\mathcal{O}_{\mathfrak{Y},\mathbb{Q},\lambda}$ and $\mathcal{D}_{\mathfrak{Y},\mathbb{Q},\lambda}^{\dagger}$ to \mathfrak{U} are isomorphic to $\mathcal{O}_{\mathfrak{U},\mathbb{Q}}$ and $\mathcal{D}_{\mathfrak{U},\mathbb{Q}}^{\dagger}$ respectively. But then the sheaf $\mathcal{I}|_{\mathfrak{U}}$ corresponds to a nonzero and proper $\mathcal{D}_{\mathfrak{U},\mathbb{Q}}^{\dagger}$ -submodule of $\mathcal{O}_{\mathfrak{U},\mathbb{Q}}$. This is in contradiction to the fact that $\mathcal{O}_{\mathfrak{U},\mathbb{Q}}$ is a simple $\mathcal{D}_{\mathfrak{U},\mathbb{Q}}^{\dagger}$ -module, cf. [14, Prop. 2.3.6]. So $\text{Supp}(\mathcal{I}) \cap \text{Supp}(\mathcal{O}_{\mathfrak{Y},\mathbb{Q},\lambda}/\mathcal{I})$ is empty. Since $\text{Supp}(\mathcal{I})$ and $\text{Supp}(\mathcal{O}_{\mathfrak{Y},\mathbb{Q},\lambda}/\mathcal{I})$ are closed subsets of \mathfrak{Y}_s and their union equals \mathfrak{Y}_s , the connectedness of \mathfrak{Y}_s implies then that one of them is empty, thus either $\mathcal{I} = 0$ or $\mathcal{I} = \mathcal{O}_{\mathfrak{Y},\mathbb{Q},\lambda}$. So $\mathcal{O}_{\mathfrak{Y},\mathbb{Q},\lambda}$ is a simple left $\mathcal{D}_{\mathfrak{Y},\mathbb{Q},\lambda}^{\dagger}$ -module, as claimed. \square

5.3. Geometric construction of simple modules

We keep the notation. In particular, $\lambda : D^{\dagger}(\mathcal{T})_{\mathbb{Q}} \rightarrow K$ is a character giving rise to the twisted sheaf $\mathcal{D}_{\mathfrak{X},\mathbb{Q},\lambda}^{\dagger}$ on the flag variety \mathfrak{X} . Let $i : \mathfrak{Y} \rightarrow \mathfrak{X}$ be a smooth closed formal subscheme with twisted sheaf $\mathcal{D}_{\mathfrak{Y},\mathbb{Q},\lambda}^{\dagger}$.

Let $\theta : Z(\mathfrak{g}) \rightarrow K$ be a character of the center $Z(\mathfrak{g})$ of $U(\mathfrak{g})$, which corresponds to the weight of \mathfrak{t} induced by λ under the classical Harish-Chandra homomorphism. We let

$$D^{\dagger}(\mathcal{G})_{\mathbb{Q},\theta} := D^{\dagger}(\mathcal{G})_{\mathbb{Q}} \otimes_{Z(\mathfrak{g}),\theta} K$$

be the corresponding central reduction. We recall the localization theorem for left $D^{\dagger}(\mathcal{G})_{\mathbb{Q},\theta}$ -modules. Recall that $\rho = \frac{1}{2} \sum_{\Phi^+} \alpha$.

THEOREM 5.4. —

- (a) *Suppose that $\lambda + \rho$ is dominant and regular (as a weight of \mathfrak{t}). The global section functor induces an equivalence of categories between coherent left $\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger$ -modules and coherent left $H^0(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger)$ -modules.*
- (b) *The \mathcal{G} -action on \mathfrak{X} induces an algebra isomorphism*

$$D^\dagger(\mathcal{G})_{\mathbb{Q}, \theta} \xrightarrow{\simeq} H^0(\mathfrak{X}, \mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger).$$

Proof. — If λ lifts to an algebraic character of T , then this summarizes the main results of [13] and [17]. The case of a general character is the main result of [18]. \square

We come to the main application of our results.

THEOREM 5.5. —

- (i) *There is an equivalence of categories*

$$i_{+, \lambda} : \text{Coh}(\mathcal{D}_{\mathfrak{Y}, \mathbb{Q}, \lambda}^\dagger) \xrightarrow{\simeq} \text{Coh}^{\mathfrak{Y}}(\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger)$$

preserving simple objects on both sides. If $\lambda \in X(T)$, let $\mathcal{B}_{\mathfrak{Y}|\mathfrak{X}, \lambda} := i_{+, \lambda} \mathcal{O}_{\mathfrak{Y}, \mathbb{Q}, \lambda}$.

- (ii) *Let $\lambda + \rho$ be dominant and regular and let \mathfrak{Y} have connected special and generic fibre. If $\lambda \in X(T)$, then $H^0(\mathfrak{X}, \mathcal{B}_{\mathfrak{Y}|\mathfrak{X}, \lambda})$ is a simple $D^\dagger(\mathcal{G})_{\mathbb{Q}, \theta}$ -module.*
- (iii) *Let $\lambda + \rho$ be dominant and regular and let $\lambda \in X(T)$. Let $\mathfrak{Y}, \mathfrak{Y}'$ be two smooth closed formal subschemes of \mathfrak{X} with connected special and generic fibre. If $H^0(\mathfrak{X}, \mathcal{B}_{\mathfrak{Y}|\mathfrak{X}, \lambda}) \simeq H^0(\mathfrak{X}, \mathcal{B}_{\mathfrak{Y}'|\mathfrak{X}, \lambda})$ as $D^\dagger(\mathcal{G})_{\mathbb{Q}, \theta}$ -modules, then $\mathfrak{Y}_s = \mathfrak{Y}'_s$.*

Proof. — The point (i) follows from Proposition 5.2 and the Berthelot–Kashiwara theorem for right modules over $(\mathcal{D}_{\mathfrak{X}, \mathbb{Q}, \lambda}^\dagger)^{opp} = \mathcal{D}_{\mathfrak{X}, \mathbb{Q}, 2\rho - \lambda}^\dagger$, cf. Theorem 4.16. The points (ii) and (iii) follow from Proposition 5.3 together with the localisation theorem. \square

Remark. — In the setting of (iii), an isomorphism $H^0(\mathfrak{X}, \mathcal{B}_{\mathfrak{Y}|\mathfrak{X}, \lambda}) \simeq H^0(\mathfrak{X}, \mathcal{B}_{\mathfrak{Y}'|\mathfrak{X}, \lambda})$ does *not* in general imply an equality of \mathfrak{Y} and \mathfrak{Y}' as closed formal subschemes of \mathfrak{X} , even in the case $\lambda = 0$.

Bibliography

- [1] A. BEILINSON & J. BERNSTEIN, “Localisation de \mathfrak{g} -modules”, *C. R. Math. Acad. Sci. Paris* **292** (1981), no. 1, p. 15-18.
- [2] P. BERTHELOT, “ \mathcal{D} -modules arithmétiques I. Opérateurs différentiels de niveau fini”, *Ann. Sci. Éc. Norm. Supér.* **29** (1996), no. 2, p. 185-272.

- [3] ———, *\mathcal{D} -modules arithmétiques. II. Descente par Frobenius*, Mém. Soc. Math. Fr., Nouv. Sér., no. 81, Société Mathématique de France, 2000, vi+136 pages.
- [4] ———, “Introduction à la théorie arithmétique des \mathcal{D} -modules”, in *Cohomologies p -adiques et applications arithmétiques, II*, Astérisque, no. 279, Société Mathématique de France, 2002, p. 1-80.
- [5] W. BORHO & J.-L. BRYLINSKI, “Differential operators on homogeneous spaces. II. Relative enveloping algebras”, *Bull. Soc. Math. Fr.* **117** (1989), no. 2, p. 167-210.
- [6] S. BOSCH, W. LÜTKEBOHMERT & M. RAYNAUD, “Formal and rigid geometry. III. The relative maximum principle”, *Math. Ann.* **302** (1995), no. 1, p. 1-29.
- [7] D. CARO, “Systèmes inductifs cohérents de \mathcal{D} -modules arithmétiques logarithmiques, stabilité par opérations cohomologiques”, *Doc. Math.* **21** (2016), p. 1515-1606.
- [8] M. DEMAZURE & P. GABRIEL, *Groupes algébriques. Tome I: Géométrie algébrique, généralités, groupes commutatifs*, Masson & Cie, Éditeur, Paris, 1970, Avec un appendice “Corps de classes local” par Michiel Hazewinkel, xxvi+700 pages.
- [9] A. GROTHENDIECK, “Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV”, *Publ. Math., Inst. Hautes Étud. Sci.* (1967), no. 32, p. 361.
- [10] R. Hotta, K. TAKEUCHI & T. TANISAKI, *D -modules, perverse sheaves, and representation theory*, Progress in Mathematics, vol. 236, Birkhäuser, 2008, translated from the 1995 Japanese edition by Takeuchi, xii+407 pages.
- [11] C. HUYGHE, D. PATEL, T. SCHMIDT & M. STRAUCH, “ \mathcal{D}^\dagger -affinity of formal models of flag varieties”, *Math. Res. Lett.* **26** (2019), no. 6, p. 1677-1745.
- [12] C. HUYGHE & T. SCHMIDT, “Algèbres de distributions et \mathcal{D} -modules arithmétiques”, *Rend. Semin. Mat. Univ. Padova* **139** (2018), p. 1-76.
- [13] ———, “ \mathcal{D} -modules arithmétiques sur la variété de drapeaux”, *J. Reine Angew. Math.* **754** (2019), p. 1-15.
- [14] ———, “Intermediate extensions and crystalline distribution algebras”, *Doc. Math.* **26** (2021), p. 2005-2059.
- [15] M. KASHIWARA, “Representation theory and D -modules on flag varieties”, in *Orbites unipotentes et représentations. III. Orbites et faisceaux pervers*, Astérisque, no. 173-174, 1989, p. 55-109.
- [16] D. MILIČIĆ, DRAGAN, “Algebraic \mathcal{D} -modules and representation theory of semisimple Lie groups”, in *The Penrose transform and analytic cohomology in representation theory*, Contemporary Mathematics, vol. 154, American Mathematical Society, 1993, p. 133-168.
- [17] C. NOOT-HUYGHE, “Un théorème de Beilinson-Bernstein pour les \mathcal{D} -modules arithmétiques”, *Bull. Soc. Math. Fr.* **137** (2009), no. 2, p. 159-183.
- [18] A. SARRAZOLA-ALZATE, “A Beilinson-Bernstein theorem for twisted arithmetic differential operators on the formal flag variety”, *J. Théor. Nombres Bordeaux* **36** (2024), no. 1, p. 1-43.
- [19] THE STACKS PROJECT AUTHORS, “The Stacks project”, 2020, <https://stacks.math.columbia.edu>.