



Annales de la Faculté des Sciences de Toulouse

MATHÉMATIQUES

ANTOINE LEMENANT AND RÉMY MOUGENOT

Boundary regularity for the polyharmonic Dirichlet problem

Tome XXXV, n° 3 (2026), p. 743–783.

<https://doi.org/10.5802/afst.1862>

© les auteurs, 2026.

Les articles des *Annales de la Faculté des Sciences de Toulouse* sont mis à disposition sous la licence Creative Commons Attribution (CC-BY) 4.0
<http://creativecommons.org/licenses/by/4.0/>



Publication membre du centre
Mersenne pour l'édition scientifique ouverte
<http://www.centre-mersenne.org/>
e-ISSN : 2258-7519

Boundary regularity for the polyharmonic Dirichlet problem ^(*)

ANTOINE LEMENANT ⁽¹⁾ AND RÉMY MOUGENOT ⁽²⁾

ABSTRACT. — In this paper we prove that any solution of the m -polyharmonic Poisson equation in a Reifenberg-flat domain with homogeneous Dirichlet boundary condition, is $\mathcal{C}^{m-1,\alpha}$ regular up to the boundary. To achieve this result we extend the Nirenberg method of translations to operators of arbitrary order, and then use some Mosco-convergence tools developed in a previous paper [12].

RÉSUMÉ. — Dans cet article, nous démontrons que toute solution de l'équation de Poisson m -polyharmonique dans un domaine Reifenberg-plat, munie de conditions aux limites de Dirichlet homogènes, est de régularité $\mathcal{C}^{m-1,\alpha}$ jusqu'au bord. Pour obtenir ce résultat, nous étendons la méthode des translations de Nirenberg à des opérateurs d'ordre arbitraire, puis nous utilisons des outils de Mosco-convergence développés dans un article précédent [12].

1. Introduction

This paper is devoted to the regularity up to the boundary of solutions of a general elliptic PDE of order $2m$ with constant coefficients. A prototype is, for instance, the following m -polyharmonic Poisson equation:

$$\begin{cases} (-\Delta)^m u = f & \text{in } \Omega \\ u \in H_0^m(\Omega). \end{cases} \quad (1.1)$$

^(*) Reçu le 25 février 2025, accepté le 6 janvier 2026.

Keywords: Polyharmonic problem, elliptic regularity, Reifenberg-flat domain.

2020 *Mathematics Subject Classification*: 35B65, 49Q99.

⁽¹⁾ Institut Universitaire de France (IUF), Université de Lorraine, CNRS, Institut Elie Cartan de Lorraine, BP 70239 54506 Vandœuvre-lès-Nancy Cedex, France — antoine.lemenant@univ-lorraine.fr

⁽²⁾ Université de Lorraine, CNRS, Institut Elie Cartan de Lorraine, BP 70239, 54506 Vandœuvre-lès-Nancy Cedex, France — remy.mougenot@univ-lorraine.fr

Article proposé par Xavier Tolsa.

For any bounded domain Ω and $f \in L^2(\Omega)$, it is easy to see that there exists a unique solution to the above problem. In the following, we also impose a mild regularity condition on the domain Ω . Specifically, we consider the relatively broad class of Reifenberg-flat domains. This concept encompasses, in particular, Lipschitz domains and has been utilized in numerous previous studies on the regularity of elliptic PDEs, as for instance [5, 6, 7, 8, 13, 14, 15, 16, 17, 18, 19, 20, 23], to mention a few. Let us first introduce some notation.

For every $r > 0$ and $x \in \mathbb{R}^N$, $B(x, r)$ is the open ball of radius r and centered at x , $B_r := B(0, r)$. Moreover for $\lambda \in [-1, 1]$ we will denote by $B^\lambda(x, r)$ the truncated ball of level λ defined by

$$B^\lambda(x, r) := B(x, r) \cap \{y \in \mathbb{R}^N \mid y_N > \lambda r + x_N\}.$$

We also denote $B^+(x, r) := B^0(x, r)$ the upper half-ball centered at x .

DEFINITION 1.1 (Reifenberg flat domain). — *Let $r_0 > 0$ and $\varepsilon_0 \in (0, 1)$. A bounded open set is (ε_0, r_0) -Reifenberg flat if for every $x \in \partial\Omega$ and $r \in [0, r_0]$, there exists a rotation R_x centered at x such that*

$$B^{\varepsilon_0}(x, r) \subset R_x(\Omega) \cap B(x, r) \subset B^{-\varepsilon_0}(x, r).$$

This definition is equivalent to the existence of a hyperplane $\mathcal{P}_x$ containing x such that

$$d_{\mathcal{H}}(\partial\Omega \cap B(x, r), \mathcal{P}_x \cap B(x, r)) \leq \varepsilon_0 r, \quad r \leq r_0,$$

where $d_{\mathcal{H}}$ is the Hausdorff distance defined by

$$d_{\mathcal{H}}(A, B) := \sup_{x \in A} \text{dist}(x, B) + \sup_{y \in B} \text{dist}(y, A),$$

and such that

$$B(x, r) \cap \{y \in \mathbb{R}^N \mid \text{dist}(y, \mathcal{P}_x) > \varepsilon_0 r\}$$

has two connected components, one lying in Ω , the other one lying in $\mathbb{R}^N \setminus \Omega$, see [15, 16, 17, 25].

The purpose of this paper is to investigate the boundary regularity for Problem (1.1) in Reifenberg flat domains. Actually, it applies to more general operators of order $2m$ of the form

$$\mathcal{A} := (-1)^m \sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \partial^\beta \partial^\alpha, \quad (1.2)$$

where $a_{\alpha,\beta} \in \mathbb{R}$ are real coefficients. We assume that $\mathcal{A}$ is symmetric, i.e. $a_{\alpha,\beta} = a_{\beta,\alpha}$, and is elliptic in the sense that there exists $C > 0$ such that for

all vectors $\xi = (\xi_\alpha)_{|\alpha|=m}$ with $\xi_\alpha \in \mathbb{R}$ it holds that

$$\sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \xi_\alpha \xi_\beta \geq C \sum_{|\alpha|=m} |\xi_\alpha|^2. \quad (1.3)$$

Also, observe that a Reifenberg-flat domain is always assumed to be bounded, by definition. Here is our main result.

THEOREM 1.2. — *Let $\alpha \in (0, 1)$ and $q \geq 2$ be such that $mq \geq N$ if $2m < N$, and $q = 2$ otherwise. There exists $\varepsilon_0 \in (0, 1)$ and $r_0 \in (0, 1]$ such that for every (ε_0, r_0) -Reifenberg-flat domain $\Omega \subset \mathbb{R}^N$, for every function $f \in L^q(\Omega)$, if $u \in H_0^m(\Omega)$ is the weak solution of $\mathcal{A}(u) = f$, then $u \in \mathcal{C}^{m-1,\alpha}(\bar{\Omega})$ and*

$$\|u\|_{\mathcal{C}^{m-1,\alpha}(\bar{\Omega})} \leq C \|f\|_{L^q(\Omega)},$$

where $C > 0$ depends on $N, \mathcal{A}, \alpha, \Omega$ and m .

What is somewhat surprising in Theorem 1.2 is that we are able to obtain fairly strong regularity up to the boundary, in the class $\mathcal{C}^{m-1,\alpha}$, although $\partial\Omega$ enjoys only very poor regularity, not better than $\mathcal{C}^{0,\gamma}$. Indeed, a Reifenberg-flat domain is less regular than Lipschitz, it could even have a fractal boundary. The high regularity of solutions in this poor regularity class of domains is possible due to the fact that $u \in H_0^m(\Omega)$ vanishes at the boundary, as well as all its k^{th} derivatives, up to the order $m - 1$, leaving a chance to be more regular up to the boundary, than the boundary itself.

For instance our result immediately implies that in dimension 3 and in the case of the bi-laplacian, if $u \in H_0^2(\Omega)$ and $\Delta^2 u \in L^2(\Omega)$, then $u \in \mathcal{C}^{1,\alpha}(\bar{\Omega})$, while Ω is only Reifenberg-flat.

Notice that for the interior regularity, if $\Delta^m(u) = f$ with $f \in L^q$ then $\Delta(\Delta^{m-1})(u) = f$ so that $\Delta^{m-1}u \in W^{2,q}$ by classical Calderon–Zygmund estimates on the standard Laplacian. Iterating this m times we arrive to the fact that $u \in W^{2m,q}$. Hence, if $mq > N$, the Sobolev embedding theorem implies that $\nabla^m u \in \mathcal{C}^{0,\alpha}$, or in other words $u \in \mathcal{C}^{m,\alpha}$. Of course this argument cannot be applied at the boundary because the Dirichlet boundary condition is not preserved while taking derivatives. But it is interesting to compare to the statement of Theorem 1.2, since up to the boundary $u \in \mathcal{C}^{2m-1,\alpha}$ and not $\mathcal{C}^{2m,\alpha}$, and so we lose one derivative with respect to what we could expect in the interior. This is due to the weak regularity of the boundary that prevents us to go beyond.

Let us furthermore emphasize that our result is sharp. Indeed, for instance for $m = 1$ it is easy to construct a harmonic function with Dirichlet boundary condition on a cone of aperture $\pi + \varepsilon_0$, that is in $\mathcal{C}^{0,\alpha}$ but not Lipschitz (see for instance [17, Remark 18]).

A similar result to our Theorem 1.2 has been partially obtained earlier in [18] in the case $m = 1$, i.e for the standard Laplacian, where the a priori estimates $u \in L^p(\Omega)$ and $f \in L^q(\Omega)$ lead to the $\mathcal{C}^{0,\alpha}(\bar{\Omega})$ regularity, provided that Ω is “flat” enough. Our result includes the one in [18], and is even more general for the case of the standard Laplacian ($m = 1$) because we do not assume any a priori condition on $u \in L^p(\Omega)$ as in [18]. The technique is also different since in [18] a monotonicity formula has been used, which is not available for the polyharmonic operator.

The result in [18] has also been extended for general elliptic operators of order 2 in [20], using some barrier arguments. This technique cannot be applied for operators of higher order since it relies on the maximum principle.

Notice that in the recent preprint [24], a similar regularity result has been obtained for the fractional Laplacian $(-\Delta)^s$. For the particular case $s = 1$, the result in [24] is consistent with the particular case of our result with $m = 1$.

For the parabolic case, an L^p theory has been developed in [8] in a general context of varying BMO coefficients, from which one can probably deduce a similar statement to our main result, after an application of the Sobolev embedding theorem. However, the technique employed in the present paper is completely different from the one in [8], and somewhat simpler.

It is worth mentioning that many works by S. Mayboroda and co-authors are dedicated to the study of the bi-harmonic operator in a general class of wild domains, including in particular Reifenberg-flat domains (see for instance [3, 4, 21, 22]). The context in these works is a bit different since they are interested in the polyharmonic measure and thus focus on the delicate problem $\Delta^2 u = 0$ in Ω with $\nabla u = f$ on the boundary. In the present paper instead, we are looking for $u = 0$ and $\nabla u = 0$ on the boundary with a source $\Delta^2 u = f$ in Ω , which is slightly different.

Our approach, which is completely self-contained in this paper, uses variational techniques that are well adapted for solutions with homogenous boundary conditions, that might not work so well with an inhomogeneous condition. On the other hand, it provides some very precise energy estimates at the boundary in that particular case, that one could probably not obtain so easily with the general tools developed for the polyharmonic measure.

We also believe that our result could be extended to more general operators of divergence type with Hölder continuous coefficients, but we do not pursue this generalization for the sake of simplicity, restricting ourselves to constant coefficients.

Ideas of proof and structure of the paper

Let us now describe the ideas behind the proof of Theorem 1.2. The main point is to arrive at a decay behavior of the energy at the boundary for a polyharmonic function of the following type: there exist $b, C, r_0 > 0$ such that for all $x \in \partial\Omega$ and $r \leq R \leq r_0$, (in the sequel, a function in $H_0^m(\Omega)$ is considered as being extended by 0 outside Ω)

$$\int_{B(x,r)} |\nabla^m u|^2 \, dx \leq C \left(\frac{r}{R} \right)^{N-b} \int_{B(x,R)} |\nabla^m u|^2 \, dx, \quad r \in (0, R). \quad (1.4)$$

If (1.4) holds, then one may say that $u \in \mathcal{C}^{m-1,\alpha}(\bar{\Omega})$ by standard Campanato theory.

Now to obtain (1.4) at a boundary point of a Reifenberg-flat domain, the first step is to prove that it holds true when the boundary is a hyperplane. Once this is established, then we can argue by compactness and obtain that it still holds true when the boundary is close enough to a hyperplane. This is done in Proposition 1.4, which uses in particular the Mosco-convergence theorem for higher order operators contained in [12]. Since a Reifenberg flat domain is ε_0 -close to a hyperplane at any small scale, we can then conclude that the same decay holds for a solution in a Reifenberg-flat domain, leading to our main $\mathcal{C}^{m-1,\alpha}$ result.

Therefore, a fairly big part of this paper is actually devoted to the proof of the energy decay property (1.4), in the case of a perfect flat boundary, i.e., when $\partial\Omega$ coincides with a hyperplane.

For the standard regularity theory for operators of order $2m$, one usually quote [1, 2], based on Fourier analysis. However, we could not find in the literature the exact statement that is needed for our purposes. Instead, we provide in this paper a complete and self-contained proof, based on the Nirenberg translation method, that allows us to control precisely all the constants.

The interesting fact is that we have to adapt this method in a non trivial way for higher order operators, compared to what is commonly known for the Laplace operator. To our knowledge, we believe this argument to be new and interesting on its own. Indeed, let us explain the difficulty.

For the recall, a usual way to treat the problem of boundary regularity in the flat case, for instance when $\Omega = \{x_N > 0\}$, is to consider a direction of translation $h \in \mathbb{R}^N \setminus \{0\}$ and then introduce the translated function

$$D_h u(x) = \frac{u(x+h) - u(x)}{|h|}, \quad x \in \Omega.$$

The main idea is the following: up to localizing with a cut-off function and provided that h goes in a horizontal direction (i.e., for $h = e_i$ and $1 \leq i \leq N - 1$), the function $D_h u$ is an admissible test function in $H_0^m(\{e_N > 0\})$ in the weak formulation of the problem. For the polyharmonic problem this leads to an a priori bound of the type

$$\|D_h \nabla^m u\|_2 \leq C.$$

Since C is uniform in h , we easily conclude by taking $h \rightarrow 0$ that actually u admits one more derivative in all the horizontal directions. We can continue inductively and actually prove that all the derivatives of the form $\partial^\alpha \nabla^m u$ are in L^2 , where α are multi-indices involving only horizontal directions. Then we only miss the vertical direction, in which the translation is not admissible as a test function.

For the standard Laplacian, the usual trick works as follows: we can recover the vertical derivatives by use of the operator. Indeed, if $-\Delta u = f$ then

$$-\partial_N^2 u = f + \sum_{i=1}^{N-1} \partial_i^2 u,$$

and this is how we obtain that $\partial_N^2 u \in L^2$ because we know that all the $\partial_i^2 u$ for $1 \leq i \leq N - 1$ are in L^2 . We then conclude that $u \in H^2$, which is a gain of regularity compared to the original information that $u \in H^1$.

For higher order operators $\mathcal{A}$, we cannot recover the estimate on $\partial_N^{2m} u$ so easily because we need to pass from $\nabla^m u \in L^2$ to $\nabla^{2m} u \in L^2$: in other words we have to win m derivatives, and not only one. Moreover, the operator is more complicated and involves multiples derivatives of different order.

As a result, our proof works by induction on the order of the operator. We first establish the regularity theorem for $m = 1$, which is the case for instance for the standard Laplace operator. Then assuming that Theorem 1.2 is true for operators $\mathcal{A}$ of order $2(m - 1)$, we prove that it holds true for operators of order $2m$. The main tool is to write a general operator $\mathcal{A}$ of order $2m$ as the sum (see Lemma 2.1):

$$\mathcal{A}(u) = \mathcal{B}(u) - \partial_N \mathcal{C}(\partial_N u), \quad (1.5)$$

where $\mathcal{B}(u)$ contains at most m -derivatives in the e_N direction, so that it belongs to L^2 , and $\mathcal{C}$ is an elliptic operator of order $2(m - 1)$, on which we can apply our induction hypothesis. Indeed, since $\mathcal{A}(u) \in L^2$ by assumption, we deduce that $\partial_N \mathcal{C}(\partial_N u) \in L^2$. Then by use of a certain Poincaré inequality, this actually yields $\mathcal{C}(\partial_N u) \in L^2$, and thus $\partial_N u \in H^{2(m-1)}$ thanks to our induction hypothesis (i.e. using that Theorem 1.2 is true for operators of order $2(m - 1)$). All this implies that $u \in H^{2m-1}$.

Now we are still missing one last derivative because we need to achieve H^{2m} instead of H^{2m-1} . This will be done by use of the operator, for a last time again. Indeed, at this stage of the proof, we have a control on all the derivatives of the form $\partial^\alpha u$ with $|\alpha| = 2m$, provided that $\alpha_N \leq 2m - 1$. In other words, to conclude that $u \in H^{2m}$ we only miss the last derivative $\partial_N^{2m} u$. But this term appears in $\mathcal{A}$ only once. Namely, we have

$$\mathcal{A} = (-1)^m a_{m e_N, m e_N} \partial_N^{2m} + \mathcal{E} \quad (1.6)$$

where $\mathcal{E}$ contains at most $2m - 1$ derivatives in the e_N direction. This is how we get $\partial_N^{2m} u \in L^2$, and finally conclude that $u \in H^{2m}$, as desired.

All this leads to the regularity result that is contained in Theorem 4.1, that we state here in the introduction to enlight what we have obtained with this method.

THEOREM 1.3 (Regularity with flat boundary). — *Let $m \in \mathbb{N}^*$. Let $\mathcal{A}$ be an operator of order $2m$ of the form (1.2) and satisfying the ellipticity condition (1.3). Let $u \in H^m(B(0, 1))$ be a weak solution of $\mathcal{A}u = f$ in $B^+(0, 1)$ with $f \in H^\ell(B^+(0, 1))$, and such that $u = 0$ on $B(0, 1) \cap \{x_N < 0\}$. Then $u \in H^{2m+\ell}(B^+(0, 1/2))$ and*

$$\|u\|_{H^{2m+\ell}(B^+(0, \frac{1}{2}))} \leq C (\|\nabla^m u\|_{L^2(B^+(0, 1))} + \|f\|_{H^\ell(B^+(0, 1))}),$$

where $C > 0$ is a constant that depends on N , ℓ , m , $\max_{\alpha, \beta} |a_{\alpha, \beta}|$, and the ellipticity constant that appears in (1.3).

We also have written the interior case, analogous to Theorem 1.3 (see Theorem 3.1). As a matter of fact, by reasoning with local charts, our paper provides a self-contained and full regularity theory for the Dirichlet problem of $2m$ -order in a smooth domain.

Observe that our method really needs to work with general operators of the form (1.2), and does not simplify for the special case of $(-\Delta)^m$. Indeed, while performing the induction argument on $(-\Delta)^m$ we would need to decompose it as the sum (1.5), in which an operator of lower order appears, which is not $(-\Delta)^{m-1}$, but more general, as in (1.2).

Then in Section 5 we use this regularity result to establish a decay property of the type (1.4) for a polyharmonic function at a boundary point of a Reifenberg-flat domain. This is obtained by use of a compactness argument (see Proposition 5.1). Once this is done, we can use a “polyharmonic” replacement argument to derive a similar decay property for a solution with a source term. This is done in Proposition 5.2, that leads to the proof of our main result, Theorem 1.2.

2. Preliminaries

As usual we denote by $H^m(\Omega)$ the Sobolev space endowed with the norm

$$\|u\|_{H^m(\Omega)}^2 = \sum_{|\alpha| \leq m} \|\partial^\alpha u\|_2^2$$

and we denote by $H_0^m(\Omega)$ the closure of C^∞ function with compact support in Ω . Since a Reifenberg-flat domain enjoys the so-called corkscrew condition, it follows that

$$H_0^m(\Omega) = \{u \in H^m(\mathbb{R}^N) \text{ s.t. } u = 0 \text{ a.e. on } \mathbb{R}^N \setminus \Omega\}. \quad (2.1)$$

This is actually [12, Corollary 10], and we refer to [12] for more details. As a consequence, throughout the paper we will consider $u \in H_0^1(\Omega)$ as being a function in $H^m(\mathbb{R}^N)$, extended by 0 on Ω^c .

We will also use (2.1) implicitly while dealing with a mixed boundary value problem in a half ball. For instance in Section 4 we will consider $u \in H^m(B(0, 1))$ such that $u = 0$ a.e. on $B(0, 1) \cap \{x_N < 0\}$. It is known (see for instance [12]) that such a function can be approximated by $v_n \rightarrow u$ in $H^m(B(0, 1))$ such that $\text{supp}(v_n) \subset B^+(0, 1)$ for all n . This fact will also be used in the proof of Proposition 5.1.

2.1. Multi-indices and the Leibniz formula

A multi-index is a vector $\alpha \in \mathbb{N}^N$. The length of α is denoted by $|\alpha|$ which is the sum of all the coefficients α_i . The partial derivative ∂^α means that we take all derivatives $\partial_{x_i}^{(\alpha_i)}$. We denote by $\alpha!$ the number $\alpha_1! \alpha_2! \dots \alpha_N!$ and we say that $\alpha \leq \beta$ when $\alpha_i \leq \beta_i$ for all $1 \leq i \leq N$.

If u, v are two smooth functions then the Leibniz formula states that

$$\partial^\alpha(uv) = \sum_{\beta \leq \alpha} \frac{\alpha!}{\alpha!(\alpha - \beta)!} \partial^\beta u \partial^{\alpha - \beta} v. \quad (2.2)$$

2.2. Elliptic operators of order $2m$

Let $m \in \mathbb{N}^*$. We consider a general operator $\mathcal{A}$ of order $2m$ of the form

$$\mathcal{A} := (-1)^m \sum_{|\alpha|=|\beta|=m} \partial^\beta (a_{\alpha,\beta} \partial^\alpha) = (-1)^m \sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \partial^\beta \partial^\alpha, \quad (2.3)$$

where $a_{\alpha,\beta} \in \mathbb{R}$ are real coefficients. We assume that $\mathcal{A}$ is symmetric, i.e., $a_{\alpha,\beta} = a_{\beta,\alpha}$ for all α, β , and is elliptic in the sense that there exists $C_A > 0$ such that for all vectors $\xi = (\xi_\alpha)_{|\alpha|=m}$ with $\xi_\alpha \in \mathbb{R}$ there holds,

$$C_A \sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \xi_\alpha \xi_\beta \geq \sum_{|\alpha|=m} |\xi_\alpha|^2. \quad (2.4)$$

2.3. Weak Formulation

For $f \in L^2(\Omega)$, an application of the Lax–Milgram theorem yields existence and uniqueness for the problem

$$\begin{cases} \mathcal{A}u = f & \text{in } \Omega \\ u \in H_0^m(\Omega) \end{cases} \quad (2.5)$$

The weak formulation of this problem is

$$\int_{\Omega} \sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \partial^\alpha u \partial^\beta \varphi \, dx = \int_{\Omega} \varphi f \, dx \quad \text{for all } \varphi \in H_0^m(\Omega). \quad (2.6)$$

2.4. Matrix formulation

For $u \in H^m(\Omega)$ we denote by $\nabla^m u$ the vector of all derivatives of order m , i.e. $\nabla^m u = (\partial^\alpha u)_{|\alpha|=m}$. Then the weak formulation (2.6) can also be written as follows,

$$\int_{\Omega} A \nabla^m u \cdot \nabla^m \varphi \, dx = \int_{\Omega} \varphi f \, dx \quad \text{for all } \varphi \in H_0^m(\Omega), \quad (2.7)$$

with $A = (a_{\alpha,\beta})$.

2.5. Polyharmonic operators

In particular, our class of operators contains the polyharmonic operator $(-\Delta)^m$ (see [10]) by taking the particular case $a_{\alpha,\beta} = \frac{m!}{\alpha!} \delta_{\alpha,\beta}$ where $\delta_{\alpha,\beta} = 1$ if $\alpha = \beta$ and 0 otherwise. In this case,

$$\sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \xi_\alpha \xi_\beta = \sum_{|\alpha|=m} \frac{m!}{\alpha!} |\xi_\alpha|^2,$$

and therefore it is clear that this operator satisfies the ellipticity condition (2.4).

Even though the following fact will not be used in this paper, it is worth mentioning that by use of a standard integration by parts valid in $H_0^m(\Omega)$, one can prove that the associated bilinear form that appears in the weak formulation for $(-\Delta)^m$ becomes

$$\begin{aligned} \int_{\Omega} \sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \partial^\alpha u \partial^\beta \varphi \, dx \\ = \begin{cases} \int_{\Omega} \Delta^{\frac{m}{2}} u \Delta^{\frac{m}{2}} \varphi \, dx & \text{if } m \text{ is even,} \\ \int_{\Omega} \nabla \Delta^{\frac{m-1}{2}} u \cdot \nabla \Delta^{\frac{m-1}{2}} \varphi \, dx & \text{if } m \text{ is odd.} \end{cases} \end{aligned}$$

2.6. Decomposition of Elliptic operators

In the sequel we will need the following Lemma, which is one of the key step in our proof.

LEMMA 2.1. — *Let $\mathcal{A}$ be an operator of order $2m$ of the form (2.3) that satisfies the ellipticity condition (2.4). Then $\mathcal{A}$ can be split in two parts, $\mathcal{A} = \mathcal{B} + \mathcal{C}$, as follows*

$$\begin{aligned} \mathcal{A} &:= (-1)^m \sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \partial^\beta \partial^\alpha \\ &= (-1)^m \underbrace{\sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N=0 \text{ or } \beta_N=0}} \partial^\beta (a_{\alpha,\beta} \partial^\alpha)}_{\mathcal{B}} + (-1)^m \underbrace{\sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N>0 \text{ and } \beta_N>0}} \partial^\beta (a_{\alpha,\beta} \partial^\alpha)}_{\mathcal{C}}, \end{aligned}$$

in such a way that

$$\mathcal{C} := (-1)^m \sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N>0 \text{ and } \beta_N>0}} \partial^\beta (a_{\alpha,\beta} \partial^\alpha) = \mathcal{D} \circ (-\partial_N^2),$$

where $\mathcal{D}$ is elliptic, as an operator of order $2m - 2$.

Proof. — By definition of $\mathcal{C}$ we can factorize by ∂_N^2 as follows:

$$\begin{aligned} \mathcal{C} &:= (-1)^m \sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N>0 \text{ and } \beta_N>0}} a_{\alpha,\beta} \partial^\beta \partial^\alpha \\ &= (-1)^{m-1} \sum_{|\alpha|=|\beta|=m-1} a_{\alpha+e_N, \beta+e_N} \partial^\beta \partial^\alpha (-\partial_N^2), \end{aligned}$$

and therefore the operator $\mathcal{D}$ of the statement of Lemma 2.1 is given by

$$\mathcal{D} := (-1)^{m-1} \sum_{|\alpha|=|\beta|=m-1} a_{\alpha+e_N, \beta+e_N} \partial^\beta \partial^\alpha.$$

We have to check that this operator is elliptic, as an operator of order $2m-2$. For that purpose we will use the ellipticity of $\mathcal{A}$ on suitably chosen vectors ξ . More precisely, let $\eta = (\eta_\alpha)_{|\alpha|=m-1}$ be any given vector indexed over all multi-indices of length $m-1$. We complete the vector η by defining a vector ξ indexed by multi-indices of length m , by writing now when $|\alpha| = m$,

$$\xi_\alpha := \begin{cases} \eta_{\alpha-e_N} & \text{if } \alpha_N > 0 \\ 0 & \text{if } \alpha_N = 0. \end{cases}$$

By testing the ellipticity of $\mathcal{A}$ with this vector ξ we get

$$\begin{aligned} \sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \xi_\beta \xi_\alpha &= \underbrace{\sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N=0 \text{ or } \beta_N=0}} a_{\alpha,\beta} \xi_\alpha \xi_\beta}_{=0} \\ &+ \sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N>0 \text{ and } \beta_N>0}} a_{\alpha,\beta} \xi_\alpha \xi_\beta \geq C_A^{-1} \sum_{|\alpha|=m} |\xi_\alpha|^2, \end{aligned}$$

which directly gives

$$\sum_{|\alpha|=|\beta|=m-1} a_{\alpha+e_N, \beta+e_N} \eta_\alpha \eta_\beta \geq C_A^{-1} \sum_{|\alpha|=m-1} |\eta_\alpha|^2,$$

and this proves that $\mathcal{D}$ is elliptic, as an operator of order $2m-2$. $\square$

2.7. Poincaré type inequalities

The following simple lemma will be needed in the proof of Theorem 3.1. The proof is elementary, by integrating on vertical rays.

LEMMA 2.2. — *For $r > 0$ we denote by $Q_r \subset \mathbb{R}^N$ the cube centered at the origin of side r , i.e.,*

$$Q_r := \{x \in \mathbb{R}^N \mid \forall 1 \leq i \leq N, |x_i| \leq r\}.$$

For $\lambda \in (0, 1)$ we also introduce

$$Q_r^\lambda := Q_r \cap \{x_N > \lambda r\}.$$

Assume that $v \in L^2(Q_r^\lambda)$ for all $\lambda \in (0, 1)$ and that furthermore v satisfies

$$\|\partial_N v\|_{L^2(Q_r)} < +\infty.$$

Then $v \in L^2(Q_r)$ and for all $\lambda \in (0, 3/4)$ it holds:

$$\|v\|_{L^2(Q_r)} \leq 4\|v\|_{L^2(Q_r^\lambda)} + 3r\|\partial_N v\|_{L^2(Q_r)}. \quad (2.8)$$

Proof. — We first assume that $v \in C^\infty(\overline{Q_r})$. We will use the notation $x = (x', x_N)$ for a point in $\mathbb{R}^N$. Then integrating along the e_N direction yields, for every $(x', t) \in Q_r$,

$$v(x', t) = v(x', s) + \int_s^t \partial_N v(x', z) dz.$$

We then integrate over $\lambda r \leq s \leq r$,

$$(1 - \lambda)rv(x', t) = \int_{\lambda r}^r v(x', s) ds + \int_{\lambda r}^r \int_s^t \partial_N v(x', z) dz ds.$$

Taking the square and using that $(a + b)^2 \leq 2(a^2 + b^2)$ it holds

$$(1 - \lambda)^2 r^2 v(x', t)^2 \leq 2 \left(\int_{\lambda r}^r v(x', s) ds \right)^2 + 2 \left((1 - \lambda)r \int_{-r}^r |\partial_N v(x', z)| dz \right)^2,$$

which becomes, by Cauchy–Schwarz inequality,

$$(1 - \lambda)^2 r^2 v(x', t)^2 \leq 2(1 - \lambda)r \int_{\lambda r}^r v(x', s)^2 ds + 4(1 - \lambda)^2 r^3 \int_{-r}^r (\partial_N v(x', z))^2 dz,$$

or differently,

$$(1 - \lambda)rv(x', t)^2 \leq 2 \int_{\lambda r}^r v(x', s)^2 ds + 4(1 - \lambda)r^2 \int_{-r}^r (\partial_N v(x', z))^2 dz.$$

We then integrate over $x' \in Q_r'$ (which is the cube in $\mathbb{R}^{N-1}$), arriving to

$$(1 - \lambda)r \int_{Q_r'} v(x', t)^2 dx' \leq 2 \int_{Q_r^\lambda} v(x)^2 dx + 4(1 - \lambda)r^2 \int_{Q_r} (\partial_N v(x))^2 dx.$$

A last integration over $-r \leq t \leq r$ finally yields

$$(1 - \lambda)r \int_{Q_r} v(x)^2 dx \leq 4r \int_{Q_r^\lambda} v(x)^2 dx + 8(1 - \lambda)r^3 \int_{Q_r} (\partial_N v(x))^2 dx,$$

or equivalently,

$$\int_{Q_r} v(x)^2 dx \leq \frac{4}{1 - \lambda} \int_{Q_r^\lambda} v(x)^2 dx + 8r^2 \int_{Q_r} (\partial_N v(x))^2 dx,$$

which proves the inequality (2.8) in the case when $v \in C^\infty(\overline{Q_r})$, because $\lambda \leq \frac{3}{4}$ by assumption.

Now let v be as in the statement, i.e. $v \in L^2(Q_\lambda)$ for all $\lambda \in (0, 1)$ and that furthermore $\|\partial_N v\|_{L^2(Q_r)} < +\infty$. We argue by approximation: for $\varepsilon > 0$, we let $v_\varepsilon := v * \rho_\varepsilon$ where ρ_ε is a standard mollifier. Then since $v \in L^2(Q_\lambda)$ for all $\lambda \in (0, 1)$, we know that $v_\varepsilon \rightarrow v$ in $L^2(Q_\lambda)$ for all $\lambda \in (0, 1)$ and by

extracting a diagonal sequence we can assume that v_ε converges to v a.e. on Q_r . Then by applying (2.8) to v_ε , and passing to the limit $\varepsilon \rightarrow 0$ we get from Fatou's Lemma that

$$\|v\|_{L^2(Q_r)} \leq \liminf_{\varepsilon \rightarrow 0} \|v_\varepsilon\|_{L^2(Q_r)} \leq 4\|v\|_{L^2(Q_r^\lambda)} + 3r\|\partial_N v\|_{L^2(Q_r)},$$

which ends the proof of Lemma 2.2. $\square$

We will also need the following standard version of the Poincaré inequality, that follows from instance from [26, Corollary 4.5.2, p. 195].

LEMMA 2.3. — *If $v \in H^m(B(0,1))$ is such that $v = 0$ a.e. in $B(0,1) \cap \{x_N < 0\}$, then*

$$\|v\|_{H^m(B(0,1))} \leq C\|\nabla^m v\|_{L^2(B(0,1))},$$

where $C > 0$ depends only on the dimension N .

2.8. Finite difference tools

We will prove the theorem using Nirenberg's "translation method". A good reference for this section is for instance [9, Section 5.8.2], but for the convenience of the reader, and since our statement are not exactly the same, we write the proofs with all details. For $h \in \mathbb{R}^N$, and $\varphi : \mathbb{R}^N \rightarrow \mathbb{R}$, we denote

$$\tau_h(\varphi)(x) = \varphi(x + h)$$

and then

$$D_h(\varphi) = \frac{\varphi(x + h) - \varphi(x)}{|h|} = \frac{\tau_h(\varphi)(x) - \varphi(x)}{|h|}.$$

The main ingredient is that a uniform L^2 -control on $D_h(\varphi)$ (with respect to h), leads to the conclusion that $\nabla \varphi \in L^2$, as stated in the following proposition.

PROPOSITION 2.4. — *Let $\omega \subset \Omega \subset \mathbb{R}^N$ be two open sets such that $\bar{\omega} \subset \Omega$ and let $u \in L^2(\Omega)$. Then the following two properties hold:*

- (1) *If $u \in H^1(\Omega)$ then for every $|\varepsilon| < \text{dist}(\omega, \Omega^c)$,*

$$\|D_{\varepsilon e_i}(u)\|_{L^2(\omega)} \leq \|\partial_i u\|_{L^2(\Omega)}.$$

- (2) *Conversely, if $u \in L^2(\Omega)$ and if there exists $C_0 > 0$ such that for all $|\varepsilon| < \text{dist}(\omega, \Omega^c)$, it holds that $\|D_{\varepsilon e_i}(u)\|_{L^2(\omega)} \leq C_0$, then $\partial_i u \in L^2(\omega)$ and we have*

$$\|\partial_i u\|_{L^2(\omega)} \leq C_0. \tag{2.9}$$

Proof. — We start by assuming that $u \in C^\infty(\Omega) \cap H^1(\Omega)$. Let ε be such that $|\varepsilon| \leq \text{dist}(\omega, \Omega^c)$. We write

$$u(x + \varepsilon e_i) - u(x) = \varepsilon \int_0^1 \partial_i u(x + t\varepsilon e_i) dt.$$

Using the Cauchy–Schwarz inequality and Fubini’s theorem, it follows that

$$\int_\omega |u(x + \varepsilon e_i) - u(x)|^2 dx \leq \varepsilon^2 \int_0^1 \int_\omega |\partial_i u(x + t\varepsilon e_i)|^2 dx dt.$$

By the change of variables $y = x + t\varepsilon e_i$, we find

$$\int_\omega |u(x + \varepsilon e_i) - u(x)|^2 dx \leq |\varepsilon|^2 \int_0^1 \int_{\omega + t\varepsilon e_i} |\partial_i u(y)|^2 dy dt.$$

If $|\varepsilon| \leq \text{dist}(\omega, \Omega^c)$, then $\omega + t\varepsilon e_i \subset \Omega$, so finally

$$\|D_{\varepsilon e_i}(u)\|_{L^2(\omega)} \leq \|\partial_i u\|_{L^2(\Omega)}.$$

Now if $u \in H^1(\Omega)$, the result follows by approximation with C^∞ functions, which concludes the proof of assertion (1).

Let us now prove the converse. We take $0 < \varepsilon < \text{dist}(\omega, \Omega^c)$ and we define $g_\varepsilon := D_{\varepsilon e_i} u$, so that the hypothesis shows that the family $(g_\varepsilon)_{\varepsilon > 0}$ is bounded in $L^2(\omega)$. Therefore, there exists a subsequence such that $g_\varepsilon \rightarrow g^{(i)}$ weakly in $L^2(\omega)$. In particular

$$\|g^{(i)}\|_{L^2(\omega)} \leq \liminf \|D_{\varepsilon e_i} u\|_{L^2(\omega)} \leq C.$$

Let $\varphi \in C_c^\infty(\omega)$. For $0 < \varepsilon < \text{dist}(\omega, \Omega^c)$, we have

$$\begin{aligned} \int_\Omega (D_{\varepsilon e_i} u(x)) \varphi(x) dx &= \int_\Omega \frac{u(x + \varepsilon e_i) - u(x)}{\varepsilon} \varphi(x) dx \\ &= - \int_\Omega u(y) \frac{\varphi(y) - \varphi(y - \varepsilon e_i)}{\varepsilon} dy \\ &= \int_\Omega u(y) D_{-\varepsilon e_i} \varphi(y) dy. \end{aligned}$$

Moreover, since $D_{-\varepsilon e_i} \varphi(y) \rightarrow \partial_i \varphi(y)$ for all y , by the dominated convergence theorem, we deduce that

$$\int_\omega g^{(i)} \varphi dx = \int_\omega u \partial_i \varphi dx,$$

which shows that $\partial_i u \in L^2(\omega)$ with $\partial_i u = g^{(i)}$ and

$$\int_\omega |\partial_i u|^2 dx \leq C_0,$$

where $C_0 > 0$ is the constant from the statement. □

By iterating the above proposition, we can prove the same for higher order derivatives. More precisely, we consider a vector of direction $(h_i)_{1 \leq i \leq N}$ with $h_i \in \mathbb{R}^N$ and the associated discretized operator D_{h_i} defined by

$$D_{h_i} \varphi := \frac{\tau_{h_i} \varphi - \varphi}{|h_i|}.$$

These operators commute two by two: for every $i, j \in \llbracket 1, N \rrbracket$, $|h_i| = |h_j|$, one has

$$D_{h_i} D_{h_j} \varphi = \frac{1}{|h_i|} \tau_{h_i} \left(\frac{\tau_{h_j} \varphi - \varphi}{|h_j|} \right) - \frac{\tau_{h_j} \varphi - \varphi}{|h_i| |h_j|} = \frac{\tau_{h_i} \tau_{h_j} \varphi - \tau_{h_i} \varphi - \tau_{h_j} \varphi + \varphi}{|h_i|^2},$$

but $\tau_{h_i} \tau_{h_j} = \tau_{h_j} \tau_{h_i}$, thus $D_{h_i} D_{h_j} = D_{h_j} D_{h_i}$. Furthermore, they commute with the operators ∂^β , namely $D_{h_i} \partial^\beta \varphi = \partial^\beta D_{h_i} \varphi$.

Then for $\alpha \in \mathbb{N}^N$ and $\varepsilon \in \mathbb{R}$ we define

$$D_\varepsilon^\alpha := \prod_{\substack{i=1 \\ \text{s.t. } \alpha_i \neq 0}}^N \prod_{l=1}^{|\alpha_i|} D_{\varepsilon e_i}. \quad (2.10)$$

Notice that by a simple change of variables, for every $u, v \in L^2(\mathbb{R}^N)$ we have

$$\int_{\mathbb{R}^N} u D_\varepsilon v \, dx = \int_{\mathbb{R}^N} v D_{-\varepsilon} u \, dx,$$

and iterating this property we obtain

$$\int_{\mathbb{R}^N} u D_\varepsilon^\alpha v \, dx = \int_{\mathbb{R}^N} v D_{-\varepsilon}^\alpha u \, dx. \quad (2.11)$$

By applying m -times Proposition 2.4, we arrive at the following direct corollary.

PROPOSITION 2.5. — *Let $\omega \subset \Omega \subset \mathbb{R}^N$ be two open sets such that $\bar{\omega} \subset \Omega$ and let $u \in L^2(\Omega)$. Then the following two properties hold:*

- (1) *If $u \in H^m(\Omega)$ then for every $|\varepsilon| < \text{dist}(\omega, \Omega^c)/m$ and $|\alpha| = m$,*

$$\|D_\varepsilon^\alpha(u)\|_{L^2(\omega)} \leq \|\partial^\alpha u\|_{L^2(\Omega)}.$$

- (2) *Conversely, if $u \in L^2(\Omega)$, and if there exists $\varepsilon_0, C_0 > 0$ such that for $|\alpha| = m$, and for all $|\varepsilon| \leq \varepsilon_0 < \text{dist}(\omega, \Omega^c)/m$, one has*

$$\|D_\varepsilon^\alpha(u)\|_{L^2(\omega)} \leq C_0,$$

then $\partial^\alpha u \in L^2(\omega)$ and we have

$$\|\partial^\alpha u\|_{L^2(\omega)} \leq C_0. \quad (2.12)$$

Proof. — We argue by induction: if $|\alpha| = 1$ then the statement is exactly Proposition 2.4.

So we assume that the statement is true for all $|\alpha| = m$ and we consider some α such that $|\alpha| = m + 1$, and $\varepsilon < \text{dist}(\omega, \Omega^c)/(m + 1)$. Then we can write

$$D_\varepsilon^\alpha = D_{\varepsilon e_i} D_\varepsilon^\beta,$$

with $|\beta| = m$ and for some e_i arbitrary chosen.

For the first part of the statement, we assume that $u \in H^{m+1}(\Omega)$. Let ω' be an open set such that $\omega \subset \omega' \subset \Omega$, and moreover such that

$$|\varepsilon| \leq \text{dist}(\omega, (\omega')^c)/m \quad \text{and} \quad |\varepsilon| \leq \text{dist}(\omega', \Omega^c).$$

This is possible because $|\varepsilon| < \text{dist}(\omega, \Omega^c)/(m + 1)$. Then the induction hypothesis says that

$$\|D_\varepsilon^\alpha(u)\|_{L^2(\omega)} = \|D_\varepsilon^\beta(D_{\varepsilon e_i} u)\|_{L^2(\omega)} \leq \|\partial^\beta D_{\varepsilon e_i} u\|_{L^2(\omega')} = \|D_{\varepsilon e_i} \partial^\beta u\|_{L^2(\omega')}.$$

But since $\partial^\beta u \in H^1(\Omega)$, we can apply Proposition 2.4 which says that

$$\|D_{\varepsilon e_i} \partial^\beta u\|_{L^2(\omega')} \leq \|\partial_i \partial^\beta u\|_{L^2(\Omega)} = \|\partial^\alpha u\|_{L^2(\Omega)},$$

which proves Proposition 2.5(1).

Now for part (2), we assume that $u \in L^2(\Omega)$, and there exists $C_0 > 0$ such that for every $|\alpha| = m + 1$, and for all $|\varepsilon| \leq \varepsilon_0 \leq \text{dist}(\omega, \Omega^c)/m$, one has

$$\|D_\varepsilon^\alpha(u)\|_{L^2(\omega)} \leq C_0.$$

Then again we write

$$D_\varepsilon^\alpha = D_{\varepsilon e_i} D_\varepsilon^\beta,$$

with $|\beta| = m$ and for some e_i arbitrary chosen. By applying the induction hypothesis we know that $\partial^\beta(D_{\varepsilon e_i} u) \in L^2(\omega)$ and

$$\|\partial^\beta(D_{\varepsilon e_i} u)\|_{L^2(\omega)} \leq C_0.$$

This actually yields

$$\|D_{\varepsilon e_i} \partial^\beta u\|_{L^2(\omega)} \leq C_0,$$

so a last application of Proposition 2.4 finally gives

$$\|\partial^\alpha u\|_{L^2(\omega)} = \|\partial_i \partial^\beta u\|_{L^2(\omega)} \leq C_0,$$

and so follows Proposition 2.5. □

2.9. Campanato spaces

We recall here some standard facts about Campanato spaces that will be used to achieve our $C^{m-1,\alpha}$ regularity result.

DEFINITION 2.6. — *Let Ω be an open set of $\mathbb{R}^N$. The Campanato space of parameter $\lambda \in \mathbb{R}^+$ is*

$$\mathcal{L}^{2,\lambda}(\Omega) := \left\{ u \in L^2(\Omega) \left| [u]_\lambda^2 := \sup_{\substack{x \in \Omega \\ 0 < r < r_0}} r^{-\lambda} \int_{\Omega \cap B(x,r)} |u - m(u; x, r)|^2 dx < \infty \right. \right\},$$

where

$$m(u; x, r) := \frac{1}{|\Omega \cap B(x, r)|} \int_{\Omega \cap B(x, r)} u \, dx.$$

PROPOSITION 2.7. — *Let Ω be an (ε_0, r_0) -Reifenberg flat domain in $\mathbb{R}^N$. If $\lambda \in (N, N + 2]$, then there exists a continuous injection*

$$\mathcal{L}^{2,\lambda}(\Omega) \hookrightarrow \mathcal{C}^{0,\alpha}(\bar{\Omega}), \quad (2.13)$$

with $\alpha := (\lambda - N)/p$.

A proof can be found in [11, Theorem 3.1] where bounded open domains with the following density condition are considered: there exists $A > 0$ such that

$$|\Omega \cap B(x, r)| \geq Ar^N, \quad 0 < r \leq r_0, \quad x \in \bar{\Omega}.$$

In our case, this is true since Reifenberg-flat domains satisfy the internal corkscrew condition (see for instance [12]).

PROPOSITION 2.8. — *Let Ω be a (ε_0, r_0) -Reifenberg flat domain and $u \in H_0^m(\Omega)$. If $\alpha \in (0, 1]$ is such that for all $x \in \Omega$ and $r \in (0, r_0)$,*

$$\int_{B(x,r)} |\nabla^m u(y)|^2 dy \leq C_0 r^{N+2\alpha-2},$$

then $u \in \mathcal{C}^{m-1,\alpha}(\bar{\Omega})$ and

$$\|u\|_{\mathcal{C}^{m-1,\alpha}(\bar{\Omega})} \leq CC_0,$$

where C depends on N and on $\text{diam}(\Omega)$.

Proof. — Let $v = \partial^\gamma u$ where $|\gamma| = m - 1$. Using the Poincaré inequality, we get

$$\begin{aligned} \int_{B(x,r)} |v(y) - m(v; x, r)|^2 dy &\leq C(N)r^2 \int_{B(x,r)} |\nabla v(y)|^2 dy \\ &\leq C(N)C_0 r^{N+2\alpha}. \end{aligned}$$

Hence, $v \in \mathcal{L}^{2,N+2\alpha}(\Omega)$ and with the injection (2.13), we know that $v \in C^{0,\alpha}(\bar{\Omega})$. But then for all $x \in \Omega$ and $r \leq r_0$,

$$\int_{B(x,r)} |v(y)|^2 dy \leq C_0 r^{N+2\alpha},$$

or in other words,

$$\int_{B(x,r)} |\nabla^{m-1} u(y)|^2 dy \leq C_0 r^{N+2\alpha},$$

and from the same argument as above we deduce that $\partial^\gamma v \in C^{0,\alpha}(\bar{\Omega})$ for all $|\gamma| = m - 2$. We can continue like this successively to arrive to the fact that $u \in \mathcal{C}^{m-1,\alpha}(\bar{\Omega})$. $\square$

3. Interior regularity by the translation method

We start with the interior regularity, which is easier since translations in all directions are admissible as test functions. This part works in a similar way as in the usual standard case of the Laplacian, but with some difficulty of working with higher order derivatives. For the convenience of the reader, and since it will be generalized later in a non-trivial way at the boundary, we write the full details.

THEOREM 3.1 (Interior regularity). — *Let $m \in \mathbb{N}^*$. Let $\mathcal{A}$ be an operator of order $2m$ of the form (2.3), and satisfying the ellipticity condition (2.4). Let $u \in H^m(B(0,1))$ be a weak solution for $\mathcal{A}u = f$ in $B(0,1)$ with $f \in H^l(B(0,1))$. Then $u \in H^{2m+l}(B(0,1/2))$ and*

$$\|u\|_{H^{2m+l}(B(0,\frac{1}{2}))} \leq C(\|u\|_{H^m(B(0,1))} + \|f\|_{H^l(B(0,1))}),$$

where $C > 0$ is a constant that depend on N , $\max_{\alpha,\beta} |a_{\alpha,\beta}|$, C_A , l and m .

Proof. — We will denote by $B = B(0,1)$. It is enough to prove the theorem for $f \in L^2(B)$. Indeed, assume that it holds true in this case. Then for $f \in H^l(B)$ one can apply the statement to the function $\partial^\alpha u$ with $|\alpha| \leq l$ that satisfies, in the sense of distributions,

$$\mathcal{A}\partial^\alpha u = \partial^\alpha f, \tag{3.1}$$

with $\partial^\alpha f \in L^2$. This would imply $\partial^\alpha u \in H^{2m}$ with

$$\|\partial^\alpha u\|_{H^{2m}(B(0, \frac{1}{2}))} \leq C(\|\partial^\alpha u\|_{H^m(B)} + \|\partial^\alpha f\|_{L^2(B)}),$$

thus we could easily conclude by an inductive argument on $|\alpha| = 0, 1, \dots, l$. The only thing is that since we would need to apply the result l times, the conclusion would hold in the smaller ball $B(0, \frac{1}{2^l})$ instead of $B(0, 1/2)$. But the constants are here arbitrary and the same result could be done from $B(0, R)$ to $B(0, r)$ for any $r < R$ without any substantial modification.

Therefore, in the sequel we will assume that $f \in L^2(B)$. The main starting point is the weak formulation of the problem which is

$$\int_B A \nabla^m u \cdot \nabla^m \varphi \, dx = \int_B f \varphi \, dx \quad \text{for every } \varphi \in H_0^m(B).$$

In the sequel we will use an inductive argument, assuming that $u \in H^{m+k}(B_k)$. We want to prove that $u \in H^{m+k+1}(B_{k+1})$ for a slightly smaller ball $B_{k+1} \subset B_k$, and arrive at $B(0, 1/2)$. For that purpose we define, for $1 \leq k \leq m$, the ball $B_k := B(0, 1/2 + (1/2)^k)$. We will also assume without mentioning it explicitly that $|h|$ is so small that Proposition 2.5 always applies with $\omega = B_{k+1}$ and $\Omega = B_k$.

Next, we fix $|\gamma| = k \leq m$. We will use the discrete derivatives defined in Section 2.8. The method of translations is based on the fact that for all $h \in \mathbb{R}^N$ such that $|h|$ is small enough, up to multiplying by a smooth cut-off function, the function $D_h u$ is admissible as a test function in the weak formulation, as being in H^m . This will provide an estimate on $D_h u$ that will allow us to win a derivative. Since we need to win m derivatives we will apply this m times.

The main ingredient is the weak formulation and the change of variables formula that yields

$$\begin{aligned} \int_{B_k} A \nabla^m D_h(D_\varepsilon^\gamma u) \cdot \nabla^m \varphi \, dx \\ = \int_{B_k} f D_{-h} D_{-\varepsilon}^\gamma \varphi \, dx, \quad \text{for every } \varphi \in H_0^m(B_k). \end{aligned} \quad (3.2)$$

To lighten the notation, from now on we will denote by $v = D_\varepsilon^\gamma u$.

Now we consider a cut-off function $\eta \in \mathcal{C}_c^\infty(B_k)$ such that for every $x \in \mathbb{R}^N$,

$$\eta(x) \in [0, 1], \quad \eta(x) = 1 \text{ on } B_{k+1}, \quad \text{and} \quad |\nabla^j \eta(x)| \leq C,$$

where $C > 0$ could depend on k . Then we take $\varphi = \eta^2 D_h D_\varepsilon^\gamma u = \eta^2 D_h v$ as a test function, for $|h| \leq 1/4$, which is admissible in the space $H_0^m(B_k)$.

Using Proposition 2.5, the term on the right-hand side of (3.2) can be estimated as

$$\left| \int_B f D_{-h} (D_{-\varepsilon}^\gamma (\eta^2 D_h v)) \, dx \right| \leq C \|f\|_{L^2(B_k)} \|\nabla \partial^\gamma (\eta^2 D_h v)\|_{L^2(B_k)}. \quad (3.3)$$

We will use this inequality later. For now we develop $\nabla^m (\eta^2 D_h v)$: for every $\alpha \in \mathbb{N}^N$ such that $|\alpha| = m$, the Leibniz formula (see (2.2)) says

$$\begin{aligned} \partial^\alpha (\eta^2 D_h v) &= \sum_{\beta \leq \alpha} \frac{\alpha!}{\beta!(\alpha - \beta)!} \partial^\beta (\eta^{2m}) \partial^{\alpha - \beta} (D_h v), \\ &= \eta^2 \partial^\alpha (D_h v) + \sum_{0 < \beta \leq \alpha} \frac{\alpha!}{\beta!(\alpha - \beta)!} \partial^\beta (\eta^2) \partial^{\alpha - \beta} (D_h v). \end{aligned}$$

In other words we have, in terms of vectors,

$$\nabla^m (\eta^2 D_h v) = \eta^2 \nabla^m D_h v + L(\eta, v), \quad (3.4)$$

where

$$L(\eta, v)_\alpha := \sum_{0 < \beta \leq \alpha} \frac{\alpha!}{\beta!(\alpha - \beta)!} \partial^\beta (\eta^2) \partial^{\alpha - \beta} (D_h v).$$

Notice that the expression in $L(\eta, v)_\alpha$ contains derivatives in v of order at most m , including the extra derivative in the direction of x_i provided by D_h . Indeed, for the particular case $\alpha = m e_i$ (i.e. when $\partial^\alpha = \partial_{x_i}^{(m)}$) then the condition $0 < \beta \leq \alpha$ means that $\beta_i > 0$ so that $|\alpha - \beta + e_i| \leq m$, as claimed. On the other hand for $\alpha \neq m e_i$ since $|\alpha| = m$, one must have $\alpha_i < m$ so that, again, $|\alpha - \beta + e_i| \leq m$.

We further decompose $L(\eta, v)$ as follows:

$$L(\eta, v) = \eta M(\eta, v) + G(\eta, v), \quad (3.5)$$

where $M(\eta, v)$ contains derivatives of v of order exactly m , and $G(\eta, v)$ contains lower order derivatives. In other words we have

$$\|M(\eta, v)\|_{L^2(B_k)} \leq C \|v\|_{H^m(B_k)} \leq C \|u\|_{H^{m+k}(B_k)}, \quad (3.6)$$

and

$$\|G(\eta, v)\|_{L^2(B_k)} \leq C \|v\|_{H^{m-1}(B_k)} \leq C \|u\|_{H^{m+k-1}(B_k)}. \quad (3.7)$$

Next, we use the decomposition (3.5) and then the discrete integration by parts formula contained in (2.11) to write

$$\begin{aligned}
 & \int_B A \nabla^m D_h v \cdot \nabla^m (\eta^2 D_h v) \, dx \\
 &= \int_B A \nabla^m D_h v \cdot \eta^2 \nabla^m D_h v \, dx \\
 & \quad + \int_B \eta A \nabla^m D_h v \cdot M(\eta, v) \, dx + \int_B A \nabla^m D_h v \cdot G(\eta, v) \, dx \\
 &= \int_B A \nabla^m D_h v \cdot \eta^2 \nabla^m D_h v \, dx \\
 & \quad + \int_B \eta A \nabla^m D_h v \cdot M(\eta, v) \, dx + \int_B A \nabla^m v \cdot D_{-h} G(\eta, v) \, dx.
 \end{aligned}$$

By (3.7) we know that

$$\left| \int_B A \nabla^m v \cdot D_{-h} G(\eta, v) \, dx \right| \leq C \|v\|_{H^m(B_k)}^2,$$

thus from to the ellipticity of A , and using also (3.7), we deduce that

$$\begin{aligned}
 & \int_B A \nabla^m D_h v \cdot \nabla^m (\eta^2 D_h v) \, dx \\
 & \geq \frac{1}{C_A} \int_B \eta^2 |\nabla^m D_h v|^2 \, dx \\
 & \quad + \int_B \eta A \nabla^m D_h v \cdot M(\eta, v) \, dx - C \|v\|_{H^m(B_k)}^2.
 \end{aligned} \tag{3.8}$$

Moreover,

$$\begin{aligned}
 & \left| \int_B \eta A \nabla^m D_h v \cdot M(\eta, v) \, dx \right| \\
 & \leq \frac{\lambda}{2} \sup |A| \int_B \eta^2 |\nabla^m D_h v|^2 \, dx + \frac{1}{2\lambda} \int_B |M(\eta, v)|^2 \, dx \\
 & \leq \frac{1}{2C_A} \int_B \eta^2 |\nabla^m D_h v|^2 \, dx + C \|v\|_{H^m(B_k)}^2.
 \end{aligned} \tag{3.9}$$

Returning to (3.2) and using (3.8), (3.9) and (3.3), we have finally obtained, using also Proposition 2.5 again, that

$$\begin{aligned}
 & \int_B \eta^2 |\nabla^m D_h v|^2 \, dx \\
 & \leq C \|v\|_{H^m(B_k)}^2 + C \|f\|_{L^2(B_k)} \|\nabla \partial^\gamma (\eta^2 D_h v)\|_{L^2(B_k)} \\
 & \leq C \|v\|_{H^m(B_k)}^2 + C \|f\|_{L^2(B)} \|v\|_{H^{k+2}(B_k)}.
 \end{aligned} \tag{3.10}$$

Since $v = D_\varepsilon^\gamma u$, with γ any multi-index with $|\gamma| = k$, and since $\eta = 1$ on B_{k+1} , the above inequality implies

$$\int_{B_{k+1}} |\nabla^{m+k+1} u|^2 dx \leq C \left(\|u\|_{H^{m+k}(B_k)}^2 + \|f\|_{L^2(B)}^2 + \|u\|_{H^{2k+2}(B_k)}^2 \right).$$

If $k \leq m-2$, this directly gives that $u \in H^{m+k+1}(B_{k+1})$ (because we assumed by induction that $u \in H^{m+k}(B_k)$ and $2k+2 \leq m+k$ only when $k \leq m-2$).

For $k = m-1$ we have to argue a bit differently. Indeed, we return back to (3.10) which provides

$$\begin{aligned} \int_B \eta^2 |\nabla^{k+1} D_h v|^2 dx \\ \leq C \left(\|v\|_{H^m(B_k)}^2 + \|f\|_{L^2(B_k)}^2 \right) + \frac{1}{4} \|\nabla^{k+1} (\eta^2 D_h v)\|_{L^2(B_k)}^2, \end{aligned} \quad (3.11)$$

Then, from Leibniz formula (3.4) yields

$$\nabla^{k+1} (\eta^2 D_h v) = \eta^2 \nabla^{k+1} D_h v + L(\eta, v), \quad (3.12)$$

where

$$\|L(\eta, v)\|_{L^2(B_k)} \leq C \|v\|_{H^{k+1}(B_k)} \leq C \|u\|_{H^{2k+1}(B_k)}. \quad (3.13)$$

we deduce that

$$\begin{aligned} \|\nabla^{k+1} (\eta^2 D_h v)\|_{L^2(B_k)}^2 &\leq 2 \|\eta^2 \nabla^{k+1} D_h v\|_{L^2(B_k)}^2 + 2 \|L(\eta, v)\|_{L^2(B_k)}^2, \\ &\leq 2 \|\eta \nabla^{k+1} D_h v\|_{L^2(B_k)}^2 + 2 \|v\|_{H^{k+1}(B_k)}^2, \end{aligned} \quad (3.14)$$

In the above we have also used that $\eta^2 \leq \eta$ because $\eta \in [0, 1]$.

Then using (3.11) and (3.14) we deduce that

$$\int_B \eta^2 |\nabla^{k+1} D_h v|^2 dx \leq C \left(\|v\|_{H^m(B_k)}^2 + \|f\|_{L^2(B_k)}^2 \right). \quad (3.15)$$

Since $2k+2 = m+k+1 = 2m$, all together we have proven, in the case when $k = m-1$, that

$$\int_{B_{k+1}} |\nabla^{2m} u|^2 dx \leq C \left(\|u\|_{H^{2m-1}(B_k)}^2 + \|f\|_{L^2(B)}^2 \right).$$

In any case we have proven that $\nabla^{m+k+1} u \in L^2(B_{k+1})$ and

$$\|\nabla^{m+k+1} u\|_{L^2(B_{k+1})} \leq C (\|u\|_{H^{m+k}(B_k)} + \|f\|_{L^2(B)}).$$

All in all, we have proved that if $u \in H^{m+k}(B_k)$ then $u \in H^{m+k+1}(B_{k+1})$. We can then argue by induction, since for $k = 0$ it is true that $u \in H^m(B)$, and since the induction works until $k = m-1$, we arrive at the conclusion that $u \in H^{2m}(B(0, 1/2))$, and so follows Theorem 3.1. $\square$

The following corollary will be needed later.

COROLLARY 3.2 (Interior regularity with $f = 0$). — *Let $\mathcal{A}$ be an operator of order $2m$ of the form (2.3), and satisfying the ellipticity condition (2.4). Let $u \in H^m(B(0, R))$ be a weak solution for $\mathcal{A}u = 0$ in $B(0, R)$. Then $\nabla^m u \in L^\infty(B(0, R/2))$ and*

$$\|\nabla^m u\|_{L^\infty(B(0, \frac{R}{2}))} \leq C_0 R^{-N/2} \|\nabla^m u\|_{L^2(B(0, R))}, \quad (3.16)$$

where $C_0 > 0$ is a constant that depends on N , $\max_{\alpha, \beta} |a_{\alpha, \beta}|$, C_A and m . In particular for all $r \leq R$ we have

$$\int_{B(x, r)} |\nabla^m u|^2 dx \leq C \left(\frac{r}{R}\right)^N \int_{B(x, R)} |\nabla^m u|^2 dx, \quad (3.17)$$

where $C := \omega_N C_0^2$.

Proof. — Assume first that $R = 1$. Then we can apply Theorem 3.1 to obtain that $\nabla^m u \in H^\ell(B(0, R/2))$ for all $\ell \in \mathbb{N}$ because here $f = 0$. Then the Sobolev embedding theorem says that, if $2\ell > N$, then $H^\ell \hookrightarrow \mathcal{C}^{0, \alpha}$, and in particular $\nabla^m u \in L^\infty$ with

$$\|\nabla^m u\|_{L^\infty(B(0, 1/2))} \leq C \|u\|_{H^m(B(0, 1))}.$$

We would like to apply this inequality to the rescaled function $u(Rx)$ but the problem is that the norm above is not homogeneous. For that purpose we consider $u - m$ where m is the average of u on $B(0, 1)$. Then we notice that $\nabla^m u = \nabla^m(u - m)$ and applying the above to $u - m$ instead of u yields

$$\|\nabla^m u\|_{L^\infty(B(0, 1/2))} \leq C \sum_{k=1}^m \|\nabla^k(u - m)\|_{L^2(B(0, 1))}.$$

By applying the Poincaré inequality we actually get

$$\|\nabla^m u\|_{L^\infty(B(0, 1/2))} \leq C \sum_{k=2}^m \|\nabla^k u\|_{L^2(B(0, 1))}.$$

Then we apply the same argument to $u - mx_i$ where m is the average of $\partial_i u$. Etc. After $m - 1$ application of Poincaré inequality we finally obtain

$$\|\nabla^m u\|_{L^\infty(B(0, 1/2))} \leq C_0 \|\nabla^m u\|_{L^2(B(0, 1))}.$$

The above inequality is now homogenous after scaling, thus applying it to the rescaled function $u(Rx)$ we finally arrive at (3.16).

Let us now prove (3.17). Let $a \in (0, 1)$ be given. Then

$$\begin{aligned} \int_{B(x, aR)} |\nabla^m u|^2 dx &\leq \|\nabla^m u\|_{L^\infty(B(x, aR))}^2 \omega_N a^N R^N \\ &\leq C_0^2 \omega_N a^N \int_{B(x, R)} |\nabla^m u|^2 dx, \end{aligned}$$

which gives (3.17) with $r = aR$, and concludes the proof of Corollary 3.2. $\square$

4. Boundary regularity in the case of a hyperplane

In this section we investigate the regularity up to the boundary for the Dirichlet problem, in the special case of a flat boundary. As before we consider an operator of order $2m$ of the form

$$\mathcal{A} := (-1)^m \sum_{|\beta|=|\alpha|=m} a_{\alpha, \beta} \partial^\alpha \partial^\beta = (-1)^m A \nabla^m \cdot \nabla^m,$$

and we focus this time on a weak solution $u \in H^m(B)$ for the problem in a half-ball:

$$\begin{cases} \mathcal{A}u = f & \text{in } B^+(0, 1) := B(0, 1) \cap \{x_N > 0\} \\ u \in H^m(B(0, 1)) & \text{and } u = 0 \text{ a.e. in } B(0, 1) \cap \{x_N < 0\}. \end{cases} \quad (4.1)$$

The weak formulation for (4.1) is

$$\int_B A \nabla^m u \cdot \nabla^m \varphi dx = \int_B f \varphi dx, \quad \text{for every } \varphi \in H_0^m(B^+).$$

Now here below is our regularity result at the boundary analogous to the interior estimate (Theorem 3.1). Notice that, compared to the interior estimate, below here in (4.2) the homogenous norm $\|\nabla^m u\|_2$ appears instead of the non homogeneous norm $\|u\|_{H^m}$. This is possible thanks to the Poincaré inequality available since a Dirichlet boundary condition is imposed on u on the flat part of $B^+(0, 1)$.

THEOREM 4.1 (Regularity with flat boundary). — *Let $m \in \mathbb{N}^*$. Let $\mathcal{A}$ be an operator of order $2m$ of the form (2.3) and satisfying the ellipticity condition (2.4). Let $u \in H^m(B(0, 1))$ a weak solution for $\mathcal{A}u = f$ in $B^+(0, 1)$ with $f \in H^\ell(B^+(0, 1))$, and such that $u = 0$ in $B(0, 1) \cap \{x_N < 0\}$. Then $u \in H^{2m+\ell}(B^+(0, 1/2))$ and*

$$\|u\|_{H^{2m+\ell}(B^+(0, \frac{1}{2}))} \leq C(\|\nabla^m u\|_{L^2(B^+(0, 1))} + \|f\|_{H^\ell(B^+(0, 1))}), \quad (4.2)$$

where $C > 0$ is a constant that depend on N , $\max_{\alpha, \beta} |a_{\alpha, \beta}|$, ℓ and m .

Proof. — The starting point of the proof is similar to the interior case, but there are some differences.

The main difference is that the translated functions $D_h u$ are admissible as a test function in $B^+(0, 1)$ only for horizontal directions, i.e. for $h \in \mathbb{R}e_i$ and $1 \leq i \leq N - 1$. Indeed, for $i = N$ the vertical translation would not preserve the boundary condition.

So let us assume first that $f \in L^2(B^+(0, 1))$ (the general case $f \in H^\ell(B^+(0, 1))$ will be treated later). For $\gamma \in \mathbb{N}^N$ containing only horizontal directions, in other words such that $\gamma_N = 0$, we can argue exactly as in the case of the interior regularity. We do not write again the details, and refer to the proof of Theorem 3.1. This reasoning yields that $\partial^\gamma \nabla^m u \in L^2(B^+(0, 1/2))$ for all $|\gamma| \leq m$ such that $\gamma_N = 0$, with the estimate

$$\|\partial^\gamma \nabla^m u\|_{L^2(B^+(0, \frac{1}{2}))} \leq C(\|\nabla^m u\|_{L^2(B^+(0, 1))} + \|f\|_{L^2(B^+(0, 1))}), \quad (4.3)$$

where $C > 0$ is a constant that depends on N , $\max_{\alpha, \beta} |a_{\alpha, \beta}|$, and m . Actually, the proof of Theorem 3.1 would give $\|u\|_{H^m(B^+(0, 1))}$ on the right-hand side but thanks to the Poincaré inequality (Lemma 2.3) and using that $u = 0$ on $B \cap \{x_N < 0\}$ we can bound

$$\|u\|_{H^m(B^+(0, 1))} \leq C\|\nabla^m u\|_{L^2(B^+(0, 1))},$$

from which we get (4.3).

Now to prove Theorem 4.1 we want to recover all the derivatives in the vertical direction, up to $2m$ -order. For this purpose we will argue by induction.

Step 1. — We suppose that $m = 1$, then we just proved that $\partial_i \partial_j u \in L^2(B^+(0, 1/2))$ for all $i \in \llbracket 1, N - 1 \rrbracket$ and $j \in \llbracket 1, N \rrbracket$. The operator $\mathcal{A}$ is of the form

$$\mathcal{A} = - \sum_{i=1}^N \sum_{j=1}^N a_{i,j} \partial_i \partial_j,$$

and by ellipticity we know that $a_{N,N} \neq 0$. From the identity $\mathcal{A}u = f \in L^2(B^+(0, 1))$ we have that

$$a_{N,N} \partial_N^2 u = -f + \sum_{(i,j) \neq (N,N)} a_{i,j} \partial_i \partial_j u \in L^2(B^+(0, 1/2)),$$

and we conclude that $u \in H^2(B^+(0, 1/2))$ with

$$\|u\|_{H^2(B^+(0, \frac{1}{2}))} \leq C(\|\nabla u\|_{L^2(B^+(0, 1))} + \|f\|_{L^2(B^+(0, 1))}).$$

This proves the desired conclusion for the particular case when $m = 1$.

Step 2. — We assume that $m \geq 2$ and we suppose that Theorem 4.1 is true for operators of order $2(m-1)$. Let us prove that it is still true for operator of order $2m$. For that purpose we decompose $\mathcal{A}$ as in the statement of Lemma 2.1:

$$\begin{aligned} \mathcal{A} &= (-1)^m \sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} \partial^\beta \partial^\alpha \\ &= (-1)^m \sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N=0 \text{ or } \beta_N=0}} a_{\alpha,\beta} \partial^\beta \partial^\alpha + (-1)^m \sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N>0 \text{ and } \beta_N>0}} a_{\alpha,\beta} \partial^\beta \partial^\alpha. \end{aligned}$$

In other words $\mathcal{A} = \mathcal{B} + \mathcal{C}$ with

$$\mathcal{B} := (-1)^m \sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N=0 \text{ or } \beta_N=0}} a_{\alpha,\beta} \partial^\beta \partial^\alpha$$

and

$$\mathcal{C} := (-1)^m \sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N>0 \text{ and } \beta_N>0}} a_{\alpha,\beta} \partial^\beta \partial^\alpha.$$

From the first part of the proof, we already know that $\mathcal{B}(u) \in L^2(B^+(0, 1/2))$ because it contains at most m derivatives in the vertical direction.

Now we consider the function $w := -\partial_N^2 u$ and we notice that $\mathcal{C}$ is an operator that acts as an operator of $2m-2$ order on the function w . Indeed,

$$\begin{aligned} \mathcal{C}(u) &= (-1)^m \sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N>0 \text{ and } \beta_N>0}} a_{\alpha,\beta} \partial^\beta \partial^\alpha u \\ &= (-1)^m \sum_{|\alpha'|=|\beta'|=m-1} a_{\alpha'+e_N, \beta'+e_N} \partial^\beta \partial^\alpha \partial_N^2 u = \mathcal{D}(w), \end{aligned}$$

where $\mathcal{D}$ is an operator of order $2m-2$. Furthermore, thanks to Lemma 2.1, we know that $\mathcal{D}$ is elliptic.

Since $u = H^m(B(0, 1))$, it follows that $w \in H^{m-2}(B(0, 1))$. Moreover, since $u = 0$ on the open set $B(0, 1) \cap \{x_N < 0\}$, the same still holds true for w . Then we can write

$$\mathcal{D}(w) = \mathcal{A}(u) - \mathcal{C}(u) = f - \mathcal{C}(u) \in L^2(B(0, 1)).$$

Since $\mathcal{D}$ is of order $2m-2$, we would like to deduce from the induction hypothesis that $w \in H^{2m-2}(B^+(0, 1))$. But unfortunately there is a gap of regularity here, because we only know that $w \in H^{m-2}(B(0, 1))$ and to use the induction hypothesis we would need that $w \in H^{m-1}(B(0, 1))$.

To overcome this problem we will use the Poincaré inequality in Lemma 2.2. Indeed, we can rewrite $\mathcal{D}(w)$ as follows:

$$\mathcal{D}(w) = -\partial_N \mathcal{D}(\partial_N u).$$

Since $\mathcal{D}(w) \in L^2(B^+(0, 1/2))$, we deduce that the distribution $z := \mathcal{D}(\partial_N u)$ verifies that

$$\partial_N z \in L^2(B^+(0, 1/2)).$$

Moreover, thanks to the interior regularity (Theorem 3.1), we know that

$$z \in L^2(B^+(0, 1/2) \cap \{\text{dist}(x, \partial B^+(0, 1/2)) > \delta_0\}).$$

By applying Lemma 2.2 we deduce that $z \in L^2(B^+(0, 1/2))$. In other words $\mathcal{D}(\partial_N u) \in L^2(B^+(0, 1/2))$ and now since $\partial_N u \in H^{m-1}(B)$, with $\partial_N u = 0$ on $B \cap \{x_N < 0\}$, the induction hypothesis applies. We conclude that $\partial_N u \in H^{2m-2}(B^+(0, 1/4))$. This fact, together with the previous analysis leads to say that $u \in H^{2m-1}(B^+(0, 1/4))$ and

$$\|u\|_{H^{2m-1}(B^+(0, \frac{1}{4}))} \leq C(\|\nabla^m u\|_{L^2(B^+(0,1))} + \|f\|_{L^2(B^+(0,1))}).$$

Now we are still missing one derivative because we need to achieve H^{2m} instead of H^{2m-1} . This will be done by use of the following observation: all in all, at this stage of the proof, we have a control on all the derivatives of the form $\partial^\alpha u$ with $|\alpha| = 2m$, provided that $|\alpha_N| \leq 2m - 1$. In other words, to conclude that $u \in H^{2m}$ we only miss the last derivative $\partial_N^{2m} u$. But this term appears in $\mathcal{A}$ only once. Namely, we have

$$\mathcal{A} = (-1)^m a_{me_N, me_N} \partial_N^{2m} + \mathcal{E} \quad (4.4)$$

where

$$\mathcal{E} := (-1)^m \sum_{\substack{|\alpha|=|\beta|=m \\ \alpha_N < m \text{ or } \beta_N < m}} a_{\alpha, \beta} \partial^\beta \partial^\alpha. \quad (4.5)$$

Since $\mathcal{E}$ contains at most $2m - 1$ derivatives in the e_N direction, we know that $\mathcal{E}(u) \in L^2(B^+(0, 1/4))$. Moreover, by ellipticity of $\mathcal{A}$ we deduce that the coefficient $a_{me_N, me_N} \neq 0$, and therefore, from (4.4) we get $\partial_N^{2m} u \in L^2(B^+(0, 1/4))$. In conclusion we have obtained,

$$\|u\|_{H^{2m}(B^+(0, \frac{1}{4}))} \leq C(\|\nabla^m u\|_{L^2(B^+(0,1))} + \|f\|_{L^2(B^+(0,1))}).$$

So follows the claim, with a radius $1/4$ instead of $1/2$. After m application of the inductive argument we arrive to the conclusion of the theorem in $B(0, (1/2)^m)$. Of course the radius $(1/2)^m$ has no importance and one could easily modify the proof in order to arrive in $B(0, 1/2)$ at the end, up to changing the constants. This finishes the proof of Theorem 4.1, in the case when $f \in L^2(B^+(0, 1))$.

We finally assume now that $f \in H^\ell(B^+(0, 1))$ and we want to get some further regularity. We cannot simply take a derivative of the equation as in

the interior case, because it would not preserve the Dirichlet boundary condition. Instead, we will use discrete derivatives again. Consider a horizontal direction $h = \varepsilon e_i$ with $1 \leq i \leq N-1$ and notice that $D_h u$ is a solution with $D_h f$ as a second member. In other words,

$$\int_B A \nabla^m D_h u \cdot \nabla^m \varphi \, dx = \int_{B_k} D_h f \varphi \, dx, \quad \text{for every } \varphi \in H_0^m(B_k). \quad (4.6)$$

Since $f \in H^1(B^+(0, 1))$, we know that $D_h f \in L^2$, uniformly in h . Applying all the reasoning as before to the function $D_h u$ leads to the estimate:

$$\|D_h u\|_{H^{2m}(B^+(0, \frac{1}{2}))} \leq C(\|\nabla^m D_h u\|_{L^2(B^+(0, 1))} + \|D_h f\|_{L^2(B^+(0, 1))}),$$

and this actually implies

$$\|\partial_i u\|_{H^{2m}(B^+(0, \frac{1}{2}))} \leq C(\|\nabla^m u\|_{L^2(B^+(0, 1))} + \|\partial_i f\|_{L^2(B^+(0, 1))}).$$

The above works for all $1 \leq i \leq N-1$. In other words, to prove that $u \in H^{2m+1}$ we only miss the derivative $\partial_N^{2m+1} u$. But now we come back to the decomposition already used in (4.4),

$$\mathcal{A} = (-1)^m a_{me_N, me_N} \partial_N^{2m} + \mathcal{E}$$

where $\mathcal{E}$ contains at most $2m-1$ derivatives in the e_N direction. Taking a derivative in the direction e_N yields,

$$(-1)^m a_{me_N, me_N} \partial_N^{2m+1} u = \partial_N f - \partial_N \mathcal{E}(u), \quad (4.7)$$

and since the second member lies in $L^2(B^+(0, 1))$ we deduce that $\partial_N^{2m+1} u \in L^2$. This proves that $u \in H^{2m+1}(B^+(0, 1/2))$ and actually,

$$\|u\|_{H^{2m+1}(B^+(0, \frac{1}{2}))} \leq C(\|\nabla^m u\|_{L^2(B^+(0, 1))} + \|f\|_{H^1(B^+(0, 1))}).$$

By a successive application of this reasoning in a sequence of balls (B_k) we finally arrive to the conclusion that

$$\|u\|_{H^{2m+\ell}(B^+(0, \frac{1}{2}))} \leq C(\|\nabla^m u\|_{L^2(B^+(0, 1))} + \|f\|_{H^\ell(B^+(0, 1))}),$$

which finishes the proof of Theorem 4.1. $\square$

The following corollary will be used in the next section.

COROLLARY 4.2 (Regularity with flat boundary and $f = 0$). — *Let $m \in \mathbb{N}^*$. Let $\mathcal{A}$ be an operator of order $2m$ of the form (2.3) and satisfying the ellipticity condition (2.4). Let $u \in H^m(B(0, R))$ be a weak solution for $\mathcal{A}u = 0$ in $B^+(0, R)$, and such that $u = 0$ in $B(0, R) \cap \{x_N < 0\}$. Then $\nabla^m u \in L^\infty(B^+(0, R/2))$ and*

$$\|\nabla^m u\|_{L^\infty(B^+(0, \frac{R}{2}))} \leq C_0 R^{-N/2} \|\nabla^m u\|_{L^2(B^+(0, R))},$$

where $C_0 > 0$ is a constant that depends on N , $\max_{\alpha, \beta} |a_{\alpha, \beta}|$, C_A and m .

Proof. — Theorem 4.1, applied to the rescaled function $u(Rx)$, says that $\nabla^m u \in H^\ell(B^+(0, R/2))$ for all $\ell \in \mathbb{N}$ because here $f = 0$. Then the Sobolev embedding theorem that says that, provided $2\ell > N$, then $H^\ell \hookrightarrow \mathcal{C}^{0,\alpha}$, and in particular $\nabla^m u \in L^\infty$. We also directly get the control on the norms because the norms that appear in the statement of Theorem 4.1 are homogeneous after scaling. $\square$

5. Energy decay for a Reifenberg-flat boundary

In this section we prove an energy decay property for the solution of a Dirichlet problem in a Reifenberg-flat domain. We begin with the case of a flat boundary, and then we argue by compactness to transfer this decay to a Reifenberg-flat boundary which is “sufficiently flat”. As before we consider an operator of order $2m$ of the form (2.3). We start with a decay property at the boundary, in the case when $f = 0$.

PROPOSITION 5.1 (Decay for a $\mathcal{A}$ -polyharmonic function near a Reifenberg-flat boundary). — *Let $a \in (0, 1/4)$, $b > 0$, $r_0 \in (0, 1]$ be given. Then there exists $\varepsilon_0 \in (0, 1)$ such that for every (ε_0, r_0) -Reifenberg flat domain Ω , $x \in \partial\Omega$, $r \leq r_0$, and for every $u \in H_0^m(\Omega)$ that is a weak solution of $\mathcal{A}(u) = 0$ in $\Omega \cap B(x, r)$, we have that*

$$\int_{B(x, ar)} |\nabla^m u|^2 dx \leq C a^{N-b} \int_{B(x, r)} |\nabla^m u|^2 dx, \quad (5.1)$$

where $C = C_0^2 \omega_N$, and $C_0 > 0$ is the constant of Corollary 4.2.

Proof. — We argue by contradiction. If the proposition is false then there exists $a \in (0, 1/4)$, $b > 0$, $r_0 \in (0, 1)$ such that for all $n \in \mathbb{N}$, there exists an open set Ω_n which is $(1/(n+1), r_0)$ -Reifenberg-flat and satisfies the following. There exists $x_n \in \partial\Omega_n$, $r_n \leq r_0$ and $u_n \in H_0^m(\Omega_n)$ such that $\mathcal{A}(u) = 0$ in $\Omega \cap B(x_n, r_n)$ and

$$C a^{N-b} \int_{B(x_n, r_n)} |\nabla^m u_n|^2 dx < \int_{B(x_n, ar_n)} |\nabla^m u_n|^2 dx. \quad (5.2)$$

By means of a translation and rotation, we reduce to the case in which $x_n = 0$ for all $n \in \mathbb{N}$ and the hyperplane induced by the definition of Reifenberg flat domain of Ω_n is $\mathcal{P}_0 := \{x_N = 0\}$.

We consider the normalized sequence

$$v_n(x) := r_n^{\frac{N}{2}-m} \|\nabla^m u_n(x)\|_{L^2(\Omega_n \cap B(0, r_n))}^{-1} u_n(r_n x).$$

The normalisation is fixed in such a way that

$$\int_{B_1} |\nabla^m v_n(x)|^2 dx = 1.$$

To lighten the notation, we will use the notation Ω_n for $\frac{1}{r_n}\Omega_n$. The functions v_n are still solutions of $\mathcal{A}(v_n) = 0$ in $\Omega_n \cap B(0, 1)$, we have $v_n = 0$ a.e. in $B(0, 1) \setminus \Omega_n$ and the sequence $(v_n)_{n \in \mathbb{N}}$ is bounded in $H^m(B(0, 1))$. Moreover, (5.2) becomes

$$Ca^{N-b} < \int_{B(0,a)} |\nabla^m v_n|^2 dx, \quad (5.3)$$

and $\Omega_n \cap B(0, 1)$ converges to $B^+(0, 1)$ for the Hausdorff distance, in the sense that

$$d_{\mathcal{H}}(\partial\Omega_n \cap B(0, 1), \{x_N = 0\} \cap B(0, 1)) \longrightarrow 0.$$

The idea is to pass to the limit and obtain a contradiction thanks to the regularity of polyharmonic functions at a flat boundary (Theorem 4.1).

By weak compactness, we can extract a subsequence, still denoted by v_n , which weakly converges to $v \in H^m(B(0, 1))$ and such that $(\nabla^i v_n)_{n \in \mathbb{N}}$ strongly converges to $\nabla^i v$ in $L^2(B(0, 1))$ for all $i \in \llbracket 0, m-1 \rrbracket$. The main objective is to show the following three facts:

CLAIMS. —

- (A₁) *The limit function v satisfies $v = 0$ a.e. in $B(0, 1) \cap \{x_N < 0\}$.*
- (A₂) *The limit function v is a weak solution of $\mathcal{A}(v) = 0$ in $B^+(0, 1)$.*
- (A₃) *The sequence $(v_n)_{n \in \mathbb{N}}$ strongly converges to v in $H^m(B(0, 1/2))$.*

Let us finish the proof by assuming that A₁, A₂ and A₃ hold. Passing to the limit in inequality (5.3) and applying Claim A₃ we obtain,

$$Ca^{N-b} \leq \int_{B(0,a)} |\nabla^m v|^2 dx. \quad (5.4)$$

On the other hand, since $\mathcal{A}v = 0$ in $B^+(0, 1)$ and $v = 0$ a.e. in $B(0, 1) \cap \{x_N < 0\}$, we can apply Corollary 4.2 to v which says that for all $a \leq 1/2$,

$$\|\nabla^m v\|_{L^\infty(B(0,a))} \leq C_0 \|\nabla^m v\|_{L^2(B(0,1))}.$$

In particular,

$$\begin{aligned} \int_{B(0,a)} |\nabla^m v|^2 dx &\leq \|\nabla^m v\|_{L^\infty(B^+(0,a))}^2 \omega_N a^N \\ &\leq C_0^2 \omega_N a^N \int_{B(0,1)} |\nabla^m v|^2 dx = C_0^2 \omega_N a^N. \end{aligned} \quad (5.5)$$

But since $C = C_0^2 \omega_N$, using (5.4) we arrive to

$$C_0^2 \omega_N a^{N-b} \leq C_0^2 \omega_N a^N,$$

which is a contradiction because $a < 1$.

So to finish the proof, we are left to prove A_1 , A_2 and A_3 . Actually, the claims A_1 and A_2 are easy to prove. Only A_3 is more delicate.

Proof of A_1 . — This fact directly follows from the Hausdorff convergence of $\partial\Omega_n \cap B(0, 1)$ to $\{x_N = 0\} \cap B(0, 1)$. Indeed, for any $x_0 \in B(0, 1) \cap \{x_N < 0\}$, there exists a ball $B(x_0, \varepsilon)$ such that $\bar{B}(x_0, \varepsilon) \subset B(0, 1) \cap \{x_N < 0\}$. From the Hausdorff convergence of Ω_n to $B^+(0, 1)$, we deduce that for n large enough, $B(x_0, \varepsilon) \subset \Omega_n^c \cap B(0, 1)$. But then we know by assumption that $v_n = 0$ on $B(x_0, \varepsilon)$. Since $v_n \rightarrow v$ strongly in L^2 , we deduce that $v = 0$ a.e. on $B(x_0, \varepsilon)$, and since x_0 is arbitrary, we conclude that $v = 0$ a.e. on $B(0, 1) \cap \{x_N < 0\}$, and the claim A_1 follows. $\square$

Proof of A_2 . Since $\partial\Omega_n \cap B(0, 1)$ converges to $\{x_N = 0\} \cap B(0, 1)$, we know that for any compact set $K \subset B^+(0, 1)$, there exist N_0 large enough such that $K \subset \Omega_n$ for all $n \geq N_0$. But then for any test function φ supported on K we have

$$\int_{B(0,1)} A \nabla^m v_n \cdot \nabla^m \varphi \, dx = 0.$$

Then from the weak convergence of v_n to v in H^m we can pass to the limit to obtain

$$\int_{B(0,1)} A \nabla^m v \cdot \nabla^m \varphi \, dx = 0.$$

Since this holds true for any compact set $K \subset B^+(0, 1)$, claim A_2 follows. $\square$

Proof of A_3 . — We show now that the sequence $(v_n)_{n \in \mathbb{N}}$ strongly converges to v in $H^m(B(0, \rho))$ for a radius $\rho \in (1/2, 1)$. Let ρ be such a radius. From now on we will write $x \cdot_A y := Ax \cdot y$ and $|x|_A^2 := x \cdot_A x$ for the norm and scalar product associated with A . By lower semicontinuity with respect to weak convergence we already know that

$$\int_{B(0,\rho)} |\nabla^m v|_A^2 \, dx \leq \liminf_{n \rightarrow +\infty} \int_{B(0,\rho)} |\nabla^m v_n|_A^2 \, dx.$$

To get the strong convergence in $B(0, \rho)$, it is enough to prove the converse, namely

$$\limsup_{n \rightarrow +\infty} \int_{B(0,\rho)} |\nabla^m v_n|_A^2 \, dx \leq \int_{B(0,\rho)} |\nabla^m v|_A^2 \, dx.$$

The idea is to modify v in order to make it admissible as a test function for v_n .

Let $\xi \in C_c^\infty(B(0, 1))$ be a smooth cut-off function such that $\xi = 1$ on $B(0, \rho)$. Since the function v belongs to $H^m(B(0, 1))$ and $v = 0$ a.e. in $B(0, 1) \cap \{x_N < 0\}$, it follows from the theory of stable domains (see [12,

Theorem 2]) that ξv belongs to $H_0^m(B^+(0, 1))$. We deduce that there exist a sequence $(w_i)_{i \in \mathbb{N}}$ in $\mathcal{C}_c^\infty(B^+(0, 1))$ which strongly converges to ξv in $H^m(\mathbb{R}^N)$. The main point is that for i being fixed, we know from the Hausdorff convergence of Ω_n that for all n large enough, $w_i \in H_0^m(\Omega_n \cap B(0, 1))$.

Now we pick $\tilde{\rho} \in (1/2, \rho)$ and consider another cut-off function $\eta_{\tilde{\rho}} \in \mathcal{C}_c^\infty(B(0, \rho))$ such that $\eta_{\tilde{\rho}} = 1$ on $B(0, \tilde{\rho})$ and

$$|\nabla^k \eta_{\tilde{\rho}}| \leq C(\rho - \tilde{\rho})^{-k}, \quad 1 \leq k \leq m,$$

for a universal constant $C > 0$.

Since v_n is a weak solution for $\mathcal{A}(v_n) = 0$ in $B(0, \rho) \cap \Omega_n$, it is a local energy minimizer and $z_{\tilde{\rho}, i} := (1 - \eta_{\tilde{\rho}})v_n + \eta_{\tilde{\rho}}w_i$ is a competitor. That is, $z_{\tilde{\rho}, i} \in H^m(B(0, \rho))$ and is equal to v_n on $\partial B(0, \rho)$. We deduce that

$$\int_{B(0, \rho)} |\nabla^m v_n|_A^2 dx \leq \int_{B(0, \rho)} |\nabla^m z_{\tilde{\rho}, i}|_A^2 dx. \quad (5.6)$$

Next, a computation reveals that, for all multi-indices α_1 satisfying $|\alpha_1| = m$ we have

$$\begin{aligned} \partial^{\alpha_1} z_{\tilde{\rho}, i} &= \partial^{\alpha_1} (v_n + \eta_{\tilde{\rho}}(w_i - v_n)) \\ &= \partial^{\alpha_1} v_n + \sum_{\alpha_1 = \alpha_2 + \alpha_3} \frac{\alpha_1!}{\alpha_2! \alpha_3!} \partial^{\alpha_2} \eta_{\tilde{\rho}} \partial^{\alpha_3} (w_i - v_n) \\ &= \partial^{\alpha_1} v_n + \eta_{\tilde{\rho}} \partial^{\alpha_1} (w_i - v_n) + \sum_{\substack{\alpha_1 = \alpha_2 + \alpha_3 \\ |\alpha_2| > 0}} \frac{\alpha_1!}{\alpha_2! \alpha_3!} \partial^{\alpha_2} \eta_{\tilde{\rho}} \partial^{\alpha_3} (w_i - v_n) \\ &= \eta_{\tilde{\rho}} \partial^{\alpha_1} w_i + (1 - \eta_{\tilde{\rho}}) \partial^{\alpha_1} v_n + \sum_{\substack{\alpha_1 = \alpha_2 + \alpha_3 \\ |\alpha_2| > 0}} \frac{\alpha_1!}{\alpha_2! \alpha_3!} \partial^{\alpha_2} \eta_{\tilde{\rho}} \partial^{\alpha_3} (w_i - v_n). \end{aligned}$$

By convexity, we get

$$\begin{aligned} &|\nabla^m z_{\tilde{\rho}, i}|_A^2 \\ &\leq \eta_{\tilde{\rho}} |\nabla^m w_i|_A^2 + (1 - \eta_{\tilde{\rho}}) |\nabla^m v_n|_A^2 \\ &\quad + C(m) \sum_{k=1}^m \frac{1}{(\rho - \tilde{\rho})^{2k}} |\nabla^{m-k} (w_i - v_n)|^2 + C(m) \sum_{k=1}^m \frac{1}{(\rho - \tilde{\rho})^{2k}} \\ &\quad \times \left| [\eta_{\tilde{\rho}} \nabla^m w_i + (1 - \eta_{\tilde{\rho}}) \nabla^m v_n] \cdot_A [\nabla^{m-k} (w_i - v_n)] \right|. \end{aligned}$$

Using (5.6), one has

$$\begin{aligned}
 & \int_{B(0,\rho)} \eta_{\tilde{\rho}} |\nabla^m v_n|_A^2 \, dx \\
 & \leq \int_{B(0,\rho)} \eta_{\tilde{\rho}} |\nabla^m w_i|_A^2 \, dx \\
 & \quad + C(m) \sum_{k=1}^m \frac{1}{(\rho - \tilde{\rho})^{2k}} \int_{B(0,\rho)} |\nabla^{m-k}(w_i - v_n)|_A^2 \, dx \\
 & \quad + C(m) \sum_{k=1}^m \frac{1}{(\rho - \tilde{\rho})^{2k}} \int_{B(0,\rho)} |[\eta_{\tilde{\rho}} \nabla^m w_i + (1 - \eta_{\tilde{\rho}}) \nabla^m v_n] \\
 & \quad \cdot_A [\nabla^{m-k}(w_i - v_n)]| \, dx.
 \end{aligned}$$

Recall that $(\nabla^k v_n)_{n \in \mathbb{N}}$ strongly converges in $L^2(B(0, 1))$ for all $k \in \llbracket 0, m-1 \rrbracket$, thus taking the limsup as $n \rightarrow +\infty$ in the above inequality we get

$$\begin{aligned}
 & \limsup_{n \rightarrow +\infty} \int_{B(0,\tilde{\rho})} |\nabla^m v_n|_A^2 \, dx \leq \limsup_{n \rightarrow +\infty} \int_{B(0,\rho)} \eta_{\tilde{\rho}} |\nabla^m v_n|_A^2 \, dx \\
 & \leq \int_{B(0,\rho)} \eta_{\tilde{\rho}} |\nabla^m w_i|_A^2 \, dx + C(m) \sum_{k=1}^m \frac{1}{(\rho - \tilde{\rho})^{2k}} \int_{B(0,\rho)} |\nabla^{m-k}(w_i - v)|_A^2 \, dx \\
 & \quad + C(m) \sum_{k=1}^m \frac{1}{(\rho - \tilde{\rho})^{2k}} \int_{B(0,\rho)} [\eta_{\tilde{\rho}} \nabla^m w_i + (1 - \eta_{\tilde{\rho}}) \nabla^m v] \\
 & \quad \cdot_A [\nabla^{m-k}(w_i - v)] \, dx.
 \end{aligned}$$

Then we take the limit as $i \rightarrow +\infty$, and we recall that w_i converges strongly in H^m to ξv , and ξv is equal to v on $B(0, \rho)$. Therefore, all the terms with $w_i - v$ disappear and we arrive at

$$\limsup_{n \rightarrow +\infty} \int_{B(0,\tilde{\rho})} |\nabla^m v_n|_A^2 \, dx \leq \int_{B(0,\rho)} \eta_{\tilde{\rho}} |\nabla^m v|_A^2 \, dx \leq \int_{B(0,\rho)} |\nabla^m v|_A^2 \, dx.$$

Finally, by letting $\rho \rightarrow \tilde{\rho}$ on the right hand side, we get the desired limsup inequality. This together with the liminf inequality and the weak-convergence, proves that v_n converges strongly to v in $H^m(B(0, \tilde{\rho}))$. Since $\tilde{\rho} > 1/2$, this achieves the proof of Claim A_3 , and the Proposition 5.1 follows. $\square$

We are now in position to prove a boundary decay property of a Dirichlet solution in a Reifenberg-flat domain.

PROPOSITION 5.2 (Energy decay for a solution at a boundary point). *Let $q \geq 2$ be such that $mq \geq N$ if $2m < N$, and $q = 2$ otherwise. Let $\eta \in (0, 1)$ be given. Then there exists $\varepsilon_0 \in (0, 1)$ such that for all $r_0 > 0$,*

for any (ε_0, r_0) -Reifenberg flat domain Ω , and for any $f \in L^q(\Omega)$, the weak solution $u \in H_0^m(\Omega)$ of $\mathcal{A}(u) = f$ satisfies

$$\int_{B(x,r)} |\nabla^m u|^2 dx \leq C r^{N-\eta} \|f\|_{L^q(\Omega)}^2, \quad \forall r \in (0, r_0), \quad x \in \partial\Omega,$$

where $C := C(A, N, m, r_0, \eta, \Omega) > 0$.

Proof of Proposition 5.2. — Consider a real $0 < b < \eta$ and fix $a \in (0, 1/4)$ satisfying

$$a \leq \left(\frac{1}{4C_A \max |A|} \right)^{\frac{1}{\eta-b}}, \quad (5.7)$$

where C_A is the ellipticity constant of A . From Proposition 5.1, for these choices of a and b , there exists $\varepsilon_0 \in (0, 1)$ such that the decay property (5.1) holds for any polyharmonic function.

Let $u \in H_0^m(\Omega)$ be the weak solution of $\mathcal{A}(u) = f$ in Ω and let $x \in \partial\Omega$. and for every $k \in \mathbb{N}$, we define $B_k := B(x, a^k r_0)$. Let v_k be the polyharmonic replacement of u in B_k , more precisely v_k is the solution of

$$\min_{w \text{ s.t. } w-u \in H_0^m(B_k \cap \Omega)} \int_{B_k} A \nabla^m w \cdot \nabla^m w dx.$$

Then Proposition 5.1 applies to v_k and yields

$$\int_{B_{k+1}} |\nabla^m v_k|^2 dx \leq C_1 a^{N-b} \int_{B_k} |\nabla^m v_k|^2 dx. \quad (5.8)$$

We will prove by induction over $k \geq k_0$ with $k_0 := \lfloor \frac{N}{\eta} \rfloor + 1$ the inequality

$$\int_{B_k} |\nabla^m u|^2 dx \leq C_I a^{k(N-\eta)} \|f\|_{L^q(\Omega)}^2, \quad (H_k)$$

for a constant $C_I > 0$ that will be fixed later.

First of all we have, by ellipticity of A ,

$$\begin{aligned} \int_{B_{k_0}} |\nabla^m u|^2 dx &\leq \int_{\Omega} |\nabla^m u|^2 dx \leq C_A \int_{\Omega} A \nabla^m u \cdot \nabla^m u dx \\ &= C_A \int_{\Omega} f u dx \leq C_A \|f\|_{L^2(\Omega)} \|u\|_{L^2(\Omega)}. \end{aligned}$$

Next by the Poincaré inequality in $H_0^m(\Omega)$,

$$\|u\|_{L^2(\Omega)} \leq C_P \|\nabla^m u\|_{L^2(\Omega)},$$

so finally,

$$\int_{B_{k_0}} |\nabla^m u|^2 dx \leq C_A^2 C_P^2 \|f\|_{L^2(\Omega)}^2 \leq C_A^2 C_P^2 |\Omega|^{1-\frac{2}{q}} \|f\|_{L^q(\Omega)}^2. \quad (5.9)$$

Hence, (H_k) holds true for $k_0 = \lfloor \frac{N}{\eta} \rfloor + 1$ provided that

$$C_A^2 C_P^2 |\Omega|^{1-\frac{2}{q}} a^{-k_0(N-\eta)} \leq C_I.$$

We assume now that (H_k) holds for some $k \geq k_0$ and we write

$$\begin{aligned} & \int_{B_{k+1}} |\nabla^m u|^2 dx \\ & \leq 2 \int_{B_{k+1}} |\nabla^m v_k|^2 dx + 2 \int_{B_{k+1}} |\nabla^m(u - v_k)|^2 dx \\ & \leq 2a^{N-b} \int_{B_k} |\nabla^m v_k|^2 dx + 2 \int_{B_{k+1}} |\nabla^m(u - v_k)|^2 dx \\ & \leq C_A 2a^{N-b} \int_{B_k} A \nabla^m u \cdot \nabla^m u dx + 2 \int_{B_k} |\nabla^m(u - v_k)|^2 dx. \\ & \leq \max(|A|) C_A 2a^{N-b} \int_{B_k} |\nabla^m u|^2 dx + 2C_A \int_{B_k} |\nabla^m(u - v_k)|_A dx. \end{aligned} \quad (5.10)$$

The second-to-last inequality is a consequence of the fact that u is a competitor for the minimization problem satisfied by v_k , and we have used as before the notation $|v|_A := Av \cdot v$. Moreover, since $u - v_k$ is A -orthogonal to v_k , the Pythagoras inequality yields

$$0 \leq \int_{B_k} |\nabla^m(u - v_k)|_A^2 dx = \int_{B_k} |\nabla^m u|_A^2 dx - \int_{B_k} |\nabla^m v_k|_A^2 dx. \quad (5.11)$$

On the other hand, since now u is a minimizer of

$$\min \left\{ \frac{1}{2} \int_{B_k} |\nabla^m w|_A dx - \int_{B_k} fw dx \mid w \in u + H_0^m(\Omega \cap B_k) \right\},$$

and since the function v_k is a competitor for this minimising problem we deduce that

$$\int_{B_k} |\nabla^m u|_A^2 dx - 2 \int_{B_k} fu dx \leq \int_{B_k} |\nabla^m v_k|_A^2 dx - 2 \int_{B_k} fv_k dx,$$

and from (5.11),

$$\int_{B_k} |\nabla^m(u - v_k)|_A^2 dx \leq 2 \int_{B_k} f(u - v_k) dx. \quad (5.12)$$

Then we use the Sobolev embedding theorem together with the Poincaré inequality (Lemma 2.3) which implies $u - v_k \in L^p$ with

$$p = \begin{cases} \frac{2N}{N-2m} & \text{if } 2m < N \\ \text{arbitrary large} & \text{if } 2m = N \\ +\infty & \text{if } 2m > N, \end{cases} \quad (5.13)$$

and moreover,

$$\begin{aligned} \|u - v_k\|_{L^p(B_k)} &\leq C_S \|\nabla^m(u - v_k)\|_{L^2(B_k)} \\ &\leq C_S \sqrt{C_A} \left(\int_{B_k} |\nabla^m(u - v_k)|_A^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

Hence, returning back to (5.12) and using Hölder's inequality,

$$\begin{aligned} \int_{B_k} |\nabla^m(u - v_k)|_A^2 dx &\leq 2 \int_{B_k} f(u - v_k) dx \\ &\leq 2 \|f\|_{L^{p'}(B_k)} \|u - v_k\|_p \\ &\leq 2 C_S \sqrt{C_A} \|f\|_{L^{p'}(B_k)} \left(\int_{B_k} |\nabla^m(u - v_k)|_A^2 dx \right)^{\frac{1}{2}}, \end{aligned}$$

which finally yields,

$$\int_{B_k} |\nabla^m(u - v_k)|_A^2 dx \leq 4 C_S^2 C_A \|f\|_{L^{p'}(B_k)}^2. \quad (5.14)$$

Here, $p' \geq 1$ is the conjugate exponent of p , defined in (5.13), so that

$$p' = \begin{cases} \frac{2N}{N+2m} & \text{if } 2m < N \\ \text{arbitrary close to } 1 & \text{if } 2m = N \\ 1 & \text{if } 2m > N. \end{cases} \quad (5.15)$$

In particular, under the assumptions of the proposition we know that $q \geq p'$. Let s be such that $sp' = q$ (with $s = +\infty$ if $q = +\infty$). Then

$$\|f\|_{L^{p'}(B_k)}^{p'} \leq |B_k|^{\frac{1}{s'}} \left(\int_{B_k} |f|^q dx \right)^{\frac{p'}{q}},$$

and (5.14) gives

$$\begin{aligned} \int_{B_k} |\nabla^m(u - v_k)|_A^2 dx &\leq 4 C_S^2 C_A C_N (a^k r_0)^{\frac{2N}{p's'}} \|f\|_{L^q(\Omega)}^2 \\ &= 4 C_S^2 C_A C_N (a^k r_0)^{2N \frac{q-p'}{p'q}} \|f\|_{L^q(\Omega)}^2. \end{aligned}$$

Returning back to (5.10) we have obtained,

$$\begin{aligned} \int_{B_{k+1}} |\nabla^m u|^2 dx &\leq \max(|A|) C_A 2 C_I a^{N-b} a^{k(N-\eta)} \|f\|_{L^q(\Omega)}^2 \\ &\quad + 8 C_A^2 C_S(p)^2 C_N (a^k r_0)^{2N \frac{q-p'}{p'q}} \|f\|_{L^q(\Omega)}^2. \end{aligned} \quad (5.16)$$

Let us check that each term in the sum on the right hand side above is less than $\frac{1}{2} a^{(k+1)(N-\eta)} C_I \|f\|_{L^q(\Omega)}^2$, which would be enough to finish the proof of (H_k) .

For the first term it will be true provided that

$$\max(|A|)C_A 2a^{N-b} \leq \frac{1}{2}a^{N-\eta},$$

which holds true thanks to our definition of a (see (5.7)).

For the second term we notice that a has the following exponent:

$$\gamma := k2N \frac{q-p'}{p'q},$$

and we want this to be greater than $(k+1)(N-\eta)$. Then using that $k \geq k_0 \geq \frac{N}{\eta}$ we estimate

$$\begin{aligned} & \gamma - (k+1)(N-\eta) \\ &= k2N \frac{q-p'}{qp'} - (k+1)(N-\eta) \\ &= \frac{1}{qp'}(2Nkq - k2Np' - Nkqp' + qp'(k\eta - N) + qp'\eta) \\ &\geq \frac{1}{qp'}(2Nkq - k2Np' - Nkqp') + \eta \\ &\geq \eta, \end{aligned} \tag{5.17}$$

provided that $2Nkq - k2Np' - Nkqp' \geq 0$ or equivalently that

$$q(2-p') \geq 2p'. \tag{5.18}$$

Then we consider three cases:

Case 1. — If $2m > N$, then $p' = 1$ and the condition in (5.18) reduces to $q \geq 2$, which is true.

Case 2. — If $2m = N$, then p' can be chosen arbitrary close to 1 thus we take p' such that

$$p' \leq \frac{2q}{1+q},$$

and then (5.18) is true.

Case 3. — If $2m < N$, then $p' = \frac{2N}{N+2m}$. Thus a simple computation reveals that (5.18) is true when $mq \geq N$, which is what we have assumed in the statement of the proposition.

Conclusion. — In all cases we have proved that the last term in (5.16) is always bounded by

$$8C_A^2 C_S(p)^2 C_N r_0^{2N \frac{q-p'}{qp'}} a^\eta a^{(k+1)(N-\eta)} \|f\|_{L^q(\Omega)}^2,$$

which is less than $\frac{1}{2}a^{(k+1)(N-\eta)}C_I\|f\|_{L^q(\Omega)}^2$ provided that

$$8C_A^2C_S(p)^2C_Nr_0^{2N\frac{q-p'}{qp'}}a^\eta \leq \frac{C_I}{2},$$

which can be true by defining C_I appropriately. This finishes the proof of (H_k) .

Now to finish the proof, let $r \in (0, a^{k_0}r_0)$. We can choose $k \geq k_0$ such that $r < a^k r_0$ and $a^{k+1}r_0 \leq r$. One has

$$\begin{aligned} \int_{B(x,r)} |\nabla^m u|^2 dx &\leq \int_{B(x, a^k r_0)} |\nabla^m u|^2 dx \\ &\leq C_I a^{k(N-\eta)} \|f\|_{L^q(\Omega)}^2 \\ &\leq C_I \left(\frac{r}{a r_0} \right)^{N-\eta} \|f\|_{L^q(\Omega)}^2. \end{aligned}$$

On the other hand if $r \in (a^{k_0}r_0, r_0)$, we can argue as for the proof of (5.9) yielding

$$\begin{aligned} \int_{B_r} |\nabla^m u|^2 dx &\leq C_A C_P |\Omega|^{1-\frac{2}{q}} \|f\|_{L^q(\Omega)}^2 \\ &\leq C_A C_P |\Omega|^{1-\frac{2}{q}} \frac{1}{(a^{k_0}r_0)^{N-\eta}} r^{N-\eta} \|f\|_{L^q(\Omega)}^2, \end{aligned} \quad (5.19)$$

and the proof is completed with

$$C := \max \left(C_I (a r_0)^{\eta-N}, C_A C_P |\Omega|^{1-\frac{2}{q}} \frac{1}{(a^{k_0}r_0)^{N-\eta}} \right). \quad \square$$

Following exactly the same proof as the one of Proposition 5.2, we can also obtain the same for interior points.

PROPOSITION 5.3 (Energy decay for a solution in the interior). — *Let $q \geq 2$ be such that $mq \geq N$ if $2m < N$, and $q = 2$ otherwise. Let $\eta \in (0, 1)$ be given. Then there exists $r_0 \in (0, 1)$ depending on N , $\max_{\alpha, \beta} |a_{\alpha, \beta}|$, and m such that for all $\Omega \subset \mathbb{R}^N$ open, for all $f \in L^q(\Omega)$, the weak solution $u \in H_0^m(\Omega)$ of $\mathcal{A}(u) = f$ satisfies*

$$\begin{aligned} \int_{B(x,r)} |\nabla^m u|^2 dx &\leq C r^{N-\eta} \|f\|_{L^q(\Omega)}^2, \\ &\forall r \in (0, r_0), \quad \forall x \in \Omega, \quad \text{s.t. } B(x, r) \subset \Omega, \end{aligned}$$

where $C := C(A, N, m, r_0, \eta, \Omega) > 0$.

Proof. — The proof is exactly the same as Proposition 5.2 using a polyharmonic replacement of u in $B(x, r)$ to obtain some decay property on the energy. The only difference is that we use the interior decay property of a polyharmonic function stated in Corollary 3.2 instead of the boundary decay of Proposition 5.1 that was used in the proof of Proposition 5.2. $\square$

6. Boundary regularity in a Reifenberg flat domain

This very short section contains the proof of our main regularity result, namely Theorem 1.2. The strategy is to use the energy decay of Propositions 5.2 and 5.3 of the derivative of the solution to estimate its Campanato space norm. Here we restate our main result.

THEOREM 6.1. — *Let $\alpha \in (0, 1)$ and $q \geq 2$ be such that $mq \geq N$ if $2m < N$, and $q = 2$ otherwise. There exists $\varepsilon_0 \in (0, 1)$ and $r_0 \in (0, 1]$ such that for every (ε_0, r_0) -Reifenberg-flat domain $\Omega \subset \mathbb{R}^N$, for every function $f \in L^q(\Omega)$, if $u \in H_0^m(\Omega)$ is the weak solution of $\mathcal{A}(u) = f$, then $u \in \mathcal{C}^{m-1, \alpha}(\bar{\Omega})$, and*

$$\|u\|_{\mathcal{C}^{m-1, \alpha}(\bar{\Omega})} \leq C \|f\|_{L^q(\Omega)},$$

where $C > 0$ depends on N, A, α, Ω and m .

Proof. — Using the energy decay of Proposition 5.2, for every $\eta \in (0, 1)$, there exists $\varepsilon_0 \in (0, 1)$ and $r_0 \in (0, 1]$ such that for all $y \in \partial\Omega$ and $r \in (0, r_0)$,

$$\int_{\Omega \cap B(y, r)} |\nabla^m u|^2 dx \leq C r^{N-\eta} \|f\|_{L^q(\Omega)}^2. \quad (6.1)$$

Furthermore, Proposition 5.3 yields

$$\begin{aligned} \int_{B(x, r)} |\nabla^m u|^2 dx &\leq C r^{N-\eta} \|f\|_{L^q(\Omega)}^2, \\ \forall r \in (0, r_1), \quad \forall x \in \Omega, \quad \text{s.t. } B(x, r) &\subset \Omega. \end{aligned} \quad (6.2)$$

All together, for any ball $B(x, r)$ with $x \in \Omega$ and $r \leq r_2 := \min(r_0, r_1)$, applying either (6.2) if $B(x, r) \subset \Omega$ or (6.1) in $B(z, 2r)$ with $z \in \partial\Omega$ such that $B(x, r) \subset B(z, 2r)$ if $B(x, r) \cap \partial\Omega \neq \emptyset$, we deduce that

$$\int_{B(x, r)} |\nabla^m u|^2 dx \leq C r^{N-\eta} \|f\|_{L^q(\Omega)}^2, \quad \forall r \in (0, r_2), \quad x \in \Omega. \quad (6.3)$$

By Proposition 2.8, we get $u \in \mathcal{C}^{m-1, \alpha}(\bar{\Omega})$ with $\alpha := 1 - \eta/2$ and

$$\|u\|_{\mathcal{C}^{m-1, \alpha}(\bar{\Omega})} \leq C \|f\|_{L^q(\Omega)},$$

and Theorem 6.1 follows. $\square$

Bibliography

- [1] S. AGMON, A. DOUGLIS & L. NIRENBERG, “Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I”, *Commun. Pure Appl. Math.* **12** (1959), p. 623-727.
- [2] ———, “Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. II”, *Commun. Pure Appl. Math.* **17** (1964), p. 35-92.
- [3] A. BARTON, S. HOFMANN & S. MAYBORODA, “Dirichlet and Neumann boundary values of solutions to higher order elliptic equations”, *Ann. Inst. Fourier* **69** (2019), no. 4, p. 1627-1678.
- [4] A. BARTON & S. MAYBORODA, “Higher-order elliptic equations in non-smooth domains: a partial survey”, in *Harmonic analysis, partial differential equations, complex analysis, Banach spaces, and operator theory. Volume 1. Celebrating Cora Sadosky’s life*, Association for Women in Mathematics Series, vol. 4, Springer, 2016, p. 55-121.
- [5] S.-S. BYUN & Y. CHO, “Lorentz–Morrey regularity for nonlinear elliptic problems with irregular obstacles over Reifenberg flat domains”, *Discrete Contin. Dyn. Syst.* **35** (2015), no. 10, p. 4791-4804.
- [6] S.-S. BYUN, Y. CHO & D. K. PALAGACHEV, “Global weighted estimates for nonlinear elliptic obstacle problems over Reifenberg domains”, *Proc. Am. Math. Soc.* **143** (2015), no. 6, p. 2527-2541.
- [7] S.-S. BYUN & L. WANG, “Elliptic equations with BMO nonlinearity in Reifenberg domains”, *Adv. Math.* **219** (2008), no. 6, p. 1937-1971.
- [8] H. DONG & D. KIM, “Higher order elliptic and parabolic systems with variably partially BMO coefficients in regular and irregular domains”, *J. Funct. Anal.* **261** (2011), no. 11, p. 3279-3327.
- [9] L. C. EVANS, *Partial differential equations*, 2nd ed., Graduate Studies in Mathematics, vol. 19, American Mathematical Society, 2010.
- [10] F. GAZZOLA, H.-C. GRUNAU & G. SWEERS, *Polyharmonic boundary value problems. Positivity preserving and nonlinear higher order elliptic equations in bounded domains*, Lecture Notes in Mathematics, vol. 1991, Springer, 2010.
- [11] M. GIAQUINTA, *Introduction to regularity theory for nonlinear elliptic systems*, Lectures in Mathematics, ETH Zürich, Birkhäuser, 1993.
- [12] J.-F. GROSJEAN, A. LEMENANT & R. MOUGENOT, “Stable domains for higher order elliptic operators”, *Comptes Rendus. Mathématique* **362** (2024), p. 1189-1203.
- [13] I. GUILLÉN-MOLA, M. PRATS & X. TOLSA, “The dimension of planar elliptic measures arising from Lipschitz matrices in Reifenberg flat domains”, 2024, <https://arxiv.org/abs/2406.11604v1>.
- [14] C. E. KENIG & T. TORO, “Poisson kernel characterization of Reifenberg flat chord arc domains”, *Ann. Sci. Éc. Norm. Supér. (4)* **36** (2003), no. 3, p. 323-401.
- [15] A. LEMENANT & E. MILAKIS, “Quantitative stability for the first Dirichlet eigenvalue in Reifenberg flat domains in $\mathbb{R}^N$ ”, *J. Math. Anal. Appl.* **364** (2010), no. 2, p. 522-533.
- [16] ———, “A stability result for nonlinear Neumann problems in Reifenberg flat domains in $\mathbb{R}^N$ ”, *Publ. Mat., Barc.* **55** (2011), no. 2, p. 413-432.
- [17] A. LEMENANT, E. MILAKIS & L. V. SPINOLO, “Spectral stability estimates for the Dirichlet and Neumann Laplacian in rough domains”, *J. Funct. Anal.* **264** (2013), no. 9, p. 2097-2135.
- [18] A. LEMENANT & Y. SIRE, “Boundary regularity for the Poisson equation in Reifenberg-flat domains”, in *Geometric partial differential equations. Based on the conference, CRM, Pisa, Italy, September 2012*, Centro di Ricerca Matematica Ennio De Giorgi (CRM) Series, vol. 15, Edizioni della Normale, 2013, p. 189-209.

- [19] Y. LIAN, W. XU & K. ZHANG, “Boundary Lipschitz regularity and the Hopf lemma on Reifenberg domains for fully nonlinear elliptic equations”, *Manuscr. Math.* **166** (2021), no. 3-4, p. 343-357.
- [20] Y. LIAN & K. ZHANG, “Boundary Hölder Regularity for Elliptic Equations on Reifenberg Flat Domains”, 2018, <https://arxiv.org/abs/1812.11354v1>.
- [21] S. MAYBORODA & V. MAZ'YA, “Regularity of solutions to the polyharmonic equation in general domains”, *Invent. Math.* **196** (2014), no. 1, p. 1-68.
- [22] ———, “Polyharmonic capacity and Wiener test of higher order”, *Invent. Math.* **211** (2018), no. 2, p. 779-853.
- [23] E. MILAKIS & T. TORO, “Divergence form operators in Reifenberg flat domains”, *Math. Z.* **264** (2010), no. 1, p. 15-41.
- [24] A. PRADE, “Boundary regularity for nonlocal elliptic equations over Reifenberg flat domains”, 2025, <https://arxiv.org/abs/2502.04107>.
- [25] E. R. REIFENBERG, “Solution of the Plateau problem for m -dimensional surfaces of varying topological type”, *Acta Math.* **104** (1960), p. 1-92.
- [26] W. P. ZIEMER, *Weakly differentiable functions. Sobolev spaces and functions of bounded variation*, Graduate Texts in Mathematics, vol. 120, Springer, 1989.