

CHRISTOPH GELLHAUS

TILMANN WURZBACHER

Holomorphic group actions with few compact orbits

Annales de la faculté des sciences de Toulouse 6^e série, tome 10,
n° 2 (2001), p. 293-298

http://www.numdam.org/item?id=AFST_2001_6_10_2_293_0

© Université Paul Sabatier, 2001, tous droits réservés.

L'accès aux archives de la revue « Annales de la faculté des sciences de Toulouse » (<http://picard.ups-tlse.fr/~annales/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Holomorphic group actions with few compact orbits (*)

CHRISTOPH GELLHAUS ⁽¹⁾, TILMANN WURZBACHER ⁽²⁾ *

RÉSUMÉ. — Pour une grande classe de variétés compactes complexes, contenant les images méromorphes des espaces de Kähler compacts, nous montrons le théorème suivant: Soit G un groupe de Lie complexe agissant holomorphiquement sur X tel qu'il y a qu'un nombre fini positif d'orbites compactes. Alors X est un fibré $G \times_I F$, où G/I est le tore d'Albanese de X et F est une fibre de l'application d'Albanese. De plus, F est connexe et son premier nombre de Betti est nul.

ABSTRACT. — For a large class of compact complex manifolds, including for example meromorphic images of compact Kähler spaces, we prove the following theorem: Let G be a complex Lie group acting holomorphically on the manifold X and suppose there is at least one, but only finitely many compact orbits. Then X is a fibre bundle $G \times_I F$, where G/I is the Albanese torus of X and F is a fibre of the Albanese map. Furthermore F is connected and has vanishing first Betti number.

1. Introduction

Holomorphic group actions on a compact complex manifold X with an open set of compact orbits are studied in [GeWu]. There it is proved that X consists only of compact orbits and is itself of a product structure, reflecting the single orbit decomposition of Borel and Remmert, provided there are no

(*) Reçu le 3 novembre 2000, accepté le 9 juillet 2001

(1) Technische Fachhochschule zu Bochum, Fachbereich Elektrotechnik, Herner Str. 45, D-44787 Bochum, Germany.

(2) Laboratoire de Mathématiques, Université de Metz et C.N.R.S., Ile du Saulcy, F-57045 Metz cedex 01, France.

* Corresponding author (email: wurzbacher@poncelet.univ-metz.fr, fax: (33) 03.87.31.52.73)

equivariantly embedded complex tori in the fibres of the Albanese mapping. This condition especially holds true if X is Kähler.

The authors would like to mention that they learned that this result was, in the Kähler case, already proved by D. Snow in 1981 ([Sn]). He used methods, which – in contrast to those of [GeWu] – depend on the fact that the connected automorphism group of a compact Kähler space acts compactifiably (cf. [F], [L]). In the algebraic case parts of this result were obtained by J. Konarski in 1981 ([K]).

Furthermore, Snow showed that the set C of compact orbits of a holomorphic action is analytic, if X is Kähler. In this note we consider the case that C is extremely small, i.e. there are only finitely many compact orbits. For manifolds in Fujiki's class \mathcal{C} (see [F]), which contains for example Kähler and Moishezon manifolds, the following observation can be made:

THEOREM. — *Let X be in \mathcal{C} , $G \times X \rightarrow X$ a holomorphic action of a connected complex Lie group such that there is at least one, but only finitely many compact orbits. Then the following holds:*

X is a fibre bundle $G \times_I F$, where G/I is the Albanese torus of X and F is a fibre of the Albanese map.

Furthermore F is connected and has vanishing first Betti number.

Remark. — Looking at a \mathbb{C}^n -action as a multidimensional generalization of a holomorphic dynamical system, the case that I is discrete, corresponds to the “suspension” of the discrete dynamical system I on F in a continuous system, the \mathbb{C}^n -action on X .

2. Proofs

Let G always denote a connected complex Lie group. If $G \times X \rightarrow X$ is a holomorphic action on a complex space, we refer to X as a holomorphic G -space. In this situation C_G will be the set of compact G -orbits in X .

For smooth X one has the Albanese map $\Psi_X : X \rightarrow \text{Alb } X$ (see [Bl]). The equivariance of ψ_X implies that there is an induced Lie group homomorphism $\lambda_X : \text{Aut}_{\mathcal{O}}(X) \rightarrow \text{Aut}_{\mathcal{O}}(\text{Alb } X)$. The kernel of λ_X restricted to the connected component of the identity in $\text{Aut}_{\mathcal{O}}(X)$ is called $L(X)$, the “linear” automorphisms of X .

Using the compactness of the irreducible components of the space of analytic cycles in complex spaces in class \mathcal{C} resp. Kähler spaces, Fujiki re-

spectively Lieberman prove that $L(X)$ carries the natural structure of a linear algebraic group (Cor. 5.8 in [F], Thm. 3.12 in [L]). For a subgroup H of $L(X)$, \bar{H} will denote the Zariski closure with respect to this structure.

Consider now a smooth holomorphic G -space X . We may assume that G acts without ineffectivity. Let $I = G \cap L(X)$ and \bar{I} the closure of I . Furthermore fix a Levi-Malcev decomposition $\bar{I}^\circ = R_{\bar{I}^\circ} \cdot S_{\bar{I}^\circ}$ of \bar{I}° .

LEMMA 1. — *Let X be a smooth holomorphic G -space in \mathcal{C} and x in X . Assume C_G is not empty. Then the following statements are equivalent:*

- (1) $G(x)$ is compact.
- (2) $I(x)$ is compact.
- (3) $\bar{I}(x)$ is compact.
- (4) $R_{\bar{I}^\circ}$ fixes x and $S_{\bar{I}^\circ}(x)$ is compact.

Proof. — Without loss of generality, we assume throughout this proof that G acts effectively on X .

(1) \iff (2) Since $\text{Aut}_{\mathcal{O}}(\text{Alb } X)^\circ = \text{Alb } X$ the isotropy groups of the induced G -action on $\text{Alb } X$ is equal to I for all points in $\text{Alb } X$. Since C_G is not empty, this implies that all G -orbits on $\text{Alb } X$ are compact. Thus, $G(x)$ is compact iff $I(x)$ is compact.

(2) \implies (3) The group of all g in $L(X)$ which stabilize the compact analytic set $I(x)$ is a Zariski closed subgroup of $L(X)$ (Lemma 2.4 in [F], Prop. 3.4 in [L]). Hence \bar{I} stabilizes $I(x)$ and consequently $\bar{I}(x) = I(x)$.

(3) \implies (4) Since the radical $R_{\bar{I}^\circ}$ is a connected solvable subgroup of $L(X)$ the Borel Fixed Point Theorem for class \mathcal{C} (cf. the proof of Prop. 6.9 in [F] resp. [So] in the Kähler case) shows that it has a fixed point on each component of $\bar{I}(x)$. Since $R_{\bar{I}^\circ}$ is normal in \bar{I} , it acts trivially on $\bar{I}(x)$. Therefore $S_{\bar{I}^\circ}$ acts transitively on the \bar{I} -components.

(4) \implies (2) Since $G/I = \lambda_X(G)$ is a subtorus of $\text{Alb } X$, I/I° is an abelian discrete group. Thus the commutator group $I' \subset I^\circ$ and $I'' \subset (I^\circ)'$. By a result of Chevalley (Thm. 13 and Thm. 15 of paragraph 14, Chapter II in [Ch]) $(I^\circ)' = (\bar{I}^\circ)'$ and this group is Zariski closed in \bar{I} .

It follows that the Zariski closure of I'' is contained in I . Since I is Zariski dense in \bar{I} , the same holds true for I'' in $(\bar{I})''$. Therefore $(\bar{I})'' \subset I$ and a fortiori $S_{\bar{I}^\circ} \subset I^\circ$. This fact shows that $I^\circ(x)$ equals $S_{\bar{I}^\circ}(x)$. Since \bar{I}

has only finitely many connected components, the assumptions imply that $\bar{I}(x)$ is compact. Obviously the same holds now for $I(x)$.

The proof of $((2) \implies (3))$ shows that indeed $I(x) = \bar{I}(x)$. \square

COROLLARY 1 (Snow). — C_G is analytic.

Proof. — If the set C_G of compact G -orbits is not empty Lemma 1 shows $C_G = C_{S_{\bar{I}^\circ}} \cap (\text{Fix } R_{\bar{I}^\circ})$. Since a fixed point set is obviously analytic it suffices to consider connected semisimple groups S .

A compact S -orbit in a complex space X in class \mathcal{C} is a homogeneous-rational manifold S/P (Thm. on p. 255 in [F] respectively [BoRe] in the Kähler case). Thus an arbitrary, but fixed Borel subgroup B of S has non-empty fixed point set on each compact orbit.

It follows that the map

$$\phi : S \times (\text{Fix } B) \rightarrow X, \quad \phi(s, x) := s \cdot x$$

has image C_S . Factorizing ϕ via

$$\bar{\phi} : S/B \times (\text{Fix } B) \rightarrow X, \quad \bar{\phi}(sB, x) := s \cdot x$$

we realize C_S as the image of an analytic space under a proper holomorphic map. The proper mapping theorem of Remmert (see e.g. [CAS]) yields the analyticity of C_S . \square

COROLLARY 2. — If C_G is not empty and A a closed G -invariant analytic set, then C_G intersects A in a non-empty set.

Proof. — By Lemma 1 it is enough to show that \bar{I} , which stabilizes A , has a compact orbit on A . This follows from the fact that for all x in X , $\bar{I}(x)$ is a constructible set with respect to the analytic Zariski topology of X since \bar{I} acts meromorphically/compactifiably (Lemma 2.4 in [F], Remark 3.7 in [L]). \square

Remark. — Lieberman states that $G \cdot A$ is Zariski open in its Zariski closure if G acts compactifiably and A is analytic. Obviously this is wrong even in the algebraic case. What is meant is that $G \cdot A$ contains a Zariski open subset of its (analytic) Zariski closure in X .

Proof of the Theorem. — Without loss of generality we assume that G acts effectively on X .

Step 1. Fibre bundle structure of the Albanese map.

By assumption C_G is not empty. Thus the induced G -action on $\text{Alb } X$ has only compact orbits, all of them isomorphic to G/I , where $I = G \cap L(x)$. Applying Cor. 2 to the analytic sets

$$\Psi_X^{-1}(\lambda_X(G)(\Psi_X(x))) = G \cdot \Psi_X^{-1}(\{\Psi_X(x)\})$$

it follows that G has at least as many compact orbits in X as in $\Psi_X(X)$. Since $\Psi_X(X)$ is connected the assumptions of the theorem imply that $\Psi_X(X)$ is only one G -orbit, which is a subtorus of $\text{Alb } X$. Universality of the Albanese torus implies that Ψ_X is surjective.

Denoting $\Psi_X^{-1}(\{0\})$ by F it is easily checked that the map

$$G \times_I F \rightarrow X, \quad [g, f] \mapsto g \cdot f$$

is G -equivariant and biholomorphic.

Step 2. The topology of the Ψ_X -fibre F .

Stein factorization of Ψ_X together with the universality of Ψ_X yields the connectivity of the Ψ_X -fibres.

Since I stabilizes F the same follows for the Zariski closure \bar{I} in $L(X)$. By Lemma 1 an \bar{I} -orbit in F is closed iff the resp. I -orbit is closed iff the resp. G -orbit is closed. Thus we have only finitely many compact \bar{I} -orbits in F . Since \bar{I}/\bar{I}° is finite the same holds true for \bar{I}° . Applying Step 1 to the action $\bar{I}^\circ \times F \rightarrow F$, $\text{Alb } F$ turns out to be \bar{I}° -homogeneous.

In [GeWu] it is shown that the Borel Fixed Point Theorem implies that the restriction morphism

$$\text{Stab}_F \text{Aut}_{\mathcal{O}}(X) \rightarrow \text{Aut}_{\mathcal{O}}(F)$$

maps $L(X)^\circ$ into $L(F)$ (for F the Albanese fibre as above).

Thus $\text{Alb } F$ is homogeneous under a subgroup of $L(F)$, which clearly says that $\text{Alb } F$ reduces to a point. By the equality $\frac{1}{2}b_1(Y) = \dim_{\mathbb{C}}(\text{Alb } Y)$ for Y in class \mathcal{C} (Cor. 1.7 in [F]) the first Betti number of F must vanish. \square

Acknowledgements. — We would like to thank J. Winkelmann for interesting discussions.

Bibliography

- [Bl] BLANCHARD (A.). — Sur les variétés analytiques complexes, *Ann. Sci. école norm. super.* 73 (1956), 157–202.
- [BoRe] BOREL (A.), REMMERT (R.). — Über kompakte homogene Kählersche Mannigfaltigkeiten, *Math. Ann.* 145 (1962), 429–439.
- [CAS] GRAUERT (H.), REMMERT (R.). — *Coherent Analytic Sheaves*, Springer, Berlin Heidelberg New York Tokyo, 1984.
- [Ch] CHEVALLEY (C.). — *Théorie des groupes de Lie II. Groupes algébriques*, Hermann, Paris, 1968.
- [F] FUJIKI (A.). — On automorphism groups of compact Kähler manifolds, *Invent. math.* 44 (1978), 225–258.
- [GeWu] GELLHAUS (C.), WURZBACHER (T.). — Holomorphic group actions with many compact orbits, *Comment. Math. Helvetici* 64 (1989), 639–649.
- [K] KONARSKI (J.). — Properties of projective orbits of affine algebraic groups, *Lect. Notes Math.*, Vol. 956, 79–91, Springer, 1982.
- [L] LIEBERMAN (D.). — Compactness of the Chow Scheme: Applications to automorphisms and deformations of Kähler manifolds, *Lect. Notes Math.*, Vol. 670, 140–186, Springer, 1978.
- [Sn] SNOW (D.). — Transformation groups of compact Kähler spaces, *Archiv der Math.* 37 (1981), 364–371.
- [So] SOMMESE (A.J.). — Extensions Theorems for reductive group actions on compact Kähler manifolds, *Math. Ann.* 218 (1975), 107–116.